

Phytobiotic Properties of Garlic, Red Ginger, Turmeric and Kencur in Growing Duck

by Diana Indrasanti

Submission date: 18-Sep-2022 01:34PM (UTC+0700)

Submission ID: 1902339629

File name: 6.pdf (213.09K)

Word count: 3455

Character count: 19202

Phytobiotic Properties of Garlic, Red Ginger, Turmeric and Kencur in Growing Ducks

Ismoyowati^{1)*}, Diana Indrasanti²⁾, Mochamad Mufti¹⁾ and Abdoreza Soleimani Farjam³⁾

¹⁾Poultry Production Laboratory, ²⁾Animal Health Laboratory, Faculty of Animal Science, Jenderal Soeman University, Jl. Dr. Soeparno 60, Purwokerto 53123, Central Java, Indonesia

³⁾Institute of Tropical Agriculture, Universiti Putra Malaysia, Malaysia

*Corresponding author email: moy.moyowati@gmail.com

Abstract. Phytobiotic properties of garlic (*Allium sativum*), turmeric (*Curcuma domestica*), red ginger (*Zingiber officinale*) and kencur (*Kaempferia galangal*) were studied using standard *in vitro* antibacterial test and *in vivo* feeding trial with ducklings. In the *in vitro* experiment, potency of aqueous extract of these phytobiotic agents were tested against *Salmonella pullorum* and *Escherichia coli*. Feeding trial was carried out for 6 week starting at day 28 using ducklings fed diets supplemented with 1% of each of four phytobiotic agents. The highest antibacterial activity against *S. pullorum* and *E. coli* was observed with garlic and no additive effect when mixture of phytobiotics was used. Weight gain, feed intake and feed conversion ratio of ducklings were not affected by inclusion of garlic, red ginger and kencur. However, 1% turmeric supplementation significantly reduced growth performance to ducklings.

Key words: phytobiotic, antibiotic, duck, medicinal plants

Abstrak. Penelitian karakteristik fitobiotik dari bawang putih (*Allium sativum*), kunyit (*Curcuma domestica*), jahe merah (*Zingiber officinale*) dan kencur (*Kaempferia galangal*) telah dilakukan secara *in vitro* melalui uji aktivitas antibakteri dan secara *in vivo* dengan perlakuan suplementasi fitobiotik didalam pakan anak itik. Pada percobaan *in vitro*, potensi aktivitas antibakteri dari ekstrak fitobiotik diuji menggunakan *Salmonella pullorum* and *Escherichia coli*. Percobaan suplementasi fitobiotik diberikan masing-masing sebesar 1% didalam pakan anak itik. Pemberian pakan perlakuan dilakukan selama 6 minggu, dimulai pada saat anak itik berumur 28 hari. Hasil penelitian menunjukkan aktivitas antibakteri terhadap *S. pullorum* dan *E. coli* paling tinggi adalah ekstrak bawang putih dan tidak ada pengaruh yang lebih baik apabila dicampur dengan ekstrak fitobiotik lainnya. Pertambahan bobot badan, konsumsi pakan dan konversi pakan anak itik tidak dipengaruhi oleh penambahan bawang putih, kunyit, jahe merah dan kencur. Akan tetapi, suplementasi kunyit nyata menurunkan performan pertumbuhan anak itik.

Kata kunci: fitobiotik, antibiotik, itik, tanaman obat-obatan.

Introduction

Duck producers are still largely using traditional approach of growing and feeding with minimum profit margin and low production efficiency in Indonesia. Many breeders use antibiotics for prevention and treatment of disease as well as growth promoting purposes. Antibiotics has long been under the spotlight as food and feed safety issue in scientific community because of antibiotic residues and resistance development. Therefore, duck producers are increasingly

enforced to adopt application of other antibiotics replacement strategies (Conway and Wang, 2000). Utilization of phytobiotics as locally available natural growth promoters (NGPs) considered as a practical and achievable way (Vidanarachchi et al., 2010; Grashorn, 2010; Yang et al., 2015). Among many medicinal plant available as phytobiotics, garlic, turmeric, red ginger and kaempferia are widely available and used as food item in South and South-East Asia.

Garlic (*Allium sativum*) contains alliin that convert to allicin upon crushing the bulbs. Allicin is reported to have strong antibacterial effect against *Campylobacter jejuni* and improves performance in broiler chickens (Tollba and Hassan, 2003; Royban et al., 2013). Turmeric (*Curcuma domestica*) contains active substances including terpenoids, alkaloids, flavonoids, essential oils, phenols and curcuminoid which often serve as antimicrobial agents against *Escherichia coli* and *Staphylococcus aureus* (Niamsa and Sittiwet, 2009). Kencur (*Kaempferia galanga*) is reported to have antimicrobial effect on gram-positive and negative bacteria (Kochuthressia, et al., 2012; Tewtrakul et al., 2005), but is not used in poultry feeding studies. Red ginger (*Zingiberofficinale*) contain phytochemicals such as zingiberol, zingiberine, bisabolene and gingerol that are potent antimicrobial agents against *Salmonella enteritidis* and of *E. coli* (Indu et al., 2006; Akintobi et al., 2013).

To date, no study has been reported on the use of garlic, red ginger, turmeric and kencur as phytobiotic agent in duck production. Accordingly, this study was aimed to examine the antibacterial potency of garlic, red ginger, turmeric and kencur aqueous extract against *Salmonella pullorum* and *E. coli* and their effectiveness to improve growth performance in growing duck.

Materials and Methods

Fresh bulbs of *Allium sativum* and rhizomes of *Curcuma domestica*, *Zingiberofficinale* and *Kaempferia galangal* are originated from Indonesia and available in traditional market. The plants were cleaned, sliced and washed with sterile distilled water and then oven-dried and milled.

In vitro experiment

Appropriate amount of each plant powder (0.5, 1, 1.5 or 2 g) was placed in a flask, and

added with 100 ml distilled water, the flasks were stirred and the extracts were obtained after filtration using filter paper. Filter paper discs of 6 mm diameter were prepared, sterilized and soaked in each of the prepared filtrates for 10 min. The saturated discs were aseptically placed over *Salmonella Shigella* Agar (SS) plates seeded with *S. pullorum* (Gram-positive) and Eosin Methylene Blue Agar (EMB) plates seeded with *E. coli* (Gram-negative). All steps were conducted under strict laboratory procedures for microbiology. The plates were incubated at 37°C for 24 h. The diameter of inhibition zones was measured in mm using caliper. The tested microorganisms were also assayed for their sensitivity against the antibiotics ampicillin (0.5g/500µl/disk) and chloramphenicol (0.5g/500µl/disk) using standard Kirby Baur disc diffusion method. The 8 treatment groups that tested were garlic 0.5-2% (GA); red ginger 0.5-2% (RG); turmeric 0.5-2% (TU); kencur 0.5-2% (KA); garlic + red ginger 0.5-2% (GR); garlic + turmeric 0.5-2% (GT); garlic + kencur 0.5-2% (GK).

In vivo experiment

This experiment was conducted to determine the effect of phytobiotic supplementation on duck growth performance. A total of 128 day-old male duck (*Anas platyrhynchos*) with 37.5g average weight were obtained from a local hatchery and distributed randomly in 1m²-sized slat floor pens (4 birds/pen) in a conventional open-sided house with 27-29°C housing temperature and 60-80% relative humidity. The birds were fed standard diet with broiler feed (BR1 code Ex), product of PT Cargill Indonesia from DOD to 3 weeks old. The feed contained metabolizable energy of 3000 kcal/ kg, 20.5% crude protein, 5% crude fat, 5.75% crude fiber, 0.95% Ca and 0.8% P. Preliminary basal feed was given for 1 week then fed treatments from 4-10 weeks later. The basal diet composition and nutrient content are presented on Table 1. Each experiment units

27
Table 1. Composition and nutrient content of basal diet (%)

Ingredient	The percentage of feed material
Corn	40
Rice bran	39
Soybean meal	12
Fish meal	8
Vitamin-mineral premix	1
Total	100
<u>Calculated analysis</u>	
Crude protein (%)	16.10
ME (kcal/kg)	2905
Crude fiber (%)	3.91
Crude fat (%)	4.36
Calcium(%)	1.82
Total Phospor (%)	1.33

9 consisted of 4 ducklings each assigned to one of eight dietary treatments with 4 replicates, or 32 pens in total. The dietary treatments were BD: basal feed (control); GA: basal feed + garlic 1%; RG: basal feed + red ginger 1%; TU: basal feed + turmeric 1%; KA: basal feed + kencur 1%; GARG: basal feed + garlic 0.5% + red ginger 0.5%; GATU: basal feed + garlic 0.5% + turmeric 0.5%; GAKA: basal feed + garlic 0.5% + kencur 0.5%. No antimicrobial, anticoccidial drugs or feed additives were included in the basal diets. Ducklings were weighed individually on 4th week until 10th week. Feed intake was recorded and feed conversion ratios (FCR) were calculated. Mortality was recorded in each subgroup. Data were subject to 9 completely randomized design (CRD) using ANOVA general linear models procedure (GLM) of SAS software. Significant effects were separated by Honest Significant Difference (HSD). Mortality data were subject to chi-square analysis 28 and statistical significance was considered at $P \leq 0.05$ test.

Results and Discussion

The results of in vitro experiment show that among the four herbal plants studied the aqueous extract of garlic showed considerable

growth-inhibiting activity against *E. coli* and *S. pullorum* with all concentration (0.5-2%) (Figure 1). Increase in the inhibitory zone was observed by increasing the concentration of the extract. Kencur and particularly turmeric and red ginger were appeared to have almost no to negligible antibacterial effect on *S. pullorum*. However, they were observed to inhibit *E. coli* at least with concentration higher than 0.5%. Combination of garlic extract with three other extracts demonstrated no superior beneficial effect compared to the sole garlic extract.

In vivo experiment result showed the effects of feeding diets supplemented with dried powder form of the four phytobiotics and the combinations on 10 feed intake, body weight (BW), weight gain (WG), feed conversion ratio (FCR) and mortality in Table 2. Following 6 39 weeks of feeding trial, no significant positive or negative effects were observed by supplementation of garlic, red ginger, kaempferia, garlic+red ginger, garlic+turmeric and garlic + kencur on all performance parameters measured. Supplementation of diet with 1% turmeric had detrimental effects on WG, feed intake and FCR compare to control birds. Mortality rate was not affected by any of the 3 dietary treatments.

The results of different studies provide evidence that some medicinal plants might indeed be potential sources of new antibacterial agents even against some antibiotic-resistant strains (Indu et al., 2006). The findings of the current study are consistent with those of Safithria et al. (2011) who found garlic extract inhibited the growth of *E. coli*. Other studies also showed that phytochemicals in garlic have strong antibacterial properties to combat 25 *E. coli*, *Salmonella*, (Johnson and Vaughn, 1969), *C. botulinum* (DeWit et al., 1979), and other pathogenic species. Garlic (*A. sativum*) has a phytochemical content of alkaloids, tannins, phylobatanin, anthraquinone and saponin (Akintobi et al., 2013). Allicin is




Figure 1. Mean inhibition zones of various phytobiotics aqueous extracts against *E. Coli* (A) and *Salmonella* (B) in comparison with standard antibiotics.

PC: positive control (AquaDest); NC: negative control (0.5g/disk Chloramphenicol for *Salmonella* and 0.5g/disk Ampicillin for *E. Coli*); GA: garlic 0.5-2%; RG: red ginger 0.5-2%; TU: turmeric 0.5-2%; KA: kaempferia 0.5-2%; GR: garlic + red ginger 0.5-2%; GT: garlic + turmeric 0.5-2%; GK: garlic + kaempferia 0.5-2%. * Mean±SEM differ versus PC (P<0.05).

Table 2. Growth performance and mortality of duck fed with diets supplemented with various phytobiotics from 4-10 weeks of age

Treatments	BW [§] (g, Wk 4)	BW (g, Wk 10)	WG (g)	Feed intake (g)	FCR	Mortality (%)
BD†	555	1254	699 ^b	2099 ^a	3.03 ^a	12.5
GA	677	1348	671 ^b	2120 ^a	3.17 ^a	0
RG	641	1293	652 ^b	2139 ^a	3.29 ^a	6.25
TU	577	1136	558 ^a	2255 ^b	4.06 ^b	0
KA	587	1222	635 ^b	2158 ^a	3.42 ^a	6.25
GARG	581	1233	653 ^b	2141 ^a	3.31 ^a	0
GATU	540	1269	728 ^b	2075 ^a	2.87 ^a	0
GAKA	557	1217	660 ^b	2132 ^a	3.24 ^a	6.25
SEM	11	15	11	12	0.08	1.64
21 OVA	NS*	NS	*	*	*	NS [#]

[§] BW: body weight; WG: weight gain; FCR: feed conversion ratio.

[†] BD: basal feed (control); GA: basal feed + garlic 1%; RG: basal feed + red ginger 1%; TU: basal feed + turmeric 1%; KA: basal feed + kaempferia 1%; GARG: basal feed + garlic 0.5% + red ginger 0.5%; GATU: basal feed + garlic 0.5% + turmeric 0.5%; GAKA: basal feed + garlic 0.5% + kaempferia 0.5%.

⁷ S: not significant; *: P < 0.05; **: P < 0.01.

^{a,b} Means within a column with no common letters differ at P < 0.05.

[#] NS: not significant with chi square test

the most potentially active component of garlic that is responsible for its characteristic odor, flavor as well as most of its biological properties (Chowdhury et al., 2002). There was no antibacterial activity in extracts of turmeric and low concentration of red ginger and kencur against *S. pullorum*. On the contrary, earlier work showed that the antibacterial activity of *Zingiber officinale*, *Curcuma longa* and *Kaempferia galanga* against Gram-positive and Gram-negative bacteria (Tewtrakul et al., 2005; Niamsa and Sittiwet, 2009; Kochuthressia, et al., 2012; Akintobi et al., 2013). The discrepancies could be associated with the different dosage used, extraction method, the method of antibacterial study, the genetic variation of plant, age of the plant or the environment. It is possible that some of the effective antibacterial active ingredients of these phytobiotics source is not extracted using aqueous extraction method. Cowan (1999) contributed a huge proportion of the plant antibacterial properties to lipid soluble or nonpolar ingredients. It has been reported that plant extracts in organic solvent provided more consistent antimicrobial activity compared to those extracted in water (Parekh et al., 2005). Furthermore, according to Longanga Otshudi et al. (2000) some plants have their full therapeutic effect only if collected at certain time or during a certain season.

It is known that lower number of some gut pathogens such as *E. coli* and *Salmonella* may improve animal performance (Kim et al., 2011). Overgrowth of some microorganisms in the intestine has been correlated with mucosal layer and villi impairment and therefore reducing the capacity of nutrient absorption (Pelicano et al., 2005). Thus, microbial activity has the potential to alter gut morphology in either positive or negative way (Rebolé et al., 2010). In our study, however, no beneficial effect of feeding phytobiotics was observed on performance parameters. Furthermore, we

observed significant negative effect of feeding 1% turmeric on feed intake, weight gain and FCR. Similarly, Durrani et al. (2006) showed that broiler fed by 1% turmeric in their diet had lower FCR than control group. This may arise from overproduction of bile with feeding of turmeric at high levels of inclusion. Al-Sultan and Gameel (2004) associated the positive effect of low dosage of turmeric to its known effect in stimulating bile production in the liver. Garlic was fed to animals in the form of crushed cloves, powder, garlic oil, water and alcohol extracts (Staba et al., 2001). Result on garlic derivative propylpropane thiosulphonate against broiler enteropathogens in vivo reported that supplementation of garlic extract 45 to 135 ml/kg beneficially reduced the numbers of pathogenic and potentially pathogenic bacteria in the intestine, and improved the morphological structure of the ileal mucosa and the productive parameters of broiler chickens (Peinado et al. 2012). Liquid garlic extract supplementation at the levels of 1.50 and 2.25 ml kg⁻¹ significantly increased body weight compared to the control group (Brzóśka et al. 2015).

Regarding the lack of positive effect of garlic, red ginger and kaempferia on performance in our study, no clear explanation could be offered. However, a critical review of the previous studies of medical plants in poultry feeding reveal that there are a huge body of reports on both side indicating beneficial effect or lack of effects. This could be attributed to lack of a standardization in effective dosage, extraction method, species and age of the phytobiotic plant sources among these studies.

Conclusion

Garlic has the most effective antibacterial activity against *S. pullorum* and *E. coli* compared to turmeric, red ginger and kaempferia. Combining garlic with these three phytobiotic showed no additive antibacterial

effects. Supplementation of garlic, red ginger and kaempferia had not improved performance of duck. One percent turmeric addition to diet of growing duck was detrimental to their growth performance.

Acknowledgement

The writers would like to extend gratitude to the Director General of Higher Education, Ministry of Education and Culture of Indonesia and Head of Research and Community Service Institution Universitas Jenderal Soedirman on supporting funds for the implementation of the National Strategic research in accordance with the Decree of the Head of LPPM No. Kept: 430/UN23.10/PN.01.00/2014 dated June 5, 2014.

References

- Akintobi OA, CC Onoh, JO Ogele, A Aldowu, OV Ojo, IO Okonko.2013. Antimicrobial activity of *zingiberofficinale* (ginger) extract against some selected pathogenic bacteria. Nature and Science 11(1): 7-15.
- Al-Sultan SI and AA Gameel. 2004.Histopathological changes in the livers of broiler chickens supplemented with turmeric (*Curcuma longa*). International Journal of Poultry Science, 3: 333-336.
- Brzóška F, B Śliwiński ,O Michalik-Rutkowska and JŚliwa. 2015. The effect of garlic (*Allium sativum* L.) on growth performance, mortality rate, meat and blood parameters in broilers. Ann. Anim. Sci., 15 (4): 961-975.
- Chowdhury SR, SD Chowdhury and TK Smith.2002. Effect of dietary garlic on cholesterol metabolism in laying hen. Journal of Poultry Science 7(2): 122-128.
- Conway PL and X Wang. 2000. Specifically targeted probiotics can reduce antibiotics usage in animal production. Asian-Australasian Journal of Animal Science 13 Supp: 358-361.
- Cowan MM. 1999.Plant products as antimicrobial agents. Clinical Microbiology Reviews 12: 564-582.
- De Wit JC, S Notermans, N Gorin and EH Kampelmacher. 1979. Effect of garlic oil or onion oil on toxin production by *Clostridium botulinum* [food poisoning bacteria] in meat slurry. Journal of Food Protection (USA) 42: 222-224.
- Durrani FR, MIsmail, A Sultan, SM Suhail, N Chand and Z Durrani. 2006. Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. Journal of Agricultural and Biological Science, 1: 9-11.
- Grashorn MA. 2010. Use of phytobiotics in broiler nutrition—an alternative to infeed antibiotics?Journal of Animal and Feed Sciences, 19: 338-347.
- Indu MN, AAM Hatha, C Abirosh, U Harsha and G Vivekanan G. 2006. Antimicrobial activity of some of the South-Indian spice against serotypes of *Escherichia coli*, *Salmonella*, *Listeria monocytogenes* and *Aeromonashydrophila*. Brazilian Journal of Microbiology, 37: 153-158.
- Johnson MG and R Vaughn. 1969. Death of *Salmonella typhimurium* and *Escherichia coli* in the presence of freshly reconstituted dehydrated garlic and onion. Applied and Environmental Microbiology, 17: 903-905.
- Kim GB, YM Seo, CH Kim and IK Paik. 2011. Effect of dietary prebiotic supplementation on the performance, intestinal microflora, and immune response of broilers. Poultry Science, 90:75-82.
- Kochuthressia KP, SJ Britto, MO Jaseentha and R Raphael. 2012. In vitro antimicrobial evaluation of *Kaempferia galangal* Linn. rhizome extract. American Journal Biotechnology and Molecular Sciences, 2(1): 1-5.
- LongangaOtshudi A, A Vercruysse and A Foriers. 2000. Contribution to the ethnobotanical, phytochemical and pharmacological studies of traditionally used medicinal plants in the treatment of dysentery and diarrhoea in Lomela area (DRC). Ethnopharmacology, 3: 411-423.
- Niamsa M and C Sittiwet. 2009. Antimicrobial activity of curcuma longa aqueous extract. Journal of Pharmacology and Toxicology, 4: 173-177.
- Parekh J, D Jadeja and S Chanda. 2005. Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. Turkish Journal of Biology, 29: 203-210.
- Pelicano ERL, PA Souza, HBA Souza, DF Figueiredo, MM Boiago, SR Carvalho and VF Bordon. 2005. Intestinal mucosa development in broiler chickens fed natural growth promoters. Brazilian Journal of Poultry Science, 7:221-229.
- Rebolé A, LT Ortiz, ML Rodríguez, C Alzueta, J Treviño and S Velasco.2010. Effects of inulin and enzyme complex, individually or in combination, on growth performance, intestinal microflora, cecal fermentation characteristics, and jejunalhistomorphology in broiler chickens fed a wheat- and barley-based diet. Poultry Science, 89:276-286.

- Robyn J, GRasschaert, D Hermans, F Pasmans and M Heyndrickx. 2013. Is allicin able to reduce campylobacter jejuni colonization in broilers when added to drinking water? Poultry Science, 9 (5): 1408-1418.
- Safithria M, M Bintang and M Poeloengan. 2011. Antibacterial activity of garlic extract against some pathogenic animal bacteria. Media Peternakan, 34: 155-158.
- Staba EJ, L Lash and JE Staba. 2001. A commentary on the effect of garlic extraction and formulation on product composition. J. Nutr. 131: 1118S-1119S.
- Tewtrakul, S, S Yuenyongsawad, S Kumee and L Atsawajaruwan. 2005. Chemical components and biological activities of volatile oil of *kaempferia galanga* Linn. Songklanakarin Journal of Science and Technology. 27(2): 503-507.
- Vidanarachchi J, LL Mikkelsen, I Sims, PA Iji and M Choct. 2010. Phytobiotics: alternatives to antibiotic growth promoters in monogastric animal feeds. Recent Advances in Animal Nutrition in Australia. 15:131-144.
- Yang C, MAK Chowdhury, Y Hou and J Gong. 2015. Phytogenic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. Pathogens, 4: 137-156.

Phytobiotic Properties of Garlic, Red Ginger, Turmeric and Kencur in Growing Duck

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | www.pjbs.org
Internet Source | 1 % |
| 2 | Peinado, M. J., R. Ruiz, A. Echavarri, and L. A. Rubio. "Garlic derivative propyl propane thiosulfonate is effective against broiler enteropathogens in vivo", Poultry Science, 2012.
Publication | 1 % |
| 3 | api.intechopen.com
Internet Source | 1 % |
| 4 | ejbio.imedpub.com
Internet Source | 1 % |
| 5 | www.pharmahealthsciences.net
Internet Source | 1 % |
| 6 | www.scialert.net
Internet Source | 1 % |
| 7 | L. Masouri, S. Salari, M. Sari, S. Tabatabaei, B. Masouri. " Effect of feed supplementation with essential oil on performance and | 1 % |

physiological parameters of broilers fed on wheat- or maize-based diets ", British Poultry Science, 2017

Publication

8

Longanga Otshudi, A.. "Contribution to the ethnobotanical, phytochemical and pharmacological studies of traditionally used medicinal plants in the treatment of dysentery and diarrhoea in Lomela area, Democratic Republic of Congo (DRC)", Journal of Ethnopharmacology, 200008

Publication

1 %

9

discovery.researcher.life

Internet Source

1 %

10

fvtm.stafpu.bu.edu.eg

Internet Source

1 %

11

www.cambridge.org

Internet Source

1 %

12

R.U. Khan, S. Naz, M. Javdani, Z. Nikousefat, M. Selvaggi, V. Tufarelli, V. Laudadio. "The use of Turmeric (*Curcuma longa*) in poultry feed", World's Poultry Science Journal, 2019

Publication

<1 %

13

Abo El-Maaty, M.A. Hayam, M.H. Rabie, A.Y. El-Khateeb. "Response of Heat-Stressed Broiler Chicks to Dietary Supplementation

<1 %

with Some Commercial Herbs", Asian Journal
of Animal and Veterinary Advances, 2014

Publication

14

Y. J. Kim. "Effect of dietary garlic bulb and
husk on the physicochemical properties of
chicken meat", Poultry Science, 02/01/2009

Publication

<1 %

15

baadalsg.inflibnet.ac.in

Internet Source

<1 %

16

portalgaruda.org

Internet Source

<1 %

17

eprints.utm.my

Internet Source

<1 %

18

mspace.lib.umanitoba.ca

Internet Source

<1 %

19

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

20

tojqi.net

Internet Source

<1 %

21

www.sasas.co.za

Internet Source

<1 %

22

D. Nwachukwu Ifeanyi, F. Asawalam Elechi.
"Laboratory evaluation of freshly prepared
juice from garlic (*Allium sativum* L.) Liliaceae
as protectants against the maize weevil,
Sitophilus zeamais (Motsch.) [Coleoptera:

<1 %

Curculionidae]", African Journal of
Biotechnology, 2014

Publication

23

Submitted to University of Sri
Jayewardenepura Nugegoda Sri Lanka

Student Paper

<1 %

24

eprints.undip.ac.id

Internet Source

<1 %

25

nrccamel.res.in

Internet Source

<1 %

26

pericles.pericles-prod.literatumonline.com

Internet Source

<1 %

27

repositorio.una.ac.cr

Internet Source

<1 %

28

www.jett.dormaj.com

Internet Source

<1 %

29

www.sciendo.com

Internet Source

<1 %

30

www.thaiscience.info

Internet Source

<1 %

31

Atefeh Sheikhlar, Goh Yong Meng, Razak
Alimon, Nicholas Romano, Mahdi Ebrahimi. "
Dietary Extract Improved the Resistance of
Sharptooth Catfish to ", Journal of Aquatic
Animal Health, 2017

Publication

<1 %

- | | | |
|----|--|------|
| 32 | Boguslaw I Olkowski. "Riboflavin content in lupine seeds and blood plasma riboflavin status in broilers fed diets containing high levels of lupine seeds", Journal of the Science of Food and Agriculture, 11/2008
Publication | <1 % |
| 33 | journal.ipb.ac.id
Internet Source | <1 % |
| 34 | repo.unand.ac.id
Internet Source | <1 % |
| 35 | seruniessays.blogspot.com
Internet Source | <1 % |
| 36 | U. Gadde, W. H. Kim, S. T. Oh, Hyun S. Lillehoj. "Alternatives to antibiotics for maximizing growth performance and feed efficiency in poultry: a review", Animal Health Research Reviews, 2017
Publication | <1 % |
| 37 | Edible Medicinal and Non-Medicinal Plants, 2016.
Publication | <1 % |
| 38 | Jolly Akullo Oder, Beatrice Kiage, Dorothy Nakimbugwe, John Kinyuru. "Effect of aqueous and organic solvent extraction on in-vitro antimicrobial activity of two varieties of fresh ginger (Zingiber officinale) and garlic (Allium sativum)", Heliyon, 2022 | <1 % |

39

S.-J. LIM. "Supplemental iron and phosphorus increase dietary inclusion of cottonseed and soybean meal in olive flounder (*Paralichthys olivaceus*)", *Aquaculture Nutrition*, 10/2008

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On