

# Determinat factor stroke prevention behavior

*by* Arif Setyo Upoyo

---

**Submission date:** 20-Dec-2021 03:06PM (UTC+0700)

**Submission ID:** 1734122823

**File name:** Determinat\_factor\_stroke\_prevention\_behavior\_1.docx (39.98K)

**Word count:** 2218

**Character count:** 13156

**Determinant Factors Stroke Prevention Behavior Among Hypertension Patient in Indonesia**

**Arif Setyo Upoyo<sup>1\*</sup>, Atyanti Isworo<sup>1</sup>, Yunita Sari, Ph.D<sup>1</sup>, Agis Taufik<sup>1</sup>, Annas Sumeru<sup>1</sup>, Akhyarul Anam<sup>1</sup>**

<sup>1</sup> Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman

Correspondence should be addressed to Arif Setyo Upoyo:

[afkarfadholi@gmail.com](mailto:afkarfadholi@gmail.com)

**Abstract**

Stroke is the leading cause of death in the World and in Indonesia. Hypertension is a major risk of stroke. Modified stroke risk factors are behavioral factors. The purpose of this study is to identify the factors that influence stroke prevention behavior in hypertensive patients. The research method uses cross sectional. Total participants were 461 hypertensive patients who came to Community Health Center. Sampling technique used consecutive sampling. The questionnaire used contains demographics, knowledge, self efficacy, awareness of stroke risk and stroke prevention behaviors which include diet, exercise, smoking and drinking alcohol. Data analysis used chi-square and logistic regression. The results showed that 65.7% of respondents showed poor stroke prevention behavior. Factors related to stroke prevention behavior include self-efficacy ( $p = 0.043$ ; 95%CI = 1.01-2.22; OR = 1.5), perception ( $p = 0.038$ ; 95%CI = 0.45-0.98; OR = 0.663), and awareness of stroke risk ( $p = <0.001$ ; 95%CI = 2.255-6.333; OR = 3.779). The dominant factor influencing stroke prevention behavior in hypertensive patients is awareness of stroke risk.

Keyword: Behavior, stroke prevention, hypertension, self awareness.

## Introduction

<sup>1</sup> Stroke is a major healthcare problem in South, East, and South-East Asia. With a majority of the world's population living in the developing countries of these regions, the the global burden of stroke will have the largest contribution from Asia [1]. The prevalence of stroke in Indonesia increases every year. <sup>13</sup> Based on the 2013 Basic Health Research, the prevalence of stroke was 7 per thousand population, while in 2018 the prevalence of stroke was 10.9 per thousand population [2].

Risk factors for stroke include factors that cannot be modified and factors that can be modified. Factors that cannot be modified include age, sex, race / ethnicity and genetics, while factors that can be modified include hypertension, diabetes, atrial fibrillation, dyslipidemia, diet, physical activity, obesity, metabolic syndrome, alcohol consumption and smoking (Slark & Sharma, 2014; Boehme et al., 2017, Upoyo et al, 2021) [3-5]. Risk factors for stroke that can be modified are influenced by patient behavior

<sup>12</sup> Hypertension is one of the main factors of stroke. Primary hypertension is a major cause of cardiovascular disease morbidity which has a significant influence on the incidence of stroke [6-7]. Goldstein states that the prevalence of stroke in hypertensive patients aged 50 years is 20% with a risk ratio of 4 and the prevalence continues to increase with increasing age [8], whereas according to Ghani et al hypertension sufferers have a risk of 2.87 times getting a stroke [7].

<sup>8</sup> The purpose of this study was to determine the factors that influence stroke prevention behavior in hypertensive patients

## **Methodology**

### **Design**

The research method used cross sectional study. The research was conducted in April - August 2019.

### **Sample**

The subjects of the study were 461 participants of hypertension. Sampling technique used consecutive sampling from 5 public health center in Banyumas Regency, Central of Java. Inclusion criteria of sample were hypertensive patient, > 35 years old and visit to public health center. Exclusion criteria were hypertensive patients with complications and refused as a respondent

### **Instrument**

The questionnaire used contained demographics, knowledge, self efficacy, awareness of stroke risk and stroke prevention behaviors. Knowledge Instrument modified from Hypertension Knowledge Level Scale (HKLS) by Erkok [9]. that divided in three kategorize: low, middle and high. Self efficacy and awareness of stroke risk used numeric score 1-10, high score if above average and low score if below or equal to average. Stroke prevention behaviors which included low salt and cholesterol diet, exercise, smoking and drinking alcohol used likert score 1-4, total score categorized good if the total score is more than average and bad if the total score is below or equal to average.

### **Data Analysis**

Data analysis used chi-square for bivariate analysis and logistic regression for multivariate analysis with SPSS16.

## **Result**

Based on table 1, it is known that the respondents are mostly female (80.9%), over 60 years old (56.8%), still married (71.4%), living together with family (49.9%), with income below the regional minimum wage (65.7%) and low education (70.9%). Respondent characteristics that significantly influence behavior include gender ( $p = 0.000$ ; 95%CI = 7.12-22.28; OR = 12.59), age ( $p = 0.000$ ; 95%CI = 0.24-0.56; OR = 0.369), marital status ( $p = 0.008$ ; 95%CI = 1.17-2.89; OR = 1.83), and income ( $p = 0.008$ ; 95%CI = 0.39-0.87; OR = 0.58). Female, old age, married and low income have a tendency to behave badly in stroke prevention behavior.

Based on table 2 it is known that the majority showed poor stroke prevention behavior (65.7%), high knowledge of hypertension (89.4%), low self efficacy (56.2%), good perception (57.3%), information about low stroke (41.4%) and low stroke risk awareness (60.1%).

Factors related to stroke prevention behavior include self-efficacy ( $p = 0.043$ ; 95%CI = 1.01-2.22; OR = 1.50), perception ( $p = 0.038$ ; 95%CI = 0.45-0.98; OR = 0.663) and awareness of stroke risk ( $p = 0.000$ ; 95%CI = 0.24-0.53; OR = 0.354).

The results of multivariate analysis showed that factors related to stroke prevention behavior were age, sex, efficacy and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk with OR 3,779.

## **Discussion**

Most respondents are female, elderly, have low education and earn below the regional minimum wage standard. The risk of hypertension increases with age.

This is related to vascular changes that cause an increased peripheral resistance. Postmenopausal women increases the risk of developing hypertension due to hormonal changes. Low education and income can affect health behaviour.

The result shows the majority (65.7%) of hypertension patients had a poor stroke prevention behaviour. The poor behaviour are high consumption of sodium and fat and also lack of regular exercise. The factors that influence stroke prevention behaviour in this study include perception, <sup>16</sup> self-efficacy, and stroke risk awareness.

The majority of the respondents have poor perception and low self-efficacy in stroke prevention. Perception and self-efficacy can influence behaviour. Imprecise perception in stroke prevention and low self-efficacy will inhibit stroke prevention behaviour [10].

One of the factors that influence perception is information. In this study, we obtained bad perception and information about stroke (41.4%). Plenty of information can affect or increase someone's knowledge and with knowledge creates awareness that eventually someone will behave in accordance with the knowledge they have.

<sup>3</sup> Self-efficacy is an important precondition for successful self-management. Many studies have shown <sup>3</sup> that self-efficacy is the strongest determinant of smoking cessation, increasing physical activity, and healthy diet [11-13]. Thus, increasing self-efficacy is an effective method to support healthy behavioural change in patients with stroke.

Respondents also indicated that the majority has a low stroke risk awareness. Many patients are not aware that hypertension is a risk of stroke so they ignore

stroke prevention behaviour. This is <sup>7</sup> in line with the research of Hetz et al (2019) <sup>15</sup> which stated that 7.6% had the chance to become stroke, 46.4% had no chance to become a stroke, and 46.0 were unsure [14]. These findings indicate that self-awareness about stroke risk is poor. This disease attitude can inhibit efforts to raise awareness about stroke in society.

Most people think that stroke is <sup>2</sup> more serious than heart attack. However, because stroke symptoms appear in variety of ways, they are not easily recognized. People with stroke report that they're confused by the symptoms such as migraine-like, headache, ear disease or food poisoning. This kind of confusion can be one of the causes of delays to the hospital. Studies suggest that myocardial infarction patients with atypical symptoms lead to increased delay <sup>2</sup> in many patients [15-16]. Dracup et al found the people who thought they were simply having gas discomfort or indigestion significantly delayed seeing a doctor [17]. This study also revealed that people who had a low stroke self-awareness had a 3.78 times risk of having poor stroke prevention behaviour. Therefore, increasing public awareness about the risk of stroke is an important factor in the stroke risk patients management.

## Conclusion

Most respondents stroke prevention behaviors are bad. Respondent characteristics that significantly influence behavior include gender, age, marital status, and income. Factors related to stroke prevention behavior include self-efficacy, perception and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk. It is necessary to intervene to increase stroke

prevention behavior in risk groups by increasing self-efficacy, perception and awareness of stroke risk.

### **Acknowledgments**

Thanks <sup>5</sup> to the Research and Education Institute of Universitas Jenderal Soedirman Indonesia for funding this research through institution research scheme.

### **Ethical aspect and Declaration of interest statement**

The study has been declared ethical by the Dr. Moewardi hospital's health research ethics commission number: 789 / VI / HREC / 2019. Informed consent was given to the participants before study.

<sup>14</sup> The authors declare there is no conflict interest in this research.

### **Author contributions**

Concept and study design (Upoyo, Sari, Isworo), data collection, data analysis and interpretations (Taufik, Sumeru, Anam), processing the draft of the manuscript (Upoyo, Isworo), critical revision of the manuscript (Sari, Upoyo, Isworo), article finalization (Upoyo, Isworo).

### **References**

- [1] Venketasubramanian, N., Yoon, B.W., Pandian, J., Navarrod, J.C. Stroke Epidemiology in South, East, and South-East Asia: A Review. Journal of Stroke. 2017;19(3):286-294 <https://doi.org/10.5853/jos.2017.00234>



- [2]Ministry of Health of the Republic of Indonesia. (2019, April 10). The reports of the results of basic health research. <https://www.litbang.kemkes.go.id/laporan-riiset-kesehatan-dasar-risikesdas/>
- [3]Slark, J., Sharma, P. Risk awareness in secondary stroke prevention: a review of the literature. *Journal of the royal society of medicine cardiovascular disease*.2014;Feb 2014: 1-6. doi: [10.1177/2048004013514737](https://doi.org/10.1177/2048004013514737)
- [4]Boehme, A.K., Esenwa, C., Elkind, M.S.V. Stroke risk factors, genetics and prevention. *Circulation Research*. 2017;120(3):472-495. doi: [10.1161/CIRCRESAHA.116.308398](https://doi.org/10.1161/CIRCRESAHA.116.308398)
- [5]Upoyo, .AS., Setyopranoto, I., Pangastuti, HS. The Modifiable Risk Factors of Uncontrolled Hypertension in Stroke: A Systematic Review and Meta-Analysis. *Stroke research and treatment*. 2021;Volume 2021 <https://doi.org/10.1155/2021/6683256>
- [6]Hanchaiphiboolkul, S., Pongvarin, N., Nidhinandana, S., Suwanwela, N.C., Puthkhao, P., Towanabut, S., Tantirittisak, T., Suwantamee, J., Samsen, M. Prevalence of stroke and stroke risk factors in thailand: Thai epidemiologic stroke (TES) study. *Journal of Medical Association Thailand*. 2011;94(4):427-36. PMID: 21591527
- [7]Ghani, L., Mihardja, L.K., Delima. Dominant risk factors of stroke in Indonesia. *Buletin Penelitian Kesehatan*. 2016;44(1):49-58. DOI : [10.22435/bpk.v44i1.4949.49-58](https://doi.org/10.22435/bpk.v44i1.4949.49-58)

- [8]Goldstein, L.B., Adams, R., Albert, M.J., Appel, L.J., Brass, L.M., Bushnell, C.D., et al. Primary prevention of ischemic stroke. *Stroke*. 2006; 37:1583–1633. <https://doi.org/10.1161/01.STR.0000223048.70103.F1>
- [9]Erkoc, S.B., Isikli, B., Mentitas, S., Kalyoneu, C. Hypertension knowledge-level scale (HK-LS): a study on development, validity and reliability. *International Journal of Environmental Research and Public Health*. 2012;9:1018-1029. doi: [10.3390/ijerph9031018](https://doi.org/10.3390/ijerph9031018)
- [10]Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing interventions. London: Silverback Publishing; 2014.
- [11]Garcia, K., Mann, T. From 'i wish' to 'i will': social-cognitive predictors of behavioral intentions. *Journal of Health Psychology*. 2003; 8(3), 347–360. DOI: [10.1177/13591053030083005](https://doi.org/10.1177/13591053030083005)
- [12]Sniehotta, F. F., Scholz, U., & Schwarzer, R.. Bridging the intention-behaviour gap: planning, self-efficacy, and action control in the adoption and maintenance of physical exercise. *Psychology & Health*. 2005;20(2), 143–160. <https://doi.org/10.1080/08870440512331317670>
- [13]Sol, B. G., van der Bijl, J. J., Banga, J. D., & Visseren, F. L. Vascular risk management through nurse-led self-management programs. *Journal of Vascular Nursing*. 2005;23(1), 20–24. DOI: [10.1016/j.jvn.2004.12.003](https://doi.org/10.1016/j.jvn.2004.12.003)
- [14]Hertz J.T., Madut D.B., William G., Maro V.P., Crump J.A., Rubach M.P. Perceptions of stroke and associated health-care-seeking behavior in Northern Tanzania: A Community-Based Study. *Neuroepidemiology*. 2019;53: 41-47. <https://doi.org/10.1159/000499069>

- [15] Moser DK, Dracup K. Gender differences in treatment-seeking delay in acute myocardial infarction. *Progress in Cardiovascular Nursing*. 1993;8:6–12. PMID: 8372090
- [16] Schwarz B, Schoberberger R, Rieder A, Kunze M. Factors delaying treatment of acute myocardial infarction. *European Heart Journal*. 1994;15:1595–1598.
- [17] Dracup K, McKinley SM, Moser DK. Australian patients' delay in response to heart attack symptoms. *The Medical Journal of Australia*. 1997;166:233–236.

Table 1 Characteristics of Respondents

Characteristics of Respondents	n	%	OR	95%CI	P
<b>Sex</b>					
Male	88	19.1	12.59	7.12-22.28	< 0.001
Female	373	80.9			
<b>Age</b>					
<60 years old	199	43.2	0.369	0.24-0.56	<0.001
>60 years old	262	56.8			
<b>Marital status</b>					
Married	329	71.4	1.83	1.17-2.89	0.008
Not yet	132	28.6			
<b>Living together</b>					
Living alone	12	2.6	1.27	0.86-1.87	0.227
With couple	230	49.9			
With family	201	43.6			
other	18	3.9			
<b>Income</b>					
Under minimum standart payment	303	65.7	0.58	0.39-0.87	0.008
Standart payment	24	5.2			
Upper minimum standart payment	134	29.1			
<b>Level Education</b>					
No	40	8.7	0.807	0.39-1.63	0.551
elementary	327	70.9			
Yunior high school	57	12.4			
Senior high scholl	34	7.4			
High education	3	0.7			

Table 2. Stroke prevention behavior and associated factors

Variable	n	%	OR	95%CI	P
<b>Stroke prevention behavior</b>					
Good	158	34.3	-	-	-
Bad	303	65.7			
<b>Knowledge</b>					
Low	0	0	0.830	0.44-1.58	0.568
middle	49	10.6			
High	412	89.4			
<b>Self efficacy</b>					
Low	259	56.2	1.50	1.01-2.22	0.043
High	202	43.8			
<b>Perception</b>					
Bad	264	57.3	0.663	0.45-0.98	0.038
Good	197	42.7			
<b>Information about Stroke</b>					
Low	191	41.4	1.19	0.811-1.766	0.366
Middle	126	27.3			
Good	144	31.2			
<b>Stroke Risk Awareness</b>					
Low	277	60.1	0.354	0.24-0.53	<0.001
High	184	39.9			

Table 3. Multivariate Analysis Results (N = 461)

Variabel	S.E.	Sig.	OR	CI95%	
				Min	Max
Sex	0.339	<0.001	0.072	0.037	0.139
age	0.261	<0.001	2.598	1.557	4.335
Marital Status	0.327	0.218	0.669	0.352	1.269
Income	0.260	0.832	0.946	0.569	1.574
Stroke risk awareness	0.263	<0.001	3.779	2.255	6.333
Perception	0.253	0.272	1.321	0.804	2.169
Self efficacy	0.261	<0.001	0.370	0.222	0.617
Living together	0.278	0.893	1.038	0.602	1.792

# Determinat factor stroke prevention behavior

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.j-stroke.org](http://www.j-stroke.org)

Internet Source

2%

2

[www.pubmedcentral.nih.gov](http://www.pubmedcentral.nih.gov)

Internet Source

1%

3

[repub.eur.nl](http://repub.eur.nl)

Internet Source

1%

4

[j-stroke.org](http://j-stroke.org)

Internet Source

1%

5

Arif Setyo Upoyo, Endang Triyanto, Agis Taufik. "Pilot study of a brief hypnotic induction: Effects on blood pressure, heart rate, and subjective distress in patients diagnosed with hypertension", International Journal of Clinical and Experimental Hypnosis, 2021

Publication

1%

6

[pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)

Internet Source

1%

7

[www.mdpi.com](http://www.mdpi.com)

Internet Source

1%

8	<a href="https://journals.lww.com">journals.lww.com</a> Internet Source	<1 %
9	<a href="https://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	<1 %
10	<a href="https://synapse.koreamed.org">synapse.koreamed.org</a> Internet Source	<1 %
11	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet Source	<1 %
12	Davina Biel, Tineke K. Steiger, Nico Bunzeck. "Age-related iron accumulation and demyelination in the basal ganglia are closely related to verbal memory and executive functioning", Scientific Reports, 2021 Publication	<1 %
13	Leandra Binar Ilyasa, Andriyani Asmuni, Nur Romdhona, Ernyasih Ernyasih. "Factors Related to Behavior Handwashing Using Primary-School-Age Children's Soap on Rt's Covid-19 Pandemic. 008/Rw. 007 Bananas, Ciputat Timur in 2021", Muhammadiyah International Public Health and Medicine Proceeding, 2021 Publication	<1 %
14	<a href="https://link.springer.com">link.springer.com</a> Internet Source	<1 %
15	<a href="http://www.karger.com">www.karger.com</a> Internet Source	

<1 %

16

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

17

Anastasia Tjan, I. Gde Raka Widian, Elysanti Dwi Martadiani, I. Made DwijaPutra Ayusta et al. "Carotid artery stiffness measured by strain elastography ultrasound is a stroke risk factor", Clinical Epidemiology and Global Health, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On