

Occurence_and_Nature.pdf

by

Submission date: 08-Mar-2023 08:06AM (UTC+0700)

Submission ID: 2031631637

File name: Occurence_and_Nature.pdf (836.25K)

Word count: 6610

Character count: 36851

BMJ Open Occurrence and nature of questionable research practices in the reporting of messages and conclusions in international scientific Health Services Research publications: a structured assessment of publications authored by researchers in the Netherlands

Reinie G Gerrits,⁸ Tessa Jansen, Joko Mulyanto,⁸ Michael J van den Berg,⁸ Niek S Klazinga, Dionne S Kringos

To cite: Gerrits RG, Jansen T, Mulyanto J, *et al.* Occurrence and nature of questionable research practices in the reporting of messages and conclusions in international scientific Health Services Research publications: a structured assessment of publications authored by researchers in the Netherlands. *BMJ Open* 2019;9:e027903. doi:10.1136/bmjopen-2018-027903

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-027903>).

Received 13 November 2018
Revised 28 February 2019
Accepted 1 March 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.
Public Health, Amsterdam UMC, University of Amsterdam, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands

Correspondence to
Reinie G Gerrits;
r.g.gerrits@amc.uva.nl

ABSTRACT

Objectives Explore the occurrence and nature of questionable research practices (QRPs) in the reporting of messages and conclusions in international scientific Health Services Research (HSR) publications authored by researchers from HSR institutions in the Netherlands.

Design In a joint effort to assure the overall quality of HSR publications in the Netherlands, 13 HSR institutions in the Netherlands participated in this study. Together with these institutions, we constructed and validated an assessment instrument covering 35 possible QRPs in the reporting of messages and conclusions. Two reviewers independently assessed a random sample of 116 HSR articles authored by researchers from these institutions published in international peer-reviewed scientific journals in 2016.

Setting Netherlands, 2016.

Sample 116 international peer-reviewed HSR publications.

Main outcome measures Median number of QRPs per publication, the percentage of publications with observed QRP frequencies, occurrence of specific QRPs and difference in total number of QRPs by methodological approach, type of research and study design.

Results We identified a median of six QRPs per publication out of 35 possible QRPs. QRPs occurred most frequently in the reporting of implications for practice, recommendations for practice, contradictory evidence, study limitations and conclusions based on the results and in the context of the literature. We identified no differences in total number of QRPs in papers based on different methodological approach, type of research or study design.

Conclusions Given the applied nature of HSR, both the severity of the identified QRPs, and the recommendations for policy and practice in HSR publications warrant discussion. We recommend that the HSR field further define and establish its own scientific norms in publication practices to improve scientific reporting and strengthen the impact of HSR. The results of our study can serve as

15

Strengths and limitations of the study

- Given the explorative nature of this study, we applied a broad and sensitive definition of 'questionable research practices' (QRPs) that allows for the identification of QRPs previously overlooked in related assessments.
- This study describes an assessment of publications and is therefore able to detect QRPs that go unnoticed in survey studies that rely on self-report.
- Although we aimed to develop a reliable measurement instrument that would guide the review process, the instrument allowed latitude for the reviewer's interpretation.
- In our assessment method, we relied on consensus among assessors, which inevitably introduces some subjectivity.
- Because publications were selected based on the title, selection bias might have occurred.

an empirical basis for continuous critical reflection on the reporting of messages and conclusions.

42 INTRODUCTION

In 2009, Chalmers and Glasziou estimated that 85% of research funding in biomedical sciences was wasted avoidably,¹ resulting in *The Lancet's* series 'Increasing value: reducing waste'. This series has stirred the international scientific community, prompting funders, regulators, academic institutions and scientific publishers to act. Funders of biomedical research have responded by organising conferences on research waste, and journal editors have initiated discussions on data

sharing and open access.² While evidence for questionable research practices (QRPs) in biomedical sciences is mounting,¹ little is known about the occurrence and nature of QRPs in the policy-oriented and management-oriented field of health services research (HSR). In particular, QRPs in the reporting of messages and conclusions have flown under the radar. The term 'questionable research practices' is commonly used to describe practices such as selective publication of results, concealing of conflicts of interests and describing a hypothesis after finding significant results.³ A questionable practice is not necessarily wrongful but does 'raise questions'. In this study, we further define the meaning of QRPs in the reporting of messages and conclusions in the field of HSR specifically.

The HSR field is an applied field of research, and produces evidence on topics such as copayments, evaluation of quality improvement efforts, cost-effectiveness of medications, patient empowerment, therapy compliance and effects of policies. Given the growing evidence for the prevalence of QRPs in the reporting of messages and conclusions in the biomedical field,^{4,5} QRPs may also occur in the HSR field. In the biomedical field, a systematic review by Chiu *et al* shows that estimates for the occurrence of questionable research practices in the interpretation of results in scientific publications vary from 10% of publications deriving discordant conclusions from study results to 100% of publications containing rhetorical practices resulting in spin, such as failure to compare risk with benefits in randomised controlled trials (RCTs).⁴

Just like biomedical researchers, health services researchers are under pressure to publish in high-impact journals to increase their citation scores and attract media attention to augment their prestige and chances for future research funding and job security.⁶⁻⁹ Unlike biomedical research, HSR findings are not easily generalised from one local or national health services setting to another, and messages and conclusions tend to be limited to a specific national context.¹⁰ A broad spectrum of quantitative and qualitative methods is used in HSR, including designs that are less subject to strict codes of execution than RCTs, such as observational and case study designs. Furthermore, HSR has difficulty creating alignment between the construction of scientific knowledge and the implementation of that knowledge in policy and practice.¹¹ This combination of HSR-specific characteristics may result in a different set of QRPs in the reporting of a scientific study. The variation of designs other than RCTs, as is more common in the biomedical field, might invite unjustified claims of causality. Moreover, the context-specific research may increase unjustified claims of generalisability, and the difficulty in translating knowledge to practice may result in unsupported recommendations or implications.

Although reporting in scientific publications is highly standardised, the discussion and conclusion sections offer researchers relative freedom when deriving messages and

conclusions from study results.⁵ We explored the occurrence and nature of QRPs in the reporting of messages and conclusions in international scientific HSR publications authored by researchers from HSR institutions in the Netherlands. We also examined the relationship between study type, methodology and design and the occurrence of QRPs. With our study, we want to fuel the debate on fostering responsible messages and conclusions, and provide a basis for the discussion on QRPs in the international HSR field.

METHODS

Setting

This study assessed scientific publications authored by researchers from 13 HSR groups, departments or institutions (hereafter referred to as 'HSR institutions') in the Netherlands, including both academic and non-academic institutions. These institutions all agreed to participate in an effort to assure the overall quality of HSR publications in the Netherlands.

Defining QRPs in the reporting of messages and conclusions in HSR

We conducted a literature review on QRPs in the reporting of messages and conclusions in biomedical research and HSR.¹²⁻¹⁴ An initial definition of QRPs in the reporting of messages and conclusions in HSR was proposed and discussed at a consensus meeting with the directors/leaders of the 13 participating institutions. This was then validated through inputs from five leading international health services researchers (10 were invited; 50% non-response), and resulted in the following amended definition: 'To report, either intentionally or unintentionally, conclusions or messages that may lead to incorrect inferences and do not accurately reflect the objectives, the methodology or the results of the study.'

Measurement instrument

We developed an extensive list of QRPs in the reporting of messages and conclusions. Items were based on the EQUATOR checklist¹⁵ and earlier checklists for identifying 'spin' (ie, 'a way to distort science reporting without actually lying')⁵ or other QRPs.^{13 14 16 17} The proposed list of QRPs was reviewed, refined and complemented using 14 semistructured interviews with the directors/leaders and representatives (n=19) of the 13 participating HSR institutions. Next, the five participating international health services researchers provided email feedback on the list resulting from these interviews; the list was adapted accordingly, resulting in 35 possible QRPs in the reporting of messages and conclusions in HSR publications.

We developed a data extraction form in Excel that contained the list of QRPs and bibliometric information, and conducted a pilot to evaluate its feasibility and usability. In the pilot, two assessors (RGG, TJ) independently assessed five international HSR publications

to identify modifications needed to improve the form, and to align the interpretation of the items. The project group discussed the proposed modifications, resulting in the final version. The data extraction form (see online supplementary material 1) and a methodology of the development of the data extraction form (see online supplementary material 2) are provided in the supplementary material.

Sample

We aimed to include 10 HSR publications from each participating HSR institution. Inclusion criteria were: published in 2016 in an international peer-reviewed scientific journal, written in English, reporting HSR findings and first-authored and/or last-authored by researchers affiliated with the respective HSR institution. As both the first author and the research institution are likely important factors influencing the occurrence of QRPs, only unique first authors were included in the publication. Moreover, not more than 10 publications per institution were included. This will ensure a maximum spread of authors and institutions across the sample.

Publication lists of the HSR institutions were retrieved either by searching publicly accessible online sources (eg, annual reports, open repositories or the research groups' website) or obtained from secretaries or librarians. All lists were verified by the respective HSR institutions. These lists included both HSR and non-HSR publications.

Two researchers (RGG, TJ) selected all titles from the 13 publication lists that were likely to indicate empirical or systematic assessment studies in HSR. Publications were included if their title fitted the definitions of HSR by Juttmann and Lohr and Steinwachs.^{18,19} These definitions are commonly used by HSR institutions (eg, in education) in the Netherlands. To select HSR studies, TJ and RGG first individually selected titles from the publication lists. Next, RGG and TJ compared their selections of titles and noted any differences. After completing the selection of the first HSR publications, selection was reviewed and approved by the research group (NSK, DSK, MJB). TJ and RGG then continued applying the selection method to the remaining publication lists. In a consensus meeting between TJ and RGG, differences in selected titles were resolved by discussing its fit with the definition. Consensus was reached on all included publications.

The HSR publications (n=717) were assigned a random number. Per institution, the publications with unique first authors with the lowest assigned number were included in the sample. Three HSR institutions did not have enough publications with unique first authors, resulting in a selection of nine, eight and two publications for these institutions. Furthermore, two publications were excluded during assessment because they concerned research protocols. These publications were replaced by another publication authored by the same institution. One publication was excluded because its methodology was considered incomprehensible by the reviewers. Ultimately, 116 HSR publications were included (16% of tot sample).

Assessment process

Two reviewers independently assessed all publications (RGG and TJ or RGG and JM). RGG has primarily qualitative HSR experience and is trained in health economics. TJ and JM have primarily quantitative HSR experience and are trained in public health, management, economics and law and medicine, respectively.

The assessment started with a test phase. During this phase, agreements and disagreements in assessments of the first 30 publications were thoroughly discussed (by RGG, TJ, NSK and DSK) to increase the accuracy of the assessments; agreement between the two reviewers (TJ, RGG) was 81% for the first 20 publications, which increased to 82% when assessing the next 10 publications. The notion emerged that it was necessary having two reviewers with complementary expertise assess each publication independently, followed by a consensus procedure and random check by the project leaders. RGG trained the third reviewer (JM).

RGG assessed all included publications, while TJ assessed the first 59 publications, and JM the remaining 57. All data were entered in the data extraction form. QRPs were coded as either 1, 'present'; 0, 'not present'; -8, 'not applicable to this study' (primarily used for items not applicable for qualitative research); or -9, 'not assessable'. To justify their assessments, the reviewers recorded their motivation for every identified QRP. At a later stage, QRPs in implications and recommendations for policy and practice were further refined into 'not mentioned' if no implication or recommendation was included in the publications, and 'not sufficiently justified', if the authors did not provide any explanation for their implications or recommendations. The reviewers held regular consensus meetings (after review of 10 publications) to discuss and reach agreement on all identified QRPs.

During the consensus meetings, the reviewers compared their assessment of all items. Inconsistencies between the individually assessed QRPs were identified, discussed and adapted. Any remaining disagreements (n=2) were resolved by a senior researcher (DSK). NSK and DSK each reassessed a random sample of six publications, so 10% of all included publications (n=12). As a result, two identified QRPs were retracted, and two QRPs were added to the reassessed publications.

Analysis

The characteristics of the included publications were described by calculating their occurrence with the percentage or mean number of publications.

We counted the total number of QRPs per publication, and the percentage of HSR publications with number of observed QRPs. The latter was visualised in a histogram. Occurrence of specific QRPs was calculated as a percentage of publications containing this particular QRP. The percentage of publications containing QRPs that were not applicable to qualitative research was calculated only for quantitative and mixed methods-based publications (n=83), (eg, the QRP: 'The relevance of

statistically significant results with small effect size is over-stated' is only applicable to quantitative research).

We used a Kruskal-Wallis test to calculate the difference in total number of QRPs applicable to all research designs by methodological approach (quantitative, qualitative, and mixed) type of research (descriptive, exploratory, hypothesis testing and measurement instruments) and study design (observational, (quasi) experimental, systematic review, economic evaluation, case study and meta-analyses). We used the Strengthening the Reporting of Observational Studies in Epidemiology checklist for observational studies in the reporting of this research.²⁰ Analyses were conducted using SPSS V.24.²¹

38

Patient and public involvement

No patients were involved in this study. This study was designed with the input provided by the participating HSR institutions at a consensus meeting at the onset of the study, and individual interviews with the directors/leaders of the 13 participating institutions. During a progress meeting with the participating institutions, preliminary (aggregated level) results were discussed to validate and complement the interpretation of findings.

Ethics approval

A waiver for ethical approval was obtained for this study. To avoid negative consequences for the authors of the included publications, each publication was assigned a unique identification number. Extracted data were entered in SPSS using this number to separate author information from the study data.

RESULTS

Characteristics of included publications

Table 1 presents the characteristics of the 116 included publications from the 13 participating HSR institutions. To summarise, 54.3% of the publications were quantitative, 28.4% were qualitative and 17.2% applied a mixed methods approach. Sixteen per cent of the publications were based on a published study protocol. The mean impact factor of the journals was 2.81, and the average number of authors was six.

Of the 116 HSR publications, the median number of QRPs per publication was six (IQR, 5.75), out of 35 possible QRPs. The distribution of the observed frequency of QRPs across publications is visualised in figure 1.

Frequency of QRPs per type

For each of the QRPs, we counted how often they were identified in the included publications. Online supplementary material 3 presents the percentage of occurrence per QRP type.

QRPs that occurred most frequently were:

- Implications for policy and practice do not adequately reflect the results in the context of the referenced literature (69.0%)*;

Table 1 Characteristics of included publications


Total (n=116)	n (%)
HSR domain	
Policy	19 (16.4)
Social factors	11 (9.5)
Financing systems	10 (8.6)
Organisational structures and processes	43 (37.1)
Health technologies	11 (9.5)
Personal behaviours	22 (19.0)
Methodological approach	
Quantitative	63 (54.3)
Qualitative	33 (28.4)
Mixed methods	20 (17.2)
Type of research	
Descriptive	31 (26.7)
Exploratory	59 (50.9)
Hypothesis testing	19 (16.4)
Measurement instruments	5 (4.3)
Other	2 (1.7)
Design	
Observational	59 (50.9)
(Quasi) experimental	9 (7.8)
Systematic review	17 (14.7)
Economic evaluation	5 (4.3)
Meta analyses	3 (2.6)
Case study	22 (19.0)
Other	1 (0.9)
Protocol published	19 (16.4)
Funder of study stated	98 (84.5)
Contributions stated	57 (49.1)
Number of included journals	80 (100.0)
Mean	
Impact factor journal (n=93 publications*)	2.81 (SD 1.45)
Number of authors (n=116)	6.12 (SD 5.53)

Occurrence of QRPs per publication.

*Not all journals had an impact factor. Mean impact factor was calculated over 93 publications.

HSR, health services research.

- *In 50.0% of publications, no implications for policy and practice were mentioned, and in 19.0% of publications, implications were mentioned without adequate justification.
- Recommendations for policy and practice do not adequately reflect the results in the context of the referenced literature (65.5%)*;
- **In 34.5% of publications, no recommendations for policy and practice were reported, and in 31.0% of publications, recommendations were mentioned without adequate justification.



13

Figure 1 Percentage of health services research publications with number of observed questionable research practices (QRPs) in the reporting of messages and conclusions.

- ▶ Contradicting evidence is poorly documented (63.8%);
- ▶ Conclusions do not adequately reflect the findings as presented in the results section (46.6%);
- ▶ Possible impact of the limitations on the results is not or poorly discussed (44.0%);
- ▶ Conclusions are not supported by the results as presented in the context of the referenced literature (43.1%).

QRPs that occurred least frequently were:

- ▶ The main source of evidence for supporting the results is based on the same underlying data (2.6%);
- ▶ Generalising findings to populations not included in the original sample is not justified (2.6%);
- ▶ Causative wording is used in the hypothesis/research question, although there is no theory to support causation (2.4%);
- ▶ Possible clinical relevance of statistically non-significant results is not addressed (2.4%);
- ▶ Generalising findings to time periods not included in the original study is not justified (0.0%).

Distribution of QRPs

Figure 2 shows the distribution of QRPs across publications. The horizontal axis shows the publications (n=116) ordered from the publication with the lowest (0) to the highest number (18) of observed QRPs in the reporting of messages and conclusions. The vertical axis shows the QRPs ordered from least (generalisation to different time period) to most (implications for practice are lacking) frequently observed. On the right vertical axis, the occurrence of QRPs is presented in number of QRPs counted. Each dot represents a QRP.

The difference in the number of QRPs by publication characteristics

Table 2 shows the associations between total number of QRPs (applicable to all study designs) and methodological


approach (quantitative, qualitative and mixed), type of research (descriptive, exploratory, hypothesis testing and measurement instruments) and study design (observational, [quasi] experimental, systematic review, economic evaluation, case study and meta-analyses). No statistically significant differences in number of QRPs was found by type of research, methodological approach or study design.

DISCUSSION

We explored the occurrence and nature of QRPs in the reporting of messages and conclusions in international scientific HSR publications authored by researchers from HSR institutions in the Netherlands, and examined the relationship between study type, methodology and design and the occurrence of QRPs. Our results indicate that HSR publications have a median of six QRPs per publication. We identified most QRPs in the reporting of implications for policy and practice, recommendations for policy and practice, contradictory evidence, study limitations and conclusions based on the results and in the context of the literature. No significant associations between number of QRPs and type of study, study design or methodological approach were identified.

Limitations and strengths

We applied a broad and sensitive definition of 'questionable', for instance by considering the absence of contradictory evidence or the absence of implications and recommendations for policy and practice as a QRP. The choice to not present contradictory evidence does not defy current publication checklists, yet this practice may hinder interpretation of findings in the full context of evidence. If authors searched for contradictory evidence, but did not mention its absence, readers of the publication would not have any clues on its existence.



The full QRP name is provided in supplementary material 3, table 1 ordered from least frequently found (Generalisation to different time period) to most frequently found (Implications for practice are lacking) QRP.

Figure 2 Distribution of questionable research practices (QRPs) in the reporting of messages and conclusions across health services research publications, ordered from lowest to highest number of observed QRPs.

Knowledge on the occurrence of QRPs is often derived from survey studies, relying on self-report.³ These studies focus on the knowledge of consciously conducted,

Table 2 Association between total number of questionable research practices and type of research, methodological approach and study design

	Median	95% CI	P value
Methodological approach			0.339
Quantitative	5	4.88 to 6.43	
Qualitative	6	4.98 to 7.62	
Mixed methods	7	5.34 to 8.46	
Type of research			0.295
Descriptive	6	4.77 to 6.78	
Exploratory	7	5.76 to 7.60	
Hypothesis testing	4	3.40 to 6.81	
Measurement instruments	5	2.14 to 6.66	
Other	5	-33.12 to 43.12	
Study design			0.159
Observational	6	5.56 to 7.21	
(Quasi) experimental	3	2.07 to 5.71	
Systematic review	6	4.61 to 8.33	
Economic evaluation	4	1.61 to 7.59	
Case studies	6	4.71 to 8.01	
Meta-analyses	5	0.50 to 10.84	

well-known QRPs. Our assessment approach allowed us to gain insight in less severe, more likely unconsciously occurring QRPs in the reporting of messages and conclusions specifically. The number of QRPs identified through assessment is generally higher than in studies relying on self-report.^{3,4} With our broad definition encompassing 35 possible QRPs, we bring to light the areas that offer possibilities for further enhancing publication practices in HSR. Consequently, this definition allows for a discussion in the field of HSR on the extent to which the identified QRPs are acceptable. This is an important strength of our applied approach.

Although we endeavoured to develop a reliable measurement instrument that would guide the review process, the instrument allowed latitude for the reviewer's interpretation. Consequently, a different group of reviewers might arrive at somewhat different scoring frequencies for observed QRPs. However, because we defined each QRP in detail, it is unlikely that there would be substantial differences in the overall distribution of different types of QRPs across publications. Our consensus method contains a degree of subjectivity, and there is the risk that one reviewer's opinion will dominate. To counteract this, NSK and DSK performed random checks on 10% of all assessments. By recording the motivation for every identified QRP, we supported the consistency of our measurement and justified our results. Because publications were selected based on the title, selection bias might have occurred. Considering we

found no relationship between study characteristics and number of QRPs, it is unlikely that a different sample would have led to different results. Inevitably, reviewers sometimes assessed publications written by authors they knew professionally or personally, and as such, a positive view of a colleague's work might have led to underestimating the QRPs in these publications.

Our study results may be representative for HSR research publications internationally. Given the fact that publication in international journals is highly standardised in terms of language (English) and format, our findings can most likely be transferred to HSR communities in other countries.

Interpretation

In HSR publications, recommendations for policy and practice warrant most attention. A study can be conducted properly, using a sound design and appropriate methodology. However, making recommendations without adequate justification could lead to incorrect inferences in policy and the management of healthcare, and undermine society's confidence in science.^{11 22–25}

Measures for safeguarding scientific soundness like those often used in biomedical research (eg, trial registration, open data policies and an improved reporting and archiving infrastructure²⁶) do not address reporting conclusions not supported by study results, and are not tailored to the observational and explorative designs most prevalent in HSR. Moreover, existing publication checklists address a report's completeness, but do not question the justification of the conclusions.⁵ If we intend to improve the reporting of HSR conclusions and recommendations, we will need to better understand the factors that influence authors when reporting the discussion and conclusions section of an HSR publication, for example, media pressure and relationships with funders.^{6 7 9 27} Journals may have influence on the reporting of a study through control of the review process.²⁸ Moreover, research institutions may prevent the occurrence of QRPs by enhancing internal integrity, training in scientific writing and communication among researchers.²⁹ Consequently, subsequent research can focus on what influences researchers when writing their scientific publications, and what factors play a role in the process from research design to the acceptance of a manuscript by a peer-reviewed journal.

A third of the HSR publications studied gave no recommendations for policy or practice, while another third did not provide an adequate justification for the recommendations. One could argue that HSR is an applied field of research, and that its ultimate goal should be to contribute to better health services and systems; researchers should therefore take responsibility for providing guidance to those who can act on the research findings instead of leaving them empty-handed. On the other hand, health services researchers may feel more comfortable committing to a more traditional interpretation of the role of academics, refraining from normative judgement. If

the latter is the dominant viewpoint, the HSR community needs to consider the role of scientific evidence in helping decision makers address the challenges they face, and informing policies and practices. Internationally, the HSR community has been promoting further strengthening of the link between HSR and practice.³⁰

In biomedical research, research being 'new' might contribute to a confused assessment of implications.³¹ This problem is amplified in HSR, where there is a limited accumulation of evidence. HSR considers a larger range of contextual factors and stakeholders in politics or management. Moreover, HSR recommendations are often based on observational or exploratory research, which is considered to be weak evidence in biomedical circles (eg, the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) checklist).³² Perhaps the norms determined by the biomedical research field make health services researchers hesitant to provide any implications or recommendations at all.

Implications and recommendations for policy and practice

The HSR field currently seems to adhere to the norms and expectations set by the biomedical field, even though HSR is multidisciplinary, and differences in approach and type of methodology pose serious challenges to observing these norms. Therefore, the HSR community needs to further define specific scientific norms appropriate to the field.

Scientific norms are developed through the forum of a scientific community.³³ This forum function is particularly strong in the Netherlands, where a community of HSR institutions work together closely. Our study was able to bring together the main Dutch academic and non-academic HSR institutions. Consequently, the results of our study help to facilitate critical reflection on the current state of research and encourage debate on how to systematically advance the reporting of messages and conclusions in HSR. Such a debate in the Dutch context is needed, given the attempts over the past decade by the Netherlands Organisation for Health Research and Development (ZonMw) to strengthen the link between research and practice. It would also be very timely, considering the ongoing, overarching Dutch research programme on responsible research practices funded by ZonMw, of which this study is a part. We recommend the HSR community to reflect on the questions our results bring forward: how do we include implications and recommendations for policy and practice in scientific publications?; how should we describe conclusions in context of literature with limited accumulation of evidence?; and what is the severity of the identified QRPs? Through this publication, we would like to urge journal editors and those working in the international field of HSR to join in this debate. After establishing norms regarding these frequently occurring QRPs, journal editors and HSR institutions may contribute to the prevention of QRPs by implementing strategies tailored to HSR research specifically.

CONCLUSIONS

QRPs in the reporting of messages and conclusions occur frequently in peer-reviewed international scientific HSR publications from Dutch HSR institutions. These QRPs differ in severity and cannot always be qualified as wrongful, but they do 'raise questions'. To ensure the applicability of HSR research in policy and practice, the HSR field should reflect on scientific norms for the reporting of conclusions and the inclusion of recommendations for policy and practice. Our study can serve as an empirical basis for continuous critical reflection on the current state of research, and encourage debate on how to systematically advance the reporting of messages and conclusions in HSR.

Correction notice This article has been corrected since it first published online. The open access licence type has been amended.

Acknowledgements We thank the Dutch HSR institutions that participated in this study: Erasmus MC, Department of Public Health; Erasmus University, Erasmus School of Health Policy and Management; Leiden University Medical Centre, Department of Medical Decision Making and the Department of Public Health and Primary Care; University Maastricht, Health Services Research group; the Netherlands Institute for Health Services Research (NIVEL); Radboud UMC, IQ healthcare; the National Institute for Public Health and the Environment (RIVM); University of Groningen, Faculty of Economics and Business; Tilburg University, Social and Behavioural Sciences, Tranzo; Trimbos Institute; University Medical Centre Utrecht, Julius Centre for Health Sciences and Primary Care; Amsterdam UMC – Vrije Universiteit Amsterdam, and the Amsterdam UMC – University of Amsterdam. Furthermore, we also would like to thank project advisers Anton Kunst, Thomas Plochg and Patrick Romano. We thank Ellen Nolte, Enrique Bernal-Delgado, Michel Wensing, Oliver Groene and Wilm Quentin for their valuable feedback on the definition of QRPs and the measurement instrument.

Contributors DSK, NSK, MJB, RGG and TJ designed the study. RGG, TJ and JM collected the data. RGG and JM analysed the data, and RGG drafted the manuscript. TJ, JM, MJB, NSK and DSK were all involved in the interpretation of the results and were major contributors to the writing of the manuscript. All authors read and approved the final manuscript and are accountable for all aspects of the work.

Funding This study was funded by ZonMw grant number 445001003.

Disclaimer The funder had no role in the study design, in the collection, analysis, and interpretation of data, in the writing of the manuscript, and in the decision to submit the paper for publication. All authors had full access to the data during the conduct of this study and take responsibility for the integrity of the data and the accuracy of the analysis.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Medical ethics review committee at Amsterdam UMC.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data are available upon request to the corresponding author.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *The Lancet* 2009;374:86–9.
- Moher D, Glasziou P, Chalmers I, et al. Increasing value and reducing waste in biomedical research: who's listening? *Lancet* 2016;387:1573–86.

- Fanelli D. How many scientists fabricate and falsify research? A systematic review and meta-analysis of survey data. *PLoS One* 2016;11:e015738.
- Chiu K, Grundy Q, Bero L. 'Spin' in published biomedical literature: a methodological systematic review. *PLoS Biol* 2017;15:e21173.
- Boutron I, Ravaud P. Misrepresentation and distortion of research in medical literature. *Proc Natl Acad Sci U S A* 2018;115:2613–9.
- Iannidis JP, Greenland S, Hlatky MA, et al. Increasing value and reducing waste in research design, conduct, and analysis. *Lancet* 2014;383:166–75.
- Singart P. Science and the media. *Res Policy* 1998;27:869–79.
- Tijink JK, Schipper K, Bouter LM, et al. How do scientists perceive the current publication culture? A qualitative focus group interview study among Dutch biomedical researchers. *BMJ Open* 2015;5:e008681.
- Bouter LM. Commentary: perverse incentives or rotten apples? *BMJ Open* 2015;5:e008681.
- Porter ME. What is value in health care? *N Engl J Med* 2010;363:2477–81.
- Van TP. Startdocument 'Systeemfalen'. The Hague, 2013.
- Arton R. The rhetoric of research. *BMJ* 1995;310:985–7.
- Boutron I, Dutton S, Ravaud P, et al. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. *JAMA* 2010;303:2058–64.
- Ochodo EA, de Haan MC, Reitsma JB, et al. Overinterpretation and misreporting of diagnostic accuracy studies: evidence of "spin". *Diagnosis* 2013;267:581–8.
- Altman DG, Simera I, Hoey J, et al. EQUATOR: reporting guidelines for health research. *Lancet* 2008;371:1149–50.
- Stevens GA, Alkema L, Black RE, et al. Guidelines for accurate and transparent health estimates reporting: the GATHER statement. *PLoS One* 2016;13:e020056.
- Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- Mann RE. Het domein van gezondheidszorgonderzoek. In: Plochg T, Klazinga NS, Mackenbach JP, eds. *Handboek gezondheidszorgonderzoek*. Houten: Bohn Stafleu van Loghum, 2013;7:22–3.
- Lohr KN, Steinwachs DM. Health services research: an evolving definition of the field. *Health Serv Res* 2002;37:7–9.
- von Elm E, Altman DG, Egger M, et al. [The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies]. *Rev Esp Salud Publica* 2008;82:344–9.
- IBM Corp. *IBM SPSS statistics for windows, version 24.0*. Armonk, NY: IBM Corp, 2016.
- Lavis JN, Ross SE, Hurley JE, et al. Examining the role of health services research in public policymaking. *Milbank Q* 2002;80:125–54.
- Hanney SR, Gonzalez-Block MA, Buxton MJ, et al. The utilisation of health research in policy-making: concepts, examples and methods for assessment. *Health Res Policy Syst* 2003;1:2.
- Kuruvilla S, Mays N, Walt G. Describing the impact of health services and policy research. *J Health Serv Res Policy* 2007;12(Suppl 1):23–31.
- Burstein P. The impact of public opinion on public policy: a review of the agenda. *Polit Res Q* 2003;56:29–40.
- Glasziou P, Altman DG, Bossuyt P, et al. Reducing waste from incomplete or unusable reports of biomedical research. *The Lancet* 2014;383:267–76.
- Pittman P. Health services research in 2020: data and methods for the future. *Health Serv Res* 2010;45(5 Pt 2):1431–41.
- Wager E, Middleton P. Technical editing of research reports in biomedical journals. *Cochrane Database Syst Rev* 2008;4:MR000002.
- Fanelli D, Costas R, Larivière V, et al. Misconduct policies, academic culture and career stage, not gender or pressures to publish, affect scientific integrity. *PLoS One* 2015;10:e0127556.
- HSR-Europe. *Health services research: helping tackle Europe's health care challenges*. Policy Brief. Utrecht: NIVEL, Technische Universiteit Berlin, National Board of Health Technology Assessment, Academic Medical Centre - University of Amsterdam, the London School of Hygiene and Tropical Medicine, 2011.
- Arton R. The hidden research paper. *JAMA* 2002;287:2775–8.
- Atkins D, Best D, Briss PA, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490.
- Methodologie deGAD. *Grondslagen van onderzoek en denken in de gedragswetenschappen*. Assen: Van Gorcum, 1994.

ORIGINALITY REPORT

18%

SIMILARITY INDEX

%

INTERNET SOURCES

18%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

Maryaline Catillon. "Trends and predictors of biomedical research quality, 1990–2015: a meta-research study", BMJ Open, 2019

Publication

2%

2

Helen Bell, Sara Garfield, Sonia Khosla, Chimnay Patel, Bryony Dean Franklin. "Mixed methods study of medication-related decision support alerts experienced during electronic prescribing for inpatients at an English hospital", European Journal of Hospital Pharmacy, 2019

Publication

1%

3

Benjamin Speich, Sara Schroter, Matthias Briel, David Moher et al. "Impact of a short version of the CONSORT checklist for peer reviewers to improve the reporting of randomised controlled trials published in biomedical journals: study protocol for a randomised controlled trial", BMJ Open, 2020

Publication

1%

4

Romana Haneef, Amélie Yavchitz, Philippe Ravaud, Gabriel Baron, Ivan Oranksey, Gary Schwitzer, Isabelle Boutron. "Interpretation of health news items reported with or without spin: protocol for a prospective meta-analysis of 16 randomised controlled trials", BMJ Open, 2017

Publication

1 %

5

Stephen R Hanney. Health Research Policy and Systems, 2006

Publication

1 %

6

Kristen A Morin, John R Dabous, Frank Wojtesek, David Marsh. "Evaluating the association between urine drug screening frequency and retention in opioid agonist treatment in Ontario, Canada: a retrospective cohort study", BMJ Open, 2022

Publication

<1 %

7

Jennifer Anne Hall, Geraldine Barrett, Judith M Stephenson, Natalie Lois Edelman, Corinne Rocca. "Desire to Avoid Pregnancy scale: clinical considerations and comparison with other questions about pregnancy preferences", BMJ Sexual & Reproductive Health, 2023

Publication

<1 %

8

Reinie G. Gerrits, Michael J. van den Berg, Niek S. Klazinga, Dionne S. Kringos. "Statistics

<1 %

in Dutch policy debates on health and healthcare", Health Research Policy and Systems, 2019

Publication

9

Anke Rohwer, Taryn Young, Elizabeth Wager, Paul Garner. "Authorship, plagiarism and conflict of interest: views and practices from low/middle-income country health researchers", BMJ Open, 2017

Publication

<1 %

10

PFW Chien. "Surveying clinical surveys", BJOG An International Journal of Obstetrics & Gynaecology, 09/2009

Publication

<1 %

11

Stephen Hanney. "Who needs what from a national health research system: lessons from reforms to the English Department of Health's R&D system", Health Research Policy and Systems, 2010

Publication

<1 %

12

Fanelli, Daniele, and Vincent Larivière. "Researchers' Individual Publication Rate Has Not Increased in a Century", PLoS ONE, 2016.

Publication

<1 %

13

P. P. Groenewegen. "Health services and systems research in Europe: overview of the literature 1995 2005", The European Journal of Public Health, 01/01/2007

<1 %

14

Anja Maria Brænd, Jørund Straand, Rune Bruhn Jakobsen, Atle Klovning. "Publication and non-publication of drug trial results: a 10-year cohort of trials in Norwegian general practice", BMJ Open, 2016

Publication

<1 %

15

Anna Catharina Maria Geraedts, Sylvana de Mik, Dirk Ubbink, Mark Koelemay, Ron Balm. "Postoperative surveillance and long-term outcome after endovascular aortic aneurysm repair in the Netherlands: study protocol for the retrospective ODYSSEUS study", BMJ Open, 2020

Publication

<1 %

16

Catherine C. Motosko, Anna K. Ault, Laura L. Kimberly, M David Gothard, Roger S. Ho, Alexis Hazen. "Analysis of Spin in the Reporting of Studies of Topical Treatments of Photoaged Skin", Journal of the American Academy of Dermatology, 2018

Publication

<1 %

17

Gerry McCartney, Frank Popham, Srinivasa Vittal Katikireddi, David Walsh, Lauren Schofield. "How do trends in mortality inequalities by deprivation and education in Scotland and England & Wales compare? A repeat cross-sectional study", BMJ Open, 2017

Publication

<1 %

18 E. E. van der Wall. "Fraud and Freud: is there an association in scientific misconduct?", Netherlands Heart Journal, 2012 $<1\%$

Publication

19 Hansen, J.. "Health services research in Europe: evaluating and improving its contribution to health care policy", Journal of Health Services Research & Policy, 2011. $<1\%$

Publication

20 J K Tijdkink, K Schipper, L M Bouter, P Maclaïne Pont, J de Jonge, Y M Smulders. "How do scientists perceive the current publication culture? A qualitative focus group interview study among Dutch biomedical researchers", BMJ Open, 2016 $<1\%$

Publication

21 Leonieke J J van Mens, Henriëtte M de Jong, Inka Fluri, Michael T Nurmohamed et al. "Achieving remission in psoriatic arthritis by early initiation of TNF inhibition: a double-blind, randomised, placebo-controlled trial of golimumab plus methotrexate versus placebo plus methotrexate", Annals of the Rheumatic Diseases, 2019 $<1\%$

Publication

22 Agnes Grudniewicz, Lindsay Hedden, Seija Kromm, Ruth Lavergne, Matthew Menear, Saskia Sivananthan. "Capacity development in $<1\%$

health systems and policy research: a survey of the Canadian context", Health Research Policy and Systems, 2014

Publication

23

huiting Li, Ping Yuan, Rong Jiang, Qin-Hua Zhao et al. "Sleep-Disordered Breathing and Nocturnal Hypoxemia in Chronic Thromboembolic Pulmonary Disease", Research Square Platform LLC, 2023

Publication

<1 %

24

Reinie G. Gerrits, Dionne S. Kringos, Michael J. van den Berg, Niek S. Klazinga. "Improving interpretation of publically reported statistics on health and healthcare: the Figure Interpretation Assessment Tool (FIAT-Health)", Health Research Policy and Systems, 2018

Publication

<1 %

25

L V Rubenstein, S Hempel, M M Farmer, S M Asch, E M Yano, D Dougherty, P W Shekelle. "Finding order in heterogeneity: types of quality-improvement intervention publications", Quality and Safety in Health Care, 2008

Publication

<1 %

26

Scott A. Mogull. "Accuracy of cited "facts" in medical research articles: A review of study methodology and recalculation of quotation error rate", PLOS ONE, 2017

Publication

<1 %

27 Rebecca Mercieca-Bebber, Michael J Palmer, Michael Brundage, Melanie Calvert, Martin R Stockler, Madeleine T King. "Design, implementation and reporting strategies to reduce the instance and impact of missing patient-reported outcome (PRO) data: a systematic review", BMJ Open, 2016

<1 %

Publication

28 Cornelis A van den Bogert, Patrick C Souverein, Cecile T M Brekelmans, Susan W J Janssen et al. "Occurrence and determinants of selective reporting of clinical drug trials: design of an inception cohort study", BMJ Open, 2015

<1 %

Publication

29 J. Timothy Harrington. "Measuring Processes and Outcomes of Care", Great Health Care, 2012

<1 %

Publication

30 Lisette Schipper, Bert Meijboom, Katrien Luijkx, Jos Schols. "Front/back office considerations in the operational access to long-term care for older people: Findings of a multiple case study", International Journal of Healthcare Management, 2013

<1 %

Publication

31 J. M. Petit, S. Loubiere, M. J. Vargas-Moniz, A. Tinland et al. "Knowledge, attitudes, and

<1 %

practices about homelessness and willingness-to-pay for housing-first across 8 European countries: a survey protocol", Archives of Public Health, 2018

Publication

32

Erik von Elm, Douglas G Altman, Matthias Egger, Stuart J Pocock, Peter C Gøtzsche, Jan P Vandenbroucke. "Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies", BMJ, 2007

Publication

33

Caroline Brall, Els Maeckelberghe, Rouven Porz, Jihad Makhoul, Peter Schröder-Bäck. "Research Ethics 2.0: New Perspectives on Norms, Values, and Integrity in Genomic Research in Times of Even Scarcer Resources", Public Health Genomics, 2017

Publication

34

M. Hermanussen, M. Bilogub, A. C. Lindl, D. Harper, L. Mansukoski, C. Scheffler. "Weight and height growth of malnourished school-age children during re-feeding. Three historic studies published shortly after World War I", European Journal of Clinical Nutrition, 2018

Publication

35

Susan M. Kiene, Haruna Lule, Katelyn M. Sileo, Kazi Priyanka Silmi, Rhoda K. Wanyenze.

<1 %

<1 %

<1 %

<1 %

"Depression, alcohol use, and intimate partner violence among outpatients in rural Uganda: vulnerabilities for HIV, STIs and high risk sexual behavior", BMC Infectious Diseases, 2017

Publication

36

Giliam de Valk, Onno Goldbach. "Towards a robust β research design: on reasoning and different classes of unknowns", Journal of Intelligence History, 2020

Publication

37

A. Cecile. J. W. Janssens. "Strengthening the reporting of genetic risk prediction studies (GRIPS): explanation and elaboration : GRIPS STATEMENT: EXPLANATION AND ELABORATION", European Journal of Clinical Investigation, 09/2011

Publication

38

Benjamin Gregory Carlisle, Adélaïde Doussau, Jonathan Kimmelman. "Patient burden and clinical advances associated with postapproval monotherapy cancer drug trials: a retrospective cohort study", BMJ Open, 2020

Publication

39

D. Moher, S. Hopewell, K. F Schulz, V. Montori, P. C Gotzsche, P J Devereaux, D. Elbourne, M. Egger, D. G Altman. "CONSORT 2010 Explanation and Elaboration: updated

<1 %

<1 %

<1 %

<1 %

guidelines for reporting parallel group randomised trials", BMJ, 2010

Publication

40

Huiberts, A., M. Hjernevik, A. Mykletun, and J. C. Skogen. "Electromagnetic hypersensitivity (EHS) in the media - a qualitative content analysis of Norwegian newspapers", JRSM Short Reports, 2013.

Publication

<1 %

41

Oscar L Rueda-Ochoa, Marco A Smiderle-Gelain, Dimitris Rizopoulos, Klodian Dhana et al. "Risk factors for longitudinal changes in left ventricular diastolic function among women and men", Heart, 2019

Publication

<1 %

42

Anisa Rowhani-Farid, Kyungwan Hong, Mikas Grewal, Jesse Reynolds, Audrey D Zhang, Joshua D Wallach, Joseph S Ross. "Consistency between trials presented at conferences, their subsequent publications and press releases", BMJ Evidence-Based Medicine, 2022

Publication

<1 %

43

Agnes Dechartres, Philippe Ravaud. "Better prioritization to increase research value and decrease waste", BMC Medicine, 2015

Publication

<1 %

44

Isabelle Boutron. "Spin in Scientific Publications: A Frequent Detrimental

<1 %

Research Practice", Annals of Emergency Medicine, 2020

Publication

45

Sara Schroter, Jason Roberts, Elizabeth Loder, Donald B Penzien, Sarah Mahadeo, Timothy T Houle. "Biomedical authors' awareness of publication ethics: an international survey", BMJ Open, 2018

Publication

<1 %

46

"Abstracts", Vox Sanguinis, 2017

Publication

<1 %

47

Benedetta Armocida, Lorenzo Monasta, Susan Sawyer, Flavia Bustreo et al. "Burden of non-communicable diseases among adolescents aged 10–24 years in the EU, 1990–2019: a systematic analysis of the Global Burden of Diseases Study 2019", The Lancet Child & Adolescent Health, 2022

Publication

<1 %

48

Daniël A. Korevaar, Jérémie F. Cohen, Johannes B. Reitsma, David E. Bruns et al. "Updating standards for reporting diagnostic accuracy: the development of STARD 2015", Research Integrity and Peer Review, 2016

Publication

<1 %

49

Isla S Humphreys, Annette von Delft, Anthony Brown, Linda Hibbert et al. "HCV genotype-3a

<1 %

T cell immunity: specificity, function and impact of therapy", Gut, 2012

Publication

50

Kellia Chiu, Quinn Grundy, Lisa Bero. "'Spin' in published biomedical literature: A methodological systematic review", PLOS Biology, 2017

Publication

<1 %

51

Lavis, J.N.. "Use of research to inform public policymaking", The Lancet, 20041030/1105

Publication

<1 %

52

Sally Hopewell, Isabelle Boutron, Douglas G Altman, Philippe Ravaud. "Incorporation of assessments of risk of bias of primary studies in systematic reviews of randomised trials: a cross-sectional study", BMJ Open, 2013

Publication

<1 %

53

Tamarinde L. Haven, Joeri K. Tijdkink, Brian C. Martinson, Lex M. Bouter. "Perceptions of research integrity climate differ between academic ranks and disciplinary fields: Results from a survey among academic researchers in Amsterdam", PLOS ONE, 2019

Publication

<1 %

54

Christiaan H. Vinkers, Herm J. Lamberink, Joeri K. Tijdkink, Pauline Heus et al. "Randomized clinical trial quality has improved over time but is still not good enough: an analysis of

<1 %

176,620 randomized controlled trials
published between 1966 and 2018", Cold
Spring Harbor Laboratory, 2020

Publication

55

Liu, L., J. Wu, J. Geng, Z. Yuan, and D. Huang.
"Geographical prevalence and risk factors for
pterygium: a systematic review and meta-
analysis", BMJ Open, 2013.

Publication

<1 %

56

Willemijn LA Schäfer. "QUALICOPC, a multi-
country study evaluating quality, costs and
equity in primary care", BMC Family Practice,
2011

Publication

<1 %

Exclude quotes

On

Exclude matches

< 1 words

Exclude bibliography

Off