

Somatic embryogenesis of the selected intergeneric hybrid between Phalaenopsis 2166 and Vanda 'Saint Valentine': Application of NAA and TDZ

by Murni Dwiati

Submission date: 20-Oct-2022 12:14PM (UTC+0700)

Submission ID: 1930333435

File name: 14._2022_Dwiati_et_al_Nusantara_Bioscience_naskah.pdf (209.14K)

Word count: 4844

Character count: 24762

3 Somatic embryogenesis of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine': Application of NAA and TDZ

MURNI DWIATI[✉], AGUS HERY SUSANTO, IMAN BUDISANTOSO

Faculty of Biology, Universitas Jenderal Soedirman, Jl. Dr. Suparno 63, Karangwangkal, Purwokerto Utara, Banyumas 53122, Central Java, Indonesia.
Tel.: +62-281-638794, Fax.: +62-281-631700, ✉email: murnidwi14@gmail.com

Manuscript received: 29 May 2022. Revision accepted: 11 August 2022.

Abstract. Dwiati M, Susanto AH, Budisantoso I. 2022. Somatic embryogenesis of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine': Application of NAA and TDZ. *Nusantara Bioscience* 14: 160-165. An intergeneric hybridization between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' has successfully produced a hybrid seedling with several characters of potentially developing into plant individuals with flowers of better performance. Therefore, identical clones of the selected hybrid should be developed into PLBs by means of *in-vitro* culture technique employing somatic embryogenesis supported by the application of plant growth regulators. This study aims to unveil the effect of NAA and TDZ in stimulating the formation of identical clones of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine'. The experiment was arranged in a factorial Randomized Complete Block Design (RCBD) involving two factors, i.e., types of plant growth regulators and the levels of concentrations of each substance. It was found that the combination of NAA and TDZ had significant effect on the growth of the identical clones. The combination of NAA 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹ resulted in the clones that potentially differentiate into PLBs. This finding indicates that NAA and TDZ should be applied appropriately to stimulate somatic embryogenesis in the intergeneric hybrid.

Keywords: Intergeneric hybrid, NAA, *Phalaenopsis* 2166, TDZ, *Vanda* 'Saint Valentine'

INTRODUCTION

The members of Family Orchidaceae are mostly known as ornamental plant species due to their distinctive characteristics of flowers. However, overexploitation and alteration in land-use have caused some orchid species vulnerable to extinction. For instance, all of the 115 identified orchid species from Mount Ungaran, Central Java, Indonesia, are listed in Appendix II of the CITES and four of them are even listed in the IUCN Red List (Kurniawan et al. 2021). On the other hands, some wild orchid species can be potentially used as parental lineages to produce hybrid varieties of desirable better performances including flower colour, shape, and resistance (Li et al. 2023).

An intergeneric hybridization between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' has resulted in several hybrid seedlings which have been characterized, both phenotypically and molecularly. Based on the leaf shape, edge, and colour, the intergeneric hybrids were likely to resemble *Phalaenopsis* 2166 as the female parent, although some variations of leaf shape and colour were also observed. In general, it could be said that maternal inheritance of the phenotypic characters in the intergeneric hybridization occurred. Hence, it is reasonable that the hybrid seedlings showed the best growth when grown in New *Phalaenopsis* (NP) medium (Dwiati et al. 2020a). Then, molecular characterization by the use of *ndhE* partial

gene revealed that 11 of 14 hybrids obtained had the same sequences of *ndhE* partial gene as that of *Phalaenopsis* 2166. The sequences have now been registered in NCBI database with accession number MH646649. The other three hybrids, i.e., F1.9, F1.11, and F1.14, showed slightly different *ndhE* sequences from that of *Phalaenopsis* 2166, and they have also been registered in NCBI database as MH646651. One of the three hybrids, i.e. F1.14, has somewhat spotted reddish-purple leaf that is predictable to produce conspicuously attractive flowers thus potentially to be developed into a large number of plant individuals (Dwiati et al. 2020b; Dwiati and Susanto 2021).

To develop the promising hybrid, an *in vitro* culture technique should be employed, by which the hybrid clones are grown using ½ MS media enriched with NAA and TDZ. In this case, NAA is used for stimulating clone formation, while TDZ is intended to promote the propagation of the somatic embryos (Mayer et al. 2010; Gantait and Sinniah 2012). Some other previous studies on the stimulation of somatic embryogenesis by the use of TDZ alone or in combination with NAA have been reported, such as those *Phalaenopsis amabilis* (L.) Blume (Mose et al. 2017), *Dendrobium aqueum* Lindl. (Parthibhan et al. 2018), commercial *Phalaenopsis* hybrids (Zanello and Cardoso 2019), and *Paphiopedilum niveum* (Raf.) Stein (Soonthornkalump et al. 2019). Therefore, the objective of this study is to demonstrate the effect of NAA and TDZ application on the stimulation of somatic embryogenesis of

the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine'. Once the somatic embryos of the hybrid are produced, they could be developed further into PLBs.

MATERIALS AND METHODS

Experimental design

The study was conducted as an experiment arranged in a factorial Randomized Complete Block Design (RCBD) using two factors, i.e., types of plant growth regulators (NAA and TDZ) and their respective levels of concentrations. The NAA concentrations consisted of 0 mgL⁻¹, 0.15 mgL⁻¹, 1.0 mgL⁻¹, while those of TDZ comprised 0 mgL⁻¹, 1.0 mgL⁻¹, 1.5 mgL⁻¹, 2.0 mgL⁻¹. Each of the 12 treatment combinations thus made was given three replications resulting in a total of 36 experimental units.

Procedures

Preparation of media

A half-strength modified MS was prepared as the basal medium. This was then supplemented with 2 gL⁻¹ peptone, 50 mL⁻¹ coconut water, 150 mL⁻¹ alkaline water, 75 mgL⁻¹ vitamin C, 0.50 mgL⁻¹ PVP, 0.25 mgL⁻¹ Na panthothenat, 0.25 mgL⁻¹ pyridoxine HCl, 2 gL⁻¹ active charcoal, and 20 gL⁻¹ sucrose. NAA solution was applied to the medium corresponding to the respective treatment. pH of individual treatment was adjusted to 5.2 by dripping NaOH or HCl necessary. Each medium was added with 1.2 g agar and sterilized in an autoclave at 121°C; 0.15MPa for 20 minutes. All the media were cooled at approximately 45°C and shaken gently for homogeneity. Each medium was then added with TDZ according the respective treatment and was poured onto Petri dish.

Planting of leaf explants

The leaves of the selected hybrid were washed under running water, air-dried and put into sterile bottles. These were then added with sterile-distilled water and Tween-20 of three drops, after which the leaves were rinsed using sterile aquadest until the foams were totally removed.

Then, the leaves were sterilized using 70% (v/v) ethanol for 5 minutes, followed by HgCl₂ for 5 minutes and rinsed three times with sterile-distilled water. The leaves were put into a sterile Petri dish lined with filter paper, where they were cut into 0.5 x 0.5 cm pieces which served as explants. These were then planted onto aseptic media in the previously prepared Petri dishes corresponding to the respective treatment. Each medium was filled with two explants and put on the culture rack placed in the dark at temperature of 22°C and air humidity of 90%. The clone growth was observed daily. Since 30th day after incubation, the explants were subjected to light exposure for 12 hours and dark exposure for 12 hours alternately until they were 108 days old.

Parameters

The parameters that were examined comprised the date of clone formation, number of embryogenic clones formed (%), thickness of clones formed (mm), clone diameter (mm), and clone colour and consistency. All parameters were examined weekly since the date of explant incubation until the clones were 21 days old. Meanwhile, the development of somatic embryos was still examined 108 days after explant incubation.

Data analysis

The quantitative data obtained were analyzed using ANOVA. When significant effect of treatments was observed, further analysis was performed using Duncan Multiple Range Test (DMRT). Meanwhile, descriptive analysis was applied to the qualitative data.

RESULTS AND DISCUSSION

Clone formation

Clone formation had been observed since the third day of explant incubation, showing clone characteristics of sufficiently friable, green, and compact. These would grow into maximum at approximately three to four weeks, depending on the treatment applied. It can be seen in Table 1 that some treatment combinations of NAA and TDZ resulted in 100% of somatic embryo formation.

Table 1. Identical clones formed in the dark condition at 14th day after explant incubation

Treatment		Clone	Clone	Clone	Somatic	Clone	Clone diameter
NAA (mgL ⁻¹)	TDZ (mgL ⁻¹)	colour	consistency	formation (%)	embryos (%)	thickness (mm)	(mm)
0.0	0.0	light green	sticky	50 ^f	50 ^c	0.565 ⁱ	0.305 ^b
0.0	1.0	light green	sticky	60 ^{ef}	70 ^b	0.698 ^c	0.279 ^b
0.0	1.5	light green	sticky	65 ^{de}	70 ^b	0.669 ^f	0.306 ²²
0.0	2.0	light green	sticky	65 ^{de}	80 ^{ab}	0.639 ^g	0.317 ^{ab}
0.5	0.0	light green	less friable	70 ^{cde}	90 ^a	0.591 ^h	0.320 ^{ab}
0.5	1.0	light green	friable	100 ^a	90 ^a	0.763 ^b	0.312 ^{ab}
0.5	1.5	light green	friable	100 ^a	90 ^a	0.759 ^{bc}	0.302 ^{ab}
0.5	2.0	fresh green	friable	100 ^a	90 ^a	0.730 ^d	0.303 ^b
1.0	0.0	fresh green	friable	90 ^{ab}	80 ^{ab}	0.740 ^{cd}	0.291 ^b
1.0	1.0	dark green	friable	95 ^a	90 ^a	0.706 ^c	0.323 ^{ab}
1.0	1.5	dark green	friable	80 ^{bc}	80 ^{ab}	0.842 ^a	0.289 ⁴
1.0	2.0	dark green	friable	75 ^{cd}	85 ^a	0.663 ^f	0.355 [#]

Note: Values followed by the same letter in the same column show non-significant difference after DMRT at α 0.05

It was found in this study that clone formation of 100 percent were obtained in the combination of NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹; NAA 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹; NAA 0.5 mgL⁻¹ and TDZ 2.0 mgL⁻¹. It seemed likely that NAA of 0.5 mgL⁻¹ was the optimum concentration for promoting clone formation. In addition, the somatic embryo thus produced was 90 percent (Table 1). It was also shown from the table that a combination of NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹ at 14th day was the most optimum treatment in producing clone thickness and clone cell diameter, i.e., 0.763 mm and 0.312 mm respectively. Light green and friable clones were obtained.

The development stage of the selected intergeneric hybrid clone began when the clone cells had reached their maximum size. At the 16th day the sufficiently friable clones started to enlarge, gradually reaching their maximum size (Figure 1A). Then, the clone cells would enlarge, followed by the formation of initial globular structures, which was indicated by the division of clone cells. The next stage was characterized by the formation of several new cells inside the previous big clone cell as shown in Figure 1B. This was found at 21st day.

Clone development

It was shown in Table 2 and Figure 2A that the combination of NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹ resulted in light green and friable clones forming dome-like protuberances of 50 percent at 21st day. Here the explants had started to be subjected to light for an hour, and at 25th day the light condition was prolonged for six hours. Then, at 28th day the somatic embryo cells under the treatment began to differentiate (Figure 2B).

Further development of globular structures as observed to occur in all treatments, except in control (NAA 0 mgL⁻¹ and TDZ 0 mgL⁻¹). On the other hands, some treatments, i.e., the combination of NAA 0.5 mgL⁻¹ and TDZ 0 mgL⁻¹; NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹; 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹; NAA 1.0 mgL⁻¹ and TDZ 1.0 mgL⁻¹ had even begun to form dome-like protuberance (Figure 2A).

At 49th day the somatic embryos got into the development stage forming structures which were characterized by the occurrence of protuberances in some parts of cell side (Figure 2B). These would then develop further into structures resembling scutella (Figure 3B).



Figure 1. Development of identical clones of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' (A) diameter of clone cells with NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹ at 16th day; (B) clone obtained at 21st day

Table 2. Clones formed in the dark condition at 21st day after explant incubation

Treatment		Clone colour	Clone consistency	Further development	Clones undergoing further development (%)
NAA (mgL ⁻¹)	TDZ (mgL ⁻¹)				
0.0	0.0	whitish green	sticky	clone not develop	0 ^d
0.0	1.0	light green	compact	initial globular	30 ^{bc}
0.0	1.5	light green	compact	tree-like	20 ^c
0.0	2.0	light green	compact	initial globular	40 ^{ab}
0.5	0.0	light green	friable	dome-like	40 ^{ab}
0.5	1.0	light green	friable	dome-like	50 ^a
0.5	1.5	light green	friable	dome-like	30 ^{bc}
0.5	2.0	fresh green	friable	tree-like	30 ^{bc}
1.0	0.0	fresh green	friable	globular	35 ^b
1.0	1.0	dark green	friable	dome-like	40 ^{ab}
1.0	1.5	dark green	friable	last globular	50 ^d
1.0	2.0	dark green	friable	globular	35 ^b

Note: Values followed by the same letter in the same column show non-significant difference after DMRT at α 0.05



Figure 2. Further development of globular structure of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' (A) dome-like protuberance with NAA 0.5 mgL⁻¹ and TDZ 1.0 mgL⁻¹ at 28th day; (B) somatic embryo with NAA 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹ at 49th day



Figure 3. The next development of somatic embryos of the selected intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' (A) somatic embryo with NAA 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹ under light condition for an hour at 70th day; (B) somatic embryo with NAA 0.5 mgL⁻¹ and TDZ 1.5 mgL⁻¹ under 12 hour-light and 12 hour-dark condition at 108th day

Discussion

Slightly different from our results that 50% clones were still formed in the medium without supply of NAA and TDZ (Table 1), recalcitrant leaf explants in *Cymbidium eburneum* Lindl. were observed in the basal medium of MS. No regeneration occurred where the explants turned brown and died within 10 weeks (Sembi et al. 2020). Similar finding (no somatic embryo formation) reported in leaf explant in *Schlotheimia plicata* Blume when cultured in MS medium in the absence of plant growth regulator (Manokari et al. 2021). Other report on indirect somatic embryogenesis is where calli of the intergeneric hybrids between *Aranda* Wan Chark Kuan 'Blue' and *Vanda coerulea* Griff. ex Lindl. appeared from the incision scar relatively fast since the third day of explant incubation (Gantait and Sinniah 2012). Wounding is considered important in assisting both direct and indirect somatic embryogenesis, because it stimulates cell division (Rojas-Herrera and Loyola-Vargas 2002).

Regeneration responses in leaf explant culture of *C. eburneum* were observed in the applications of NAA and

BA, where the highest response (83.3%) was obtained in the combination of 2 mgL⁻¹ NAA and 0.5 mgL⁻¹ BA (Sembi et al. 2020). In our study we use TDZ instead of BA for stimulating further development of the clones into somatic embryos, since TDZ has been generally used in orchid tissue culture to enhance somatic embryogenesis (Vu et al. 2012; Mahendran and Bai 2016). It was also reported that TDZ showed better efficacy over other purine-types of cytokinins such as BA in inducing somatic embryogenesis in orchids (Bhattacharyya et al. 2018). The fastest PLB induction in the intergeneric hybrids between *Aranda* Wan Chark Kuan 'Blue' and *V. coerulea* using leaf explants was observed in the treatment of TDZ 1.5 mgL⁻¹ (Gantait and Sinniah 2012). Some other studies showed that TDZ was proved very effective in inducing somatic embryogenesis in several orchid species, such as *Renanthera* Tom Thumb 'Qilin' (Wu et al. 2012), *P. bilis* (Mose et al. 2017), and *P. niveum* (Soonthornkalump et al. 2019). TDZ can be used in replace of cytokinin, so that it can be applied as plant growth regulator along with auxin in an in-vitro culture media

(Kou et al. 2016). As plant growth regulator, TDZ should be applied not exceed 3 mgL^{-1} . The high level of TDZ (3 to 5 mgL^{-1}) will inhibit cytokinin oxidase (Soonthornkalump et al. 2019). Basically, variation in the application of growth regulators, especially auxin and cytokinin, in an in vitro culture media, could affect somatic embryogenesis (Guo et al. 2011; Moradi et al. 2017).

Since 30th day after incubation, the explants of the selected intergeneric hybrid of *Phalaenopsis* 2166 x *Vanda* 'Saint Valentine' were subjected 12 hour-exposure to light and another 12 hours in the dark. As a comparison, the explants of *Phalaenopsis* Classic spotted pink started to form last globular structure at 23rd day and the embryos began to form coleoptelar (Pereira et al. 2019). Meanwhile, direct somatic embryogenesis from leaf explants in *Phalaenopsis* 'Little Steve' revealed that somatic embryos were formed at 30th day after explant incubation in the dark (Kuo et al. 2005).

Somatic embryogenesis in *P. amabilis* was reported to begin at 8th week when grown in NP media supplemented with TDZ. The most rapid somatic embryogenesis was obtained with TDZ 3.0 mgL^{-1} at 11th day using leaf explants, while the slowest one was found with TDZ 3.0 mgL^{-1} using stem explants (Mose et al. 2017). In this study we found that somatic embryogenesis of the intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' began at 21st day in the modified $\frac{1}{2}$ MS medium. Meanwhile, *Vanda tricolor* Lindl. shoots produced from somatic embryogenesis showed the best development when subcultured in the New Dogashima (ND) medium without application of any plant growth regulator (Ashihah et al. 2022).

Regeneration of somatic embryogenesis in orchids began at 15th to 30th day in the concentration of TDZ ranging between 0.001 and 5 mgL^{-1} (Shen et al. 2018). Reduced auxin in the last development stage of somatic embryogenesis was needed, especially for stimulating PLB proliferation and differentiation (Yang and Zhang 2010). TDZ concentration and its interaction with light spectra were found highly determining direct somatic embryogenesis in *Phalaenopsis* orchids. The concentration of 3 mgL^{-1} in interaction with red and far red light spectra was the efficient treatment to induce direct somatic embryogenesis in the orchids without somaclonal variation (Boldaji et al. 2021).

While no callus was formed in our study, pre-embryo is a further development of callus in the indirect somatic embryogenesis, which is characterized by two bipolar centers of meristems. These structures will develop into root and stem meristem respectively (Seth et al. 2017; Shen et al. 2018). Histological examination shows that callus resulting from somatic embryogenesis will develop sequentially into PLBs, which consist of some meristematic tissues undergoing further development to form roots, stem, and leaves (Sherif et al. 2018).

Some factors have direct effects on the somatic embryogenesis of orchids. They are genotypes, growth regulators, and media (Campos et al. 2017; Zanello and Cardoso 2019). Half strength MS is the most common media used, in which N is in form of nitrate (NH_4NO_3) in a

sufficiently high concentration, i.e., 1.7 mgL^{-1} . In addition, KNO_3 of 1.9 mgL^{-1} is also contained in $\frac{1}{2}\text{MS}$ media. Most plants absorb N in form of nitrate. Both NH_4NO_3 and KNO_3 can be used to stimulate somatic embryogenesis (Zanello and Cardoso 2019). NP was reported as the media with N in form of nitrate suitable for *P. amabilis*. This media contained NH_4NO_3 of 82 mgL^{-1} , KNO_3 of 424 mgL^{-1} , $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ of 443.04 mgL^{-1} , and $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ of 256.4 mgL^{-1} (Mose et al. 2017). Nitrogen, in form of either potassium nitrate or calcium nitrate, is very good to stimulate somatic embryogenesis, while that in form of ammonium nitrate less stimulate somatic embryogenesis. Nevertheless, explants in the absence of ammonium nitrate in the growth media will fail to undergo somatic embryogenesis (Méndez-Hernández et al. 2019).

The presence of TDZ in the culture media of *Oncidium flexuosum* (Kunth) Lindl. without light would stimulate regeneration of PLBs. Pre-embryos having no chlorophyll were formed in the dark condition, so that somatic embryogenesis occurred in the absence of chlorophyll. After treatment with no growth regulator and incubation in the light condition, embryos would be green initiating to form PLBs (Zanello and Cardoso 2019). It was proved that in the early stages of PLB formation, characteristics of somatic embryonic callus similar to zygotic embryo development were observed, indicating that PLBs were truly somatic embryos of orchids (Lee et al. 2013).

Somatic embryogenesis could result from proliferation of young PLBs that were cultured in the MVW media containing NAA 0.1 mgL^{-1} . The increasing accumulation of endogenous auxin through the application of exogenous auxin in the early stages of somatic embryogenesis was needed. In the next stages of development reduced level of auxin enabled rapid proliferation and differentiation of meristems, which in turn would stimulate the emergence of shoots. Then, the plantlets thus produced were moved into MVW media without growth regulator (Soonthornkalump et al. 2019).

In conclusions, our present study found that $\frac{1}{2}$ MS medium supplemented with the combination of NAA 0.5 mgL^{-1} and TDZ 1.5 mgL^{-1} produced identical clones of the intergeneric hybrid between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine' that showed high potential of differentiating into PLBs. This indicates that somatic embryogenesis in the selected hybrid could be stimulated by proper application of both NAA and TDZ.

ACKNOWLEDGEMENTS

The authors would like to express their gratefulness to the Institute of Research and Community Service of Universitas Jenderal Soedirman, Central Java, Indonesia, for funding this project with the scheme of Applied Research Grand financial year 2019 with director decree number 158/UN23/14/PN01.00/2019. Appreciation is also addressed to Supriyono for laboratory work assistance.

REFERENCES

- Ashihah FR, Rineksane IA, Astuti A. 2022. New Dogashima medium as subculture medium improve the growth of *Vanda tricolor* shoots from embryogenesis. IOP Conf Ser Earth Environ Sci 985(1): 012006. DOI: 10.1088/1755-1315/985/1/012006.
- Bhattacharyya P, Paul P, Kumaria S, Tandon P. 2018. Transverse thin cell layer (t-TCL)-mediated improvised micropropagation protocol for endangered medicinal orchid *Dendrobium aphyllum* Roxb: An integrated phytomolecular approach. Acta Physiologiae Plantarum 40: 1-14. DOI: 10.1007/s11738-018-2703-y.
- Boldaji HN, Dylami SD, Aliniaefard S, Norouzi M. 2021. Efficient method for direct embryogenesis in *Phalaenopsis* orchid. Intl J Hort Sci Technol 9 (2): 37-50. DOI: 10.22059/ijhst.2020.296696.339.
- Campos NA, Panis B, Carpentier SC. 2017. Somatic embryogenesis in coffee: The evolution of biotechnology and the integration of omics technologies offer great opportunities. Front Plant Sci 8: 1-12. DOI: 10.3389/fpls.2017.01460.
- Dwiyati M, Hardiyati T, Susanto AH, Chasanah T, Palupi D. 2020a. Growth medium for intergeneric hybrids between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine'. IOP Conf Ser Earth Environ Sci 593 (1): 012023. DOI: 10.1088/1755-1315/593/1/012023.
- Dwiyati M, Susanto AH, Prayoga L. 2020b. Intergeneric hybrids of *Phalaenopsis* 2166 x *Vanda* 'Saint Valentine' showing maternal inheritance: Genetic analysis based on *ndhE* partial gene. Biodiversitas 21 (11): 5138-5145. DOI: 10.13057/biodiv/d211119.
- Dwiyati M, Susanto AH. 2021. Intergeneric hybridization between *Phalaenopsis* 2166 and *Vanda* 'Saint Valentine': Characterization of parents using *ndhE* cpDNA partial sequence. J Trop Biodivers Biotechnol 6 (3): 1-6. DOI: 10.22146/JTBB.65658.
- Gantait S, Sinniah UR. 2012. Rapid micropropagation of monopodial orchid hybrid (*Aranda* Wan Chark Kuan "Blue" x *Vanda coerulea* Griff. ex. Lindl.) through direct induction of protocorm-like bodies from leaf segments. Plant Growth Regul 68 (2): 129-140. DOI: 10.1007/s10725-012-9698-y.
- Guo B, AbbasiBH, Zeb A, Xu LL, Wei YH. 2011. Thidiazuron: A multi-dimensional plant growth regulator. Afr J Biotechnol 10 (45): 8984-9000. DOI: 10.5897/ajb11.636.
- Kou Y, Yuan C, Zhao Q, Liu G, Nie J, Ma Z, Cheng C, Teixeira da Silva JA, Zhao L. 2016. Thidiazuron triggers morphogenesis in *Rosa canina* L. protocorm-like bodies by changing incipient cell fate. Front Plant Sci 7: 1-13. DOI: 10.3389/fpls.2016.00557.
- Kuo HL, Chen JT, Chang WC. 2005. Efficient plant regeneration through direct somatic embryogenesis from leaf explants of *Phalaenopsis* 'Little Steve'. In Vitro Cell Dev Biol Plant 41 (4): 453-456. DOI: 10.1079/IVP2005644.
- Kurniawan FH, Nazar L, Anjrawati R, Sasono HD, Rahayuningsih M. 2021. Orchids of Mount Ungaran (Indonesia) compiled from a decade of data collections between 2010 and 2021. Nusantara Biosci 13 (2): 238-252. DOI: 10.13057/nusbiosci/n130214.
- Lee YI, Hsu ST, Yeung EC. 2013. Orchid protocorm-like bodies are somatic embryos. Am J Bot 100 (11): 2121-2131. DOI: 10.3732/ajb.1300193.
- Li C, Dong N, Zhao Y, Wu S, Liu Z, Zhai J. 2021. A review for the breeding of orchids: Current achievements and prospects. Hort Plant J 7 (5): 380-392. DOI: 10.1016/j.hpj.2021.02.006.
- Mahendran G, Bai VN. 2016. Direct somatic embryogenesis of *Malaxis densiflora* (A. Rich.) Kuntze. J Genet Eng Biotechnol 14 (1): 77-81. DOI: 10.1016/j.jgeb.2015.11.003.
- Manokari M, Priyadarshini S, Shekhawat MS. 2021. Direct somatic embryogenesis using leaf explants and short term storage of synseeds in *Spathoglottis plicata* Blume. Plant Cell Tissue Organ Cult 145 (2): 321-331. DOI: 10.1007/s11240-021-02010-9.
- Mayer JLS, Stancato GC, Appezzato-Da-Glória B. 2010. Direct regeneration of protocorm-like bodies (PLBs) from leaf apices of *Oncidium flexuosum* Sims (Orchidaceae). Plant Cell Tissue Organ Cult 103 (3): 411-416. DOI: 10.1007/s11240-010-9782-9.
- Méndez-Hemández HA, Ledezma-Rodríguez M, Avilez-Montalvo RN, Juárez-Gómez YL, Skeete A, Avilez-Montalvo J, De-La-Peña C, Loyola-Vargas VM. 2019. Signaling overview of plant somatic embryogenesis. Front Plant Sci 10: 1-15. DOI: 10.3389/fpls.2019.00077.
- Moradi S, Daylami SD, Arab M, Vahdati K. 2017. Direct somatic embryogenesis in *Epipactis veratrifolia*, a temperate terrestrial orchid. J Hort Sci Biotechnol 92 (1): 88-97. DOI: 10.1080/14620316.2016.1228434.
- Mose W, Indrianto A, Purwantoro A, Semiarti E. 2017. The influence of thidiazuron on direct somatic embryo formation from various types of explant in *Phalaenopsis amabilis* (L.) Blume orchid. Hayati J Biosci 24 (4): 201-205. DOI: 10.1016/j.hjb.2017.11.005.
- Parthibhan S, Rao M V, Teixeira da Silva JA, Kumar TS. 2018. Somatic embryogenesis from stem thin cell layers of *Dendrobium aequum*. Biologia Plantarum 62 (3): 439-450. DOI: 10.1007/s10535-018-0769-4.
- Pereira JAF, Ferreira LT, De Moraes MB, Uliisses C. 2019. Somatic embryos from *Phalaenopsis* classic spotted pink (Orchidaceae) protocorms. Ciencia Rural 49 (7): 1-5. DOI: 10.1590/0103-8478cr20180822.
- Rojas-Herrera R, Loyola-Vargas VM. 2002. Induction of a class III acidic chitinase in foliar explants of *Coffea arabica* L. during somatic embryogenesis and wounding. Plant Sci 163 (4): 705-711. DOI: 10.1016/S0168-9452(02)00156-5.
- Sembi JK, Pathak P, Vema J. 2020. Regeneration competence of leaf explants in *Cymbidium eburneum* Lindl. J Orchid Soc India 34: 17-21.
- Seth S, Rath SC, Rout GR, Panigrahi J. 2017. Somatic embryogenesis in *Abutilon indicum* (L.) sweet and assessment of genetic homogeneity using SCoT markers. Plant Biosyst 151 (4): 704-714. DOI: 10.1080/11263504.2016.1211193.
- Shen HJ, Chen JT, Chung HH, Chang WC. 2018. Plant regeneration via direct somatic embryogenesis from leaf explants of *Tolumnia* Louise Elmore 'Elsa'. Bot Stud 59 (1): 1-7. DOI: 10.1186/s40529-018-0220-3.
- Sherif NA, Franklin Benjamin JH, Senthil Kumar T, Rao MV. 2018. Somatic embryogenesis, acclimatization and genetic homogeneity assessment of regenerated plantlets of *Anoetochilus elatus* Lindl., an endangered terrestrial jewel orchid. Plant Cell Tissue Organ Cult 132 (2): 303-316. DOI: 10.1007/s11240-017-1330-4.
- Soonthomkalump S, Nakkanong K, Meesawat U. 2019. In vitro cloning via direct somatic embryogenesis and genetic stability assessment of *Paphiopedilum niveum* (Rchb.f.) Stein: The endangered Venus's slipper orchid. In Vitro Cell Dev Biol Plant 55 (3): 265-276. DOI: 10.1007/s11627-019-09981-7.
- Wu K, Zeng S, Teixeira da Silva JA, Chen Z, Zhang J, Yang Y, Duan J. 2012. Efficient regeneration of *Renanthera* Tom Thumb "Qilin" from leaf explants. Scientia Hort 135: 194-201. DOI: 10.1016/j.scienta.2011.11.028.
- Yang X, Zhang X. 2010. Regulation of somatic embryogenesis in higher plants. Critical Rev Plant Sci 29 (1): 36-57. DOI: 10.1080/07352680903436291.
- Zanello CA, Cardoso JC. 2019. PLBs induction and clonal plantlet regeneration from leaf segment of commercial hybrids of *Phalaenopsis*. J Hort Sci Biotechnol 00 (00): 1-5. DOI: 10.1080/14620316.2019.1600384.

Somatic embryogenesis of the selected intergeneric hybrid between Phalaenopsis 2166 and Vanda 'Saint Valentine': Application of NAA and TDZ

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

repository.ubaya.ac.id

Internet Source

2%

2

Saikat Gantait, Nirmal Mandal, Somnath Bhattacharyya, Prakash Kanti Das. "An elite protocol for accelerated quality-cloning in *Gerbera jamesonii* Bolus cv. Sciella", In Vitro Cellular & Developmental Biology - Plant, 2010

Publication

1%

3

www.sciencegate.app

Internet Source

1%

4

DF Alfauzaan, Warid. "Consumer preference for some genotypes of ornamental chili (*Capsicum* spp.) population F2", IOP Conference Series: Earth and Environmental Science, 2021

Publication

1%

5

jurnal.unmuhjember.ac.id

Internet Source

1%

6	Hafiz Muhammad Rizwan, Muhammad Irshad, Bizhu He, Shuang Liu, Xiaocao Lu, Yueting Sun, Dongliang Qiu. "Role of reduced nitrogen for induction of embryogenic callus induction and regeneration of plantlets in <i>Abelmoschus esculentus</i> L.", South African Journal of Botany, 2020 Publication	1 %
7	revistabionatura.com Internet Source	1 %
8	jurnal.uns.ac.id Internet Source	1 %
9	journal.ipb.ac.id Internet Source	1 %
10	mdpi-res.com Internet Source	<1 %
11	www.scilit.net Internet Source	<1 %
12	Sujatha Govindaraj. "Chapter 22 Thidiazuron: A Potent Phytohormone for In Vitro Regeneration", Springer Science and Business Media LLC, 2018 Publication	<1 %
13	www.floriculturealliance.org Internet Source	<1 %

14 Madi Waheed Al-Mayahi Ahmed. <1 %
"Thidiazuron-induced in vitro bud
organogenesis of the date palm (Phoenix
dactylifera L.) CV. Hillawi", African Journal of
Biotechnology, 2014
Publication

15 www.schinafish.cn <1 %
Internet Source

16 jjbs.hu.edu.jo <1 %
Internet Source

17 "Somatic Embryogenesis in Ornamentals and
Its Applications", Springer Nature, 2016 <1 %
Publication

18 Seyed Mohammad Hashemi, Mohammadreza
Naghavi, Mehdi Ghorbani, Peiman Zandi. <1 %
"Effects of Abiotic Elicitors on Expression and
Accumulation of Three Candidate
Benzophenanthridine Alkaloids in Cultured
Greater Celandine Cells ", Research Square,
2020
Publication

19 www.ijcmas.com <1 %
Internet Source

20 www.tandfonline.com <1 %
Internet Source

21 garuda.ristekbrin.go.id
Internet Source

<1 %

22

J Rattin, P Wagner, D Ferreyro, A Riverti, E Giardina, A Benedetto. "Roots Partially Drive Super Sweet Maize Yield", Journal of Experimental Agriculture International, 2017

Publication

<1 %

23

open.library.ubc.ca

Internet Source

<1 %

24

www.coursehero.com

Internet Source

<1 %

25

www.qucosa.de

Internet Source

<1 %

26

123dok.com

Internet Source

<1 %

27

M. Manokari, S. Priyadharshini, Phanikanth Jogam, Abhijit Dey, Mahipal S. Shekhawat. "Meta-topolin and liquid medium mediated enhanced micropropagation via ex vitro rooting in Vanilla planifolia Jacks. ex Andrews", Plant Cell, Tissue and Organ Culture (PCTOC), 2021

Publication

<1 %

28

researcherslinks.com

Internet Source

<1 %

www.ajol.info

30

Arun Kumar Khajuria, NS Bisht, Nisha Bhagat. "In vitro organogenesis and plant regeneration of *Thymus serpyllum* L.: an important aromatic medicinal plant", *In Vitro Cellular & Developmental Biology - Plant*, 2020

Publication

<1 %

31

Etienne Yergeau, David W. Sommerville, Emilie Maheux, Vladimir Vujanovic, Chantal Hamel, Joann K. Whalen, Marc St-Arnaud. "Relationships between *Fusarium* population structure, soil nutrient status and disease incidence in field-grown asparagus", *FEMS Microbiology Ecology*, 2006

Publication

<1 %

32

core.ac.uk
Internet Source

<1 %

33

estudogeral.sib.uc.pt
Internet Source

<1 %

34

Leimapokpam Tikendra, Thoungamba Amom, Potshangbam Nongdam. "Molecular genetic homogeneity assessment of micropropagated *Dendrobium moschatum* Sw. - A rare medicinal orchid, using RAPD and ISSR markers", *Plant Gene*, 2019

Publication

<1 %

- | | | |
|----|--|------|
| 35 | Yung-I Lee, Shan-Te Hsu, Edward C. Yeung. "Orchid protocorm-like bodies are somatic embryos", American Journal of Botany, 2013
Publication | <1 % |
| 36 | kb.psu.ac.th
Internet Source | <1 % |
| 37 | nopr.niscair.res.in
Internet Source | <1 % |
| 38 | nveo.org
Internet Source | <1 % |
| 39 | vascbio.brandnews.at
Internet Source | <1 % |
| 40 | www.iufro.org
Internet Source | <1 % |
| 41 | "Protocols for Micropropagation of Selected Economically-Important Horticultural Plants", Springer Science and Business Media LLC, 2013
Publication | <1 % |
| 42 | Ahmed, Sheikh Rashel, Md Moniruzzaman Shohag Howlader, Pijush Sutradhar, and Sabina Yasmin. "An efficient protocol for in vitro regeneration of Stevia rebaudiana", Asian Journal of Medical and Biological Research, 2016.
Publication | <1 % |

43 Amitav Bhattacharya. "Soil Water Deficit and Physiological Issues in Plants", Springer Science and Business Media LLC, 2021 $<1\%$
Publication

44 Marlúcia Souza Souza Pádua, Luciano Viela Paiva, Luis Gustavo Texeira da Silva, Luciano Coutinho Silva et al. "In vitro development and acclimatization of dendezeiro (Elaeis guineensis)", Revista Árvore, 2014 $<1\%$
Publication

45 István Mosonyi. "In vitro regeneráció és szaporodás különleges útja Spathiphyllum hibridek esetén", Corvinus University of Budapest, 2014 $<1\%$
Publication

46 Tim Wing Yam, Joseph Arditti. "Micropropagation of Orchids", Wiley, 2017 $<1\%$
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On