

Immunomodulatory effects of black solo garlic (*Allium sativum* L.) on streptozotocin-induced diabetes in Wistar rats

by Sarie Yono

Submission date: 22-Nov-2022 08:23AM (UTC+0700)

Submission ID: 1960792870

File name: jurnal_heliyon.pdf (398.99K)

Word count: 3423

Character count: 18746



Research article

Immunomodulatory effects of black solo garlic (*Allium sativum* L.) on streptozotocin-induced diabetes in Wistar rats

Saryono^{a,*}, Desiyani Nani^a, Atikah Proverawati^b, Sarmoko^c

^a Nursing Department, Faculty of Health Sciences, University of Jenderal Soedirman, Dr. Soeparno Street Purwokerto 53123, Indonesia

^b Nutrition Department, Faculty of Health Sciences, University of Jenderal Soedirman, Dr. Soeparno Street Purwokerto 53123, Indonesia

^c Pharmacy Department, Faculty of Health Sciences, University of Jenderal Soedirman, Dr. Soeparno Street Purwokerto 53123, Indonesia

ARTICLE INFO

Keywords:

Allium sativum
Black solo garlic
Immunomodulator
Interferon
Interleukin
Streptozotocin-induced diabetes

ABSTRACT

Diabetes mellitus is a chronic disease that leads to different complications. Therefore, this study aims to investigate the immunomodulatory effects of the black solo garlic on streptozotocin (STZ)-induced diabetic rats. The Wistar rats were grouped into six groups of: normal control, negative control, treatment dose of 6.5 g/kg, 13.5 g/kg, and 26 g/kg body weight, and positive control glibenclamide. In addition to normal control, rats were induced with STZ on day 8–11. Also, steeping black solo garlic glibenclamide was administered on the day 12–19. The experimental animals were sacrificed on day 20 and tumor necrosis factor (TNF- α), interleukin-6 (IL-6), interleukin-1 β (IL-1 β) and interferon gamma (IFN- γ) were measured using ELISA. The results showed that the administration of steeping black solo garlic significantly decreased levels of IL-1 β , IL-6 and TNF- α as well as increased IFN- γ with the immunity of STZ-induced rats.

1. Introduction

Diabetes mellitus (DM) is a disease characterized by impaired metabolism of carbohydrates, proteins and fats due to insufficient secretion of insulin. Furthermore, this complication is caused by decreased insulin sensitivity in the tissues, and it is the leading cause of kidney disease, blindness, and amputations [1]. Diabetes mellitus is one of chronic inflammatory disease that suppresses cellular immunity in diabetic [2].

Diabetes mellitus is a chronic condition that occurs when the pancreas do not create enough insulin, or when the body cannot use the insulin adequately. Insulin deficiency in diabetes increases blood glucose resulting in impaired microcirculation as well as increased oxidative stress resulting in prolonged inflammation [3]. This condition is different from inflammation in general. Therefore, agent that can lower blood glucose as well as an anti-inflammatory is required. Diabetes mellitus is associated with high levels of cytokines TNF- α , IL-6, IL-1 β , and decreased interferon- γ [4, 5]. TNF- α may change permeability of glomerular and albuminuria by promoting local reactive oxygen species (ROS) production [6]. IL-6 is a powerful predictor of the development of diabetes complications such as diabetic nephropathy [7].

The body's immunity can increase through nutritious food, adequate rest, and routine exercise. In addition, some plants have beneficial effect as immunomodulator such as garlic (*Allium sativum* L.). Fermentation of

garlic produced a black garlic with a sour taste rather than a pungent flavor [8]. Black garlic contains reduced sugars, polyphenols, flavonoids, Amadori and Heyns, leucine, isoleucine, phenylalanine, S-allyl-cysteine, and alkaloids content higher than those of fresh garlic [9]. Black garlic have several functions, such as an antioxidants, anti-bacterial, anti-allergic, anti-diabetic, anti-inflammatory, and anti-carcinogenic effects [10]. Black garlic increases natural killer cell cytotoxicity and increases the proliferation of B and T lymphocytes and macrophages. The previous study showed that black ordinary garlic extract increases the immune system [11] and recover kidney cells [12].

One of type garlic is single clove or solo garlic (*Allium sativum* 'Solo garlic'), which contains unstable compounds such as allicin [13]. The allicin ratio in a single clove is equivalent to 5–6 of ordinary garlic. In addition, black solo garlic (BSG) contains more S-allyl-cysteine six time than fresh garlic, which increase antioxidant activity [14]. Furthermore, the content of polyphenols, vitamins, minerals and flavonoids are increased during the fermentation [8]. However, there has not been any study on immunomodulatory as well as an anti-inflammatory of black solo garlic. Therefore, this study aims to evaluate the immunomodulatory and anti-inflammatory effect of black solo garlic in streptozotocin (STZ)-induced rats. The present study was designed to investigate the changes of proinflammatory mediators (TNF- α , IL-6, IL-1 β , IFN- γ) to verify the protective effects of BSG in DM rats.

* Corresponding author.

E-mail address: saryono2016@unsoed.ac.id (Saryono).

<https://doi.org/10.1016/j.heliyon.2021.e08493>

Received 23 June 2021; Received in revised form 2 September 2021; Accepted 24 November 2021

2405-8440/© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

2. Materials and methods

2.1. Plant

The fresh solo garlies were obtained from the Brebes region in Central Java and validated by the Plant Taxonomy Laboratory at the Faculty of Biology at Jenderal Soedirman University, Indonesia.

2.2. Fermentation of solo garlies

The selected garlies were approximately the same size and were fermented in the fermentation modified apparatus by using rice cooker. Furthermore, they were arranged in different layers and covered with tissue paper. The rice cooker was set in the warm mode (temperature 60–80 °C) left for 21 days and discoloration was observed once every 3 days. After 21 days, the garlic turned into a black chewy texture.

2.3. Preparation of aqueous black solo garlic extract

Black solo garlic were peeled and weighed according to the dose, then mashed using a pestle and mortar. The crushed garlic was placed in a glass, then hot water (at a temperature of 80–90 °C) was added and stirred for 15 min. The mixture was filtered to obtain steeping black solo garlic. The steeping were further analysed by using GC-MS, and we found the same results as reported by Tran et al. (2018) [15].

2.4. Animals

The study was conducted using a pre- and post-test approach with a control group design after obtaining ethical eligibility from ethics committee, Faculty of Health Sciences, Jenderal Soedirman University No: 152/EC/KEPK/VII/2020. White male Wistar strain rats aged around 6–7 weeks with a body weight range of 150–200 g were used. They were given *ad libitum* access to standard food and distilled water.

2.5. Streptozotocin-induced diabetic rats

Streptozotocin was dissolved in 2.5 ml of citrate buffer 0.05 M and the experimental animals were induced by STZ 50 mg/kg body weight on the day 8th. In addition, they were fasted for 6–8 h before STZ induction. Induction was performed for 3 days.

2.6. Treatments

The total number of animals was 30 rats, which were divided randomly into six group: (i) normal, (ii) STZ-induced rat, (iii) STZ-induced rat treated with glibenclamide 0.09 mg/kg body weight (positive control), STZ-induced rat treated with steeping black solo garlic dose

of (iv) 6.5 (low dose), (v) 13.5 (medium dose), and (vi) 26 g/kg body weight (high dose). After STZ-induction for 3 days, the animals were treated with steeping black solo garlic for 7 days on the day 11 to 19th. The same schedule was also applied for glibenclamide treatment (Figure 1). Briefly, the black solo garlic powder was measured based on the bodyweight of rats to obtain those three doses. The measured powder was diluted in the 3.6 mL hot water and stirred for 15 min to obtain steeping of black solo garlic. Steeping black solo garlic was given per oral once at night based on the dose and body weight of each group.

2.7. Measurement of cytokines

Blood was drawn using a hematocrit capillary pipette which was inserted into the *plexus orbitalis* at the corner of the eye in the day 11st and 19th. The ELISA kits were used to check cytokine IL-6, TNF- α , IL-1 β , and IFN- γ (BT Laboratories, Shanghai) based on the manufacturer protocol. Briefly, this ELISA kit uses the Sandwich-ELISA principle. The plate has been pre-coated with an antibody specific to rat cytokine. The optical density (OD) was measured using the ELISA Reader (Labotrone, Germany) at a wavelength of 450 nm. The OD value was proportional to the concentration of rat cytokine. Concentration of rat cytokine in the samples was calculated by comparing the OD of the samples to the standard curve.

2.8. Statistical analysis

All data were presented as mean \pm SEM. GraphPad Prism 8 (California, US) was used to generate the figure. Furthermore, the mean scores among the treatment groups were compared with the one-way ANOVA with a Tukey post-hoc test. Pre- and post-test treatment were analyzed by t-test. $p < 0.05$ was considered as significant.

3. Results

3.1. The effect of steeping black solo garlic on pro-inflammatory cytokines

The results showed that STZ induction significantly increased IL-1 β , IL-6, and TNF- α level compared normal group ($p < 0.0001$) (Figure 2). The administration of black solo garlic at dose of 6.5 g/kg, 13.5 g/kg and 26 g/kg body weight had significantly reduced the level of IL-1 β , IL-6, and TNF- α compared to untreated diabetic rats ($p < 0.0001$). Interestingly, the treatment of black solo garlic at dose of 13.5 g/kg and 26 g/kg body weight was able to reduce IL-1 β , IL-6, and TNF- α level lower than those of glibenclamide. The effects on IL-1 β , IL-6, and TNF- α were dose dependent, the reduced level of those cytokines in rats administrated with the medium and high black solo garlic dose (13.5 and 26 g/kg, respectively) being significantly greater than that in the rats receiving low black solo garlic dose (6.5 g/kg). Therefore, treatment of the black




Figure 1. Study design in animal experiments. STZ, streptozotocin; BSG, black solo garlic.




Figure 2. Treatment of black solo garlic reduced pro-inflammatory cytokines. (A) IL-6, (B) IL-1 β , (C) TNF- α level in the treatment of black solo garlic in 6.5, 13.5, 26 g/kg doses and glibenclamide 0.09 mg/kg in streptozotocin-induced rats were determined by ELISA. Statistical significance for the difference among groups: ****p < 0.0001; ***p < 0.0005; *p < 0.05. Statistical significance for the difference between the data of pre-test group vs post-test groups: ##p < 0.01; #p < 0.05.

solo garlic dose of 13.5 g/kg are equal to those 26 g/kg body weight as an anti-inflammatory agent.

3.2. The effect of steeping black solo garlic on IFN- γ

Streptozotocin induction produced a significant decreased of IFN- γ compared to the normal group (p < 0.0001) (Figure 3). Black solo garlic (13.5–26 g/kg)-treated diabetic rats had significantly increased IFN- γ level compared to the untreated diabetic rats (p < 0.0001). The effects on IFN- γ were dose dependent, IFN- γ level of rats having been administered

medium or high black solo garlic dose being significantly higher than those in the diabetic rats receiving the low black solo garlic dose. Moreover, the treatment of black solo garlic at dose of 13.5 g/kg and 26 g/kg body weight was able to increase IFN- γ level higher than those of glibenclamide.

4. Discussion

The administration of steeping black solo garlic decreases the level of IL-1 β , IL-6 and TNF- α and increases IFN- γ in experimental animals. These




Figure 3. Treatment of black solo garlic increased IFN- γ . IFN- γ level in the treatment of black solo garlic in 6.5, 13.5, 26 g/kg doses and glibenclamide 0.09 mg/kg in streptozotocin-induced rats was determined by ELISA. Statistical significance for the difference among groups: ****p < 0.0001; ***p < 0.0005; **p < 0.01. Statistical significance for the difference between the data of pre-test group vs post-test groups: ##p < 0.01; #p < 0.05.

results suggested that black solo garlic is potential to be used as an immunomodulator and antiinflammation in diabetic condition to prevent complication.

Induction of STZ generates free radicals which oxidize β -pancreatic cells by means of alkylating DNA, damaging mitochondria, and inhibiting the O-GlcNAcase enzyme. The oxidation process induces toxicity through a free radical chain reaction. This results in an inflammatory process to organ malfunction [16], which causes the release of major pro-inflammatory cytokines such as IL-1 β and TNF- α . The present study showed that the administration of steeping black solo garlic significantly reduced levels of IL-1 β , IL-6 and TNF- α and increased IFN- γ in experimental animals.

Black solo garlic contains S-allyl-cysteine that has pharmacological effect as an antidiabetic, antioxidant, and anti-inflammatory with a higher bioactivity than regular types. Furthermore, the S-allyl-cysteine level with distilled water is higher than ethanol solvent [17]. Black solo garlic contains organosulphur compounds with potent antioxidant activity and free radical scavengers [18]. During inflammation, many pro-inflammatory mediators are generated and the administration of black solo garlic suppresses them to avoid tissue damage. Antioxidant compounds such as S-allyl cysteine, S-allyl mercaptocysteine, and allicin exhibit antioxidant activity and they inhibit inflammation by suppressing the activity of the NF- κ B signaling [19]. Meanwhile, S-allyl-cysteine has been proven to scavenged superoxide anions, hydrogen peroxide, hydroxyl radicals, peroxynitric radicals, and peroxyl radicals produced in neuron, as well as hypochloric acid and singlet oxygen in microglial cells.

Chemical compounds content of solo garlic is similar to ordinary type; however, some compounds have higher amounts such as flavonoids, total phenols, S-allyl-cysteine, and minerals [20]. Flavonoids can function as antioxidants to reduce or terminate free radical chain reactions [21]. S-allyl cysteine ameliorates cells by regulating peroxisomal proliferator activator receptor- α (PPAR- α), sterol regulatory element binding protein 1c (SREBP-1c), and decreasing levels of reactive oxygen species [22]. In addition, black solo garlic contains minerals such as Cu, Mn, and Zn, which have important role in the activity of oxidant enzymes such as superoxide dismutase. The antioxidant activity reduces reactive oxygen species and increases glutathione peroxidase, catalase, superoxide dismutase, reduced glutathione (GSH) and malondialdehyde [23, 24].

Black garlic suppresses the toll-like receptor 4 (TLR4) signals in macrophages. TLR4 acts to activate myeloid differentiation factor 88 (MyD88). It releases pro-inflammatory mediators such as IL-1 β , IL-6, and TNF- α [25]. These various mechanisms underlie several studies which explaining the role of black garlic in suppressing the formation of TNF- α , IL-1 β , and interferon- γ [26, 27]. Although BSG affected TLR4, the effect on NLRP as a component of inflammasome that have critical role in generating IL-1 β is interesting to be explored in the future study.

Declarations

Author contribution statement

Saryono: Conceived and designed the experiments; Performed the experiments; Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Desiyani Nani: Conceived and designed the experiments; Analyzed and interpreted the data.

Atikah Proverawati: Performed the experiments; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Sarmoko: Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Funding statement

This work was supported by LPPM Jenderal Soedirman University (No. T/324/UN23.18/PT.01.03/2020).

21

Data availability statement

Data will be made available on request.

Declaration of interests statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

Acknowledgements

The authors are grateful to Dr. Arif Fadlan for discussion on data analysis. The authors also acknowledge the support of Agil Patria Putra during preparing this manuscript.

References

- [1] Y. Sari, A.S. Upoyo, A. Isworo, A. Taufik, A. Sumeru, D. Anandari, et al., Foot self-care behavior and its predictors in diabetic patients in Indonesia, BMC Res. Notes 13 (2020) 38.
- [2] S. Tsalamandris, A.S. Antonopoulos, E. Oikonomou, G.-A. Papamikroulis, G. Vogiatzi, S. Papaioannou, et al., The role of inflammation in diabetes: current concepts and future perspectives, Eur. Cardiol. 14 (2019) 50–59.
- [3] G.K. Kolluru, S.C. Bir, C.G. Kevil, Endothelial dysfunction and diabetes: effects on angiogenesis, vascular remodeling, and wound healing, Int. J. Vasc. Med. 2012 (2012) 918267.
- [4] S. Mirza, M. Hossain, C. Mathews, P. Martinez, P. Pino, J.L. Gay, et al., Type 2-diabetes is associated with elevated levels of TNF- α and IL-6 and adiponectin and low levels of leptin in a population of Mexican Americans: a cross-sectional study, Cytokine 57 (2012) 136–142.
- [5] R. Kartika, D. Purnamasari, S. Pradipta, R.A. Larasati, H. Wibowo, Impact of low interferon- γ and IL-10 levels on TNF- α and IL-6 production by PHA-induced PBMCs in type 2 diabetes mellitus, J. Inflamm. Res. 13 (2020) 187–193.
- [6] A. Ortiz, C. Bustos, J. Alonso, R. Alcázar, M.J. López-Aranda, J.J. Plaza, et al., Involvement of tumor necrosis factor- α in the pathogenesis of experimental and human glomerulonephritis, Adv. Nephrol. Necker. Hosp. 24 (1995) 53–77.
- [7] Y.-C. Chang, L.-M. Chuang, The role of oxidative stress in the pathogenesis of type 2 diabetes: from molecular mechanism to clinical implication, Am. J. Transl. Res. 2 (2010) 316–331.
- [8] T.-C. Chang, H.-D. Jang, W.-D. Lin, Biochemical properties of black garlic aged under different temperatures of commercial rice wine extracts in Taiwan, Food Meas. 15 (2021) 509–518.
- [9] S. Kimura, Y.-C. Tung, M.-H. Pan, N.-W. Su, Y.-J. Lai, K.-C. Cheng, Black garlic: a critical review of its production, bioactivity, and application, J. Food Drug Anal. 25 (2017) 62–70.
- [10] N.B. Sembiring, Y. Iskandar, A review of component and pharmacology activities of black garlic, Trad. Med. J. 24 (2019) 178.
- [11] D. Wang, Y. Feng, J. Liu, J. Yan, M. Wang, J. Sasaki, et al., Black garlic (allium sativum) extracts enhance the immune system, Med. Aromat. Plant Sci. Biotechnol. (2010).
- [12] H. Lee, J.-W. Heo, A.-R. Kim, M. Kwon, S. Nam, J.-S. Lim, et al., Z-ajoene from crushed garlic alleviates cancer-induced skeletal muscle atrophy, Nutrients 11 (2019).
- [13] B. Zhai, C. Zhang, Y. Sheng, C. Zhao, X. He, W. Xu, et al., Hypoglycemic and hypolipidemic effect of S-allyl-cysteine sulfoxide (alliin) in DIO mice, Sci. Rep. 8 (2018) 3527.
- [14] Z. Qiu, Z. Zheng, B. Zhang, D. Sun-Waterhouse, X. Qiao, Formation, nutritional value, and enhancement of characteristic components in black garlic: a review for maximizing the goodness to humans, Compr. Rev. Food Sci. Food Saf. 19 (2020) 801–834.
- [15] G.-B. Tran, S.-M. Dam, N.-T.T. Le, Amelioration of single clove black garlic aqueous extract on dyslipidemia and hepatitis in chronic carbon tetrachloride intoxicated swiss albino mice, Int. J. Hepatol. 2018 (2018) 9383950.
- [16] M.L. Graham, J.L. Janecek, J.A. Kittredge, B.J. Hering, H.-J. Schuurman, The streptozotocin-induced diabetic nude mouse model: differences between animals from different sources, Comp. Med. 61 (2011) 356–360.
- [17] N.A. Thach, Effect of extraction conditions on polyphenols, flavonoids, s-allyl cysteine content and antioxidant activity of black garlic extracts, JST 55 (2018) 18.
- [18] A.Y. Nasr, Protective effect of aged garlic extract against the oxidative stress induced by cisplatin on blood cells parameters and hepatic antioxidant enzymes in rats, Toxicol. Rep. 1 (2014) 682–691.
- [19] G. Yang, S. Li, B. Li, L. Cheng, P. Jiang, Z. Tian, et al., Protective effects of garlic-derived S-allylmercaptocysteine on IL-1 β -stimulated chondrocytes by regulation of MMPs/TIMP-1 ratio and type II collagen expression via suppression of NF- κ B pathway, BioMed Res. Int. 2017 (2017) 8686207.
- [20] J.H. Ryu, D. Kang, Physicochemical properties, biological activity, health benefits, and general limitations of aged black garlic: a review, Molecules 22 (2017).

- [21] X. Lu, N. Li, X. Qiao, Z. Qiu, P. Liu, Composition analysis and antioxidant properties of black garlic extract, *J. Food Drug Anal.* 25 (2017) 340–349.
- [22] J.-C. Tsai, Y.-A. Chen, J.-T. Wu, K.-C. Cheng, P.-S. Lai, K.-F. Liu, et al., Extracts from fermented black garlic exhibit a hepatoprotective effect on acute hepatic injury, *Molecules* 24 (2019).
- [23] K.M. Naji, E.S. Al-Shaibani, F.A. Alhadi, S.A. Al-Soudi, M.R. D'souza, Hepatoprotective and antioxidant effects of single clove garlic against CCl₄-induced hepatic damage in rabbits, *BMC Compl. Alternative Med.* 17 (2017) 411.
- [24] H. Xiong, X. Jin, C. You, Expression of the CD59 glycoprotein precursor is upregulated in an estrogen receptor-alpha (ER- α)-Negative and a tamoxifen-resistant breast cancer cell line in vitro, *Med. Sci. Mon.* 24 (2018) 7883–7890.
- [25] B.R. You, J.-M. Yoo, S.Y. Baek, M.R. Kim, Anti-inflammatory effect of aged black garlic on 12-O-tetradecanoylphorbol-13-acetate-induced dermatitis in mice, *Nutr. Res. Pract.* 13 (2019) 189–195.
- [26] S.A. Almatroodi, S. Anwar, A. Almatroudi, A.A. Khan, F. Alrumaihi, M.A. Alsahli, et al., Hepatoprotective effects of garlic extract against carbon tetrachloride (CCl₄)-induced liver injury via modulation of antioxidant, anti-inflammatory activities and hepatocyte architecture, *Appl. Sci.* 10 (2020) 6200.
- [27] S. Saryono, A. Proverawati, *The Potency of Black Garlic as Anti-atherosclerotic: Mechanisms of Action and the Prospectively*, AIP Publishing, 2019, p. 20027.

Immunomodulatory effects of black solo garlic (*Allium sativum* L.) on streptozotocin-induced diabetes in Wistar rats

ORIGINALITY REPORT

12%
SIMILARITY INDEX

9%
INTERNET SOURCES

6%
PUBLICATIONS

2%
STUDENT PAPERS

PRIMARY SOURCES

1 repository.ubaya.ac.id 1 %
Internet Source

2 www.cetjournal.it 1 %
Internet Source

3 journals.lww.com 1 %
Internet Source

4 www.dovepress.com 1 %
Internet Source

5 html.rhhz.net <1 %
Internet Source

6 Submitted to Chester College of Higher Education <1 %
Student Paper

7 Rona Kartika, Dyah Purnamasari, Saraswati Pradipta, Rahma A Larasati, Heri Wibowo. " <1 %

Impact of Low Interferon- γ and IL-10 Levels on TNF- α and IL-6 Production by PHA-Induced PBMCs in Type 2 Diabetes Mellitus

8

Submitted to University of Mustansiriyah

Student Paper

<1 %

9

mdpi-res.com

Internet Source

<1 %

10

Flora, S.J.S.. "Prevention of arsenic-induced hepatic apoptosis by concomitant administration of garlic extracts in mice", Chemico-Biological Interactions, 20090212

Publication

<1 %

11

www.karger.com

Internet Source

<1 %

12

Hesham N. Mustafa. "The role of curcumin in streptozotocin-induced hepatic damage and the trans-differentiation of hepatic stellate cells", Tissue and Cell, 2016

Publication

<1 %

13

J. Puppini-Rontani, A.P.P. Fugolin, A.R. Costa, L. Correr-Sobrinho, C.S. Pfeifer. "In vitro performance of 2-step, total etch adhesives modified by thiourethane additives", International Journal of Adhesion and Adhesives, 2020

Publication

<1 %

14

akjournals.com

Internet Source

<1 %

15 pubag.nal.usda.gov <1 %
Internet Source

16 ri.conicet.gov.ar <1 %
Internet Source

17 ri.uaemex.mx <1 %
Internet Source

18 www.spandidos-publications.com <1 %
Internet Source

19 Araldi, Elisa, and Yajaira Suarez. "MicroRNAs as regulators of endothelial cell functions in cardiometabolic diseases", Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids, 2016.
Publication

20 Soodeh Karami, Shiva Shamshiri, Mohammad Abdollahi, Roja Rahimi. "An Evidence-based Review of Medicinal Plants used in Traditional Persian Medicine for Treatment of Osteoarthritis", Current Drug Discovery Technologies, 2021
Publication

21 downloads.hindawi.com <1 %
Internet Source

22 harvest.usask.ca <1 %
Internet Source

23

Internet Source

<1 %

24

mdpi.com

Internet Source

<1 %

25

wprim.whoocc.org.cn

Internet Source

<1 %

26

www.39kf.com

Internet Source

<1 %

27

www.elsevier.es

Internet Source

<1 %

28

www.nature.com

Internet Source

<1 %

29

www.tjnpr.org

Internet Source

<1 %

30

Orawan Wongmekiat, Edward J. Johns.
"Contribution of Endothelial Nitric Oxide
Synthase in the Blunted Renal Responses to
Volume Expansion in Diabetic Rats",
Experimental Physiology, 2001

Publication

<1 %

31

Jun Ho Kim, Su Hyun Yu, Yun Jeong Cho, Jeong
Hoon Pan et al. " Preparation of -Allylcysteine-
Enriched Black Garlic Juice and Its Antidiabetic
Effects in Streptozotocin-Induced Insulin-
Deficient Mice ", Journal of Agricultural and
Food Chemistry, 2017

<1 %

32

de Souza Bastos, Alliny, Dana T. Graves, Ana Paula de Melo Loureiro, Carlos Rossa Junior, Samia Cruz Tfaile Corbi, Fausto Frizzera, Raquel Mantuanelli Scarel-Caminaga, Niels Olsen Camara, Oelisoa M. Andriankaja, Meire I. Hiyane, and Silvana Regina Perez Orrico. "Diabetes and increased lipid peroxidation are associated with systemic inflammation even in well-controlled patients", Journal of Diabetes and its Complications, 2016.

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On