

# Renoprotective Effects of Lycopene in Tomato Extracts on Rat Exposed to Cadmium

*by* Hernayanti Hernayanti

---

**Submission date:** 05-Nov-2021 01:00PM (UTC+0700)

**Submission ID:** 1693677738

**File name:** rnayanti\_2020\_IOP\_Conf.\_Ser.\_Earth\_Environ.\_Sci.\_593\_012005.pdf (297.98K)

**Word count:** 2173

**Character count:** 10561

## 4 Renoprotective Effects of Lycopene in Tomato Extracts on Rat Exposed to Cadmium

Hernayanti<sup>1</sup>, S Santoso, S Lestari, L Prayogo, Kamsinah, and Rochmatino

Faculty of Biology, Jenderal Soedirman University, Purwokerto, Indonesia

E-mail : hernayantioentoro@gmail.com

**Abstract.** Cadmium is a heavy metal widely used in human life. When it enters the body, cadmium might bind to metallothionein protein and impair renal function. The renal function decrease usually is characterized by raise of  $\beta$ 2-microglobulin, creatinine, ureum and uric acid levels, which possibly be cured by lycopene, and antioxidant found in tomato. The objective of the study was to determine the effective lycopene dose as a renoprotective of Cd exposure. Twenty four Wistar rats used in this study were divided into six groups (four individuals in each group). Group I was the negative control, Group II was given 5.6 mg/kgBW CdSO<sub>4</sub> as a positive control. Group III, IV and V were given 5.6 mg/kgBW CdSO<sub>4</sub> and tomato extract of 0.36, 0.72 and 1.08 mg/kgBW/day respectively. Group VI exposed to CdSO<sub>4</sub>, and after seven days an additional 0.3 mg/kgBW/day of dimercaprol was administered for 14 days as a comparison to tomato. The parameters observed were levels of Cd, creatinine, and  $\beta$ 2-microglobulin in blood level. Observations were made on day 0 and day 22. The data were analyzed by ANOVA (F-test), followed by the Duncan test. The results showed that all treatments and dimercaprol could reduce the levels of Cd, creatinine, ureum, uric acid, and  $\beta$ 2-Microglobulin. The optimal decrease was observed in the rats administered with 1.08 mg/kgBW/day. Tomato extract dose 1.08 mg/kgBW was the highest in lowering levels of blood Cd,  $\beta$ 2-M, creatinine, urea, and uric acid levels.

### 1. Introduction

Industrial waste containing Cd can pollute the atmosphere, soil, and water. Cadmium is a persistent compound in an environment with a half-life of 30-40 years [1, 2]. Consequently, exposure to both acute and chronic Cd is very harmful to human health, especially to the kidney, which is the main target of Cd. Cadmium is also one of the causes of hypertension and human heart disease (atherosclerotic heart disease) [3, 4, 5]. Poisoning of Cd, Itai-Itai disease, occurring in Japan was caused by water pollution in Kumamoto. In Indonesia, it happened in Jakarta Bay. A Kaliadem resident who consumed green mussels had a high risk for Cd exposure (RQ > 1) with hypertension symptom [6]. In the body, cadmium binds to the metallothionein protein [7, 8]. This bonding is stable and can lead to increase free radicals in the liver and kidneys, resulting in oxidative stress characterized by decreased of superoxide dismutase (SOD) and Glutathione Peroxidase (GPx), leading to a decrease of renal function [3, 9]. An antioxidant supplement such as vitamin C, Vitamin E, and Selenium prevents Cd toxicity. They reduce Cd absorption by the kidneys and help to eliminate Cd out of the body without damaging the kidneys [10]. However, antioxidants in the form of food supplements are expensive and only the upper-middle-class society that might afford it. Usually, Cd toxicity treatment involves application of a chemical chelation compound, dimercaprol, but it is not



2  
Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

recommended because it can damage the kidneys and cause hypertension [3]. It is necessary to find an alternative treatment of easily available and cheap natural antioxidants such as tomatoes to overcome Cd toxicity. Tomato contains an active compound called lycopene, a potent antioxidant because of its capacity to reduce free radical compounds in the body [10]. This research aimed to understand renoprotective effects of lycopene on rat exposed by Cd with measuring of their  $\beta$ 2-M level, and renal function creatinine, urea, and uric acid level, as well as to find the effective dose of lycopene as renoprotective on rat exposed by Cd in terms of decreasing  $\beta$ 2-M, creatinine, urea, and uric acid level.

## 2. Methods

We selected a golden jubilee or golden tomato. The ethanol tomato extract (by maceration method) was used as a chelating agent for Cd poisoning. Five kg of raw tomato was dissolved in 96%-ethanol for 3x24 hours, and to thicken the extract the macerate was evaporated by vacuum rotary evaporator to produce 0.5 g extract. Tomato extract was administered 14 days after rat exposures to CdSO<sub>4</sub>. Wistar rats were fed with AD II pellets and distilled water for drinking ad libitum. They (24) were divided into six groups, C1: negative control (non-CdSO<sub>4</sub>, non-tomato extract), C2: positive control (exposed to 5.6 mg/kgBW CdSO<sub>4</sub> for 14 days), C3, C4, C5 were exposed to 5.6 mg/kgBW CdSO<sub>4</sub> for seven days and given tomato extract at a dose of 0.36, 0.72 and 1.08 mg/kgBW/day for 14 days, respectively. C6 was exposed to CdSO<sub>4</sub> for seven days and dimercaprol for 14 days at a dose of 0.3 mg/kgBW. Dose of 0.36 mg/kgBW equals to lycopene dose of 15 mg/kgBW in humans, 0.72 mg/kgBW correspond to 30 mg/kgBW in human, and 1.08 mg/kgBW was to 45 mg/kgBW. The blood was collected on day 0 and 15, using a hematocrit capillary pipette on the vein orbitalis plexus of rat, then collected in Eppendorf tube (3 ml). The blood was divided into two parts, i.e., 0.5 ml for blood Cd analysis, and 2.5 ml for measurements of creatinine, uric acid, urea, and  $\beta$ 2-M. The 2.5 ml blood was centrifuged for 10 minutes (4,000 rpm). Cadmium was measured by AAS at 228.6 nm wavelength and a strong current of 3.5 mA0. Creatinine level was examined by Jaffe kinetic method and read at spectrophotometer (492 nm). Urea and uric acid were measured by Dyas method with a wavelength of 546 nm. Data of Cd, creatinine, uric acid, urea, and  $\beta$ 2-M levels were analyzed by ANOVA, followed by Duncan test, to find out the differences in the treatments.

## 3. Results

The levels of cadmium, creatinine,  $\beta$  2M, urea and uric acid after administrated by lycopene is presented in Table 1.

**Table 1.** Cadmium, creatinine,  $\beta$  2M, urea and uric acid levels after lycopene administered (C1: healthy control, C2: CdSO<sub>4</sub> + 5.6 mg/kgBW lycopene, C3: CdSO<sub>4</sub> + 0.36 mg/kgBW lycopene, C4: CdSO<sub>4</sub> + 0.72 mg/kgBW lycopene, C5: CdSO<sub>4</sub> + 1.08 mg/kgBW lycopene, C6: CdSO<sub>4</sub> + 0.3 mg/kgBW dimercaprol. Column followed by the same letter is not significantly different with  $p < 0.05$ .)

Treatment	Cadmium (ppm)	Creatinin (mg/dl)	$\beta$ 2M (mg/dl)	Urea (mg/dl)	Uric acid (mg/dl)
C1	0.2 ± 0.08 <sup>a</sup>	0.61 ± 0.09 <sup>a</sup>	96.21 ± 16.98 <sup>a</sup>	13.5 ± 1.62 <sup>a</sup>	2.63 ± 0.12 <sup>a</sup>
C2	2.08 ± 0.17 <sup>b</sup>	1.23 ± 0.16 <sup>b</sup>	188.15 ± 17.91 <sup>b</sup>	45.38 ± 2.97 <sup>b</sup>	6.17 ± 0.56 <sup>b</sup>
C3	1.36 ± 0.04 <sup>c</sup>	1.00 ± 0.20 <sup>c</sup>	140.38 ± 28.07 <sup>c</sup>	28.75 ± 2.01 <sup>c</sup>	4.67 ± 0.36 <sup>c</sup>
C4	1.18 ± 0.36 <sup>c</sup>	0.78 ± 0.11 <sup>a</sup>	111.05 ± 10.04 <sup>d</sup>	17.50 ± 1.01 <sup>d</sup>	2.92 ± 0.20 <sup>a</sup>
C5	0.84 ± 0.05 <sup>d</sup>	0.69 ± 0.08 <sup>a</sup>	84.58 ± 14.48 <sup>a</sup>	19.70 ± 1.05 <sup>d</sup>	3.05 ± 0.19 <sup>d</sup>

Treatment	Cadmium (ppm)	Creatinin (mg/dl)	<sup>8</sup> $\beta$ 2M (mg/dl)	Urea (mg/dl)	Uric acid (mg/dl)
C6	1.14±0.40 <sup>c</sup>	0.82±0.05 <sup>a</sup>	113.6±15.22 <sup>d</sup>	22.32±1.15 <sup>d</sup>	3.23±1.2 <sup>d</sup>

#### 4. Discussion

The highest Cd level was in C2 after giving the rat with CdSO<sub>4</sub>. It is because Cd in the body binds to metallothionein (Cd + Mt) and increase Reactive Oxygen Species (ROS) such as 1O<sub>2</sub>, O<sub>2</sub><sup>-</sup> and OH<sup>-</sup> which leads to lipid peroxidation that accumulates and damages the kidney especially renal proximal tubule [7, 8]. The damage of proximal renal tubules by Cd leads to an increase in blood creatinine, urea, uric acid, and  $\beta$ 2-M level (Table 1) [11, 12].

The  $\beta$ 2-M increase in the rats (normal value 80-150 ng/ml) is due to renal dysfunction and inhibits salt reabsorption, reduction of water reabsorption, and consequently an increase in urine volume (polyuria) [13]. Creatinine is a creatine and phosphocreatine metabolites which are filtered in the glomerulus and reabsorbed in the kidney tubules. Kidney dysfunction causes Glomerular Filtration Rate (GFR) to decrease, followed by decrease ability to filtrate creatinine, and increase of serum creatinine (normal value 0.3-0.9 mg/dl) [14].

After administering the tomato extract and dimercaprol for 14 days, improvements were shown in renal function characterized by decreased levels of blood Cd, creatinine urea, uric acid, and  $\beta$ 2-M, which returned to normal levels. The analysis showed highly significant differences between control and treatment groups. Lycopene and flavonoids in tomatoes can neutralize free radical of Cd +Mt by giving H<sup>+</sup> as an electron donor, result in improvements of kidney organ. Lycopene in tomato fruit may reduce free radicals 20 times greater than vitamin C, and 10 times larger than vitamin E [10, 15]. Lycopene in tomato extract dose of 1.08 mg/kgBW was the best in lowering levels of blood Cd,  $\beta$ 2-M, creatinine, urea, and uric acid levels compared to dose of 0.36 mg/kgBW, 0.72 mg/kgBW, and dimercaprol dose of 0.3 mg/kgBW.

#### 5. Conclusion

Tomato extract with dose of 1.08 mg/kgBW was the best in lowering levels of blood Cd,  $\beta$ 2-M, creatinine, urea, and uric acid levels.

#### 6. Acknowledgment

We thank the Institute for Research and Community Service, Universitas Jenderal Soedirman for the financial support to which this research was completed.

#### References

- [1] Jacques P F, Lyass A, Massaro J M, Vasan R S and D'Agostino R B 2013 Relationship of lycopene intake and consumption of tomato products to incident CVD *British Journal Nutrition* **110** 545–51
- [2] Satarug S and Garret S H 2010 cadmium, environmental, exposure and health outcomes *Environ Health Perspect.* **118** 182–90
- [3] Caciari T, Sancini A, Fioravanti M, Cappozella A and Casale T 2013 Cadmium and hypertension in exposed workers: a meta-analysis *Int J Occup Med Environ Health* **26** 440–56
- [4] Gallagher C M and Meliker J R 2010 Blood and urine cadmium, blood pressure and hypertension: a systemic review and meta-analysis *Environ Health Perspect* **118** 1676–84
- [5] Messner B and Bernhard D 2010 Cadmium and cardiovascular disease: cell biology, pathophysiology and epidemiological relevance *Biometals* **23** 811–22

- [6] Safitri F Z 2015 Level of environmental health effects contents of heavy metal cadmium (Cd) in green mussels (*Perna viridis*), which consumed by Kaliadem Muara Angke people, North Jakarta *Skripsi* (Jakarta: UIN Syarif Hidayatullah)
- [7] Chen L, Jin T, Huang B, Norberg G and Norberg M 2006 Critical exposure level of Cd for elevated urinary methallothionein an occupational population study in China *Toxicol Appl Pharmacol.* **215** 93–9
- [8] Cho M R, Kang H W, Jeon S H and Cho M H 2010 Time dependent changes of cadmium and methallothionein after short-term exposure to cadmium in rats *Toxicol Res* **26** 131–6
- [9] Hijova E, Nistiar F and Kuchta M 2004 Influence of acute cadmium exposure on plasma antioxidant parameters in rats *Bulletin of the Veterinary Reseach Institute Pulawy* **48** 155–7
- [10] Agarwal S and Rao A 2000 Tomato lycopene and its role in human health and chronic diseases *Journal of Canadian Medical Association* **163** 739–44
- [11] Bernhoft R A 2013 Cadmium toxicity and treatment *The Scientific World Journal* **13** 1–7
- [12] Johnson D W, Graham R D J, Timothy H M, Marie J L and Matthew P D 2012 Chronic kidney disease and automatic reporting of estimated glomerular filtration rate: new developments and revised recommendations *Med J Aust* **197** 222–3
- [13] Li L, Dong M and Wang X G 2016 The implication and significance of beta-2 microglobulin: a conservative multifunctional regulator *Chin Med J* **129** 448–54
- [14] Derelanko M J 2000 *Toxicologist's Pocket Handbook* (New York Washington, D.C: CRC Press. Boca Raton London) p 98
- [15] Holzapfel N P, Holzapfel B M, Champ S, Feldhusen J, Clement J and Huttmacher D W 2013 The potential role of lycopene for the prevention and therapy of prostate cancer: from molecular mechanisms to clinical evidence *Int J Mol Sci.* **14** 14620–46

# Renoprotective Effects of Lycopene in Tomato Extracts on Rat Exposed to Cadmium

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

14%

INTERNET SOURCES

9%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.journal.atmph-specialissues.org](http://www.journal.atmph-specialissues.org)

Internet Source

4%

2

[eprints.undip.ac.id](http://eprints.undip.ac.id)

Internet Source

3%

3

Hasnadhiazahra Rohadi, Muhammad Ilyas, Nuraeni Ekowati. "Preservation Technique of Filamentous Fungi Based on Inactive Metabolism at Indonesian Culture Collection (InaCC)", IOP Conference Series: Earth and Environmental Science, 2020

Publication

2%

4

[seminar.bio.unsoed.ac.id](http://seminar.bio.unsoed.ac.id)

Internet Source

2%

5

[beta.space.iqnito.com](http://beta.space.iqnito.com)

Internet Source

2%

6

[mro.massey.ac.nz](http://mro.massey.ac.nz)

Internet Source

1%

7

AS Piranti, DRUS Rahayu, ER Ardli, N Setyaningrum, DS Widyartini, I Insan. "Water

1%

Quality Status of Segara Anakan Cilacap  
Indonesia for Biota Life", IOP Conference  
Series: Earth and Environmental Science, 2020

Publication

8

[www.e-sc.org](http://www.e-sc.org)

Internet Source

1 %

9

Muslihudin Muslihudin, Tyas Retno Wulan, Tri Sugiarto, Sotyania Wardhianna, Sri Wijayanti. "Migrant Workers Empowerment through Productive Migrant Village Programs in Banyumas, Indonesia", Society, 2021

Publication

1 %

10

[bionursing.fikes.unsoed.ac.id](http://bionursing.fikes.unsoed.ac.id)

Internet Source

1 %

11

Gheorghe Coman, Camelia Draghici. "Chapter 3 Heavy Metals Activity Mechanisms at Cellular Level and Possible Action on Childrens Bodies", Springer Science and Business Media LLC, 2011

Publication

<1 %

12

Sehrawat, A.. "Evaluation of possible mechanisms of protective role of Tamarix gallica against DEN initiated and 2-AAF promoted hepatocarcinogenesis in male Wistar rats", Life Sciences, 20060905

Publication

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On