

Turn_Rimbawanto_2019

by Efka Aris

Submission date: 28-Mar-2023 08:21PM (UTC+0700)

Submission ID: 2049012014

File name: 30._Artikel_58183-178400-2-PB_1.docx (40.13K)

Word count: 1844

Character count: 9644

Effect of urea-cassava waste and protected soybean meal supplementation on dry matter, crude fiber, and crude protein intake of ram

D N Cahyo¹, R Nurvaif², B Hartoyo³, E A Rimbawanto³

Postgraduate Student of Faculty of Animal Science, Gadjah Mada University,
Yogyakarta, Indonesia

² **3**. Charoen Pokphand Indonesia Breeding Farm Banten

³ Faculty of Animal Science, Jenderal Soedirman University, Purwokerto, Indonesia

Corresponding author: fk.aris.r@gmail.com

Abstract. The purpose of this research is to measure the effect of urea-cassava waste **10** *Leucaena leucocephala* tannin extract protected soybean meal supplementation in ration on dry matter (DM), crude fiber (CF), and crude protein (CP) intake of ram and to determine the best supplementation level based on these three variables. This research was **4** held from October 18⁰¹, 2017 until January 10⁰¹, 2018 at Experimental Farm and Animal Feed and Nutrition Laboratory, Faculty of Animal Science, Jenderal Soedirman University, Purwokerto. The material used in this research were \pm 12 months old ram (initial weight 23-25 kg) as many as 15 heads, control ration (rice straw: concentrate = 30:70), supplement (urea-cassava waste : protected soybean meal = 54 : 46%), and experimental cage. The method of this research was *in vivo* experiment with completely randomized design followed by orthogonal polynomial if the treatment had significant effect. Treatments tested in this research was the percentage of supplementation that consist of Ro (control), (Ro + 5% DM supplement), R2 (R0+10% DM supplement), **23** (R0+15% DM supplement), and R4 (R0+20% DM supplement) with three replications. The result of ANOVA showed **12** that supplementation of urea cassava waste and protected soybean meal in male sheep ration had no significant effect ($P>0,05$) on male sheep DM and CF intake but had very significant effect ($P<0,01$) on CP intake with linear shaped curve ($Y=113,005 + 1,86X$). The conclusion of this research is supplementation of urea-cassava waste and protected soybean meal as much as 20% DM in ration did not affect DM and CF intake and could optimize CP intake of ram.

1. Introduction

Cut and carry feeding system, which implemented in major lamb farms in Indonesia cannot fulfill the energy and protein needs of fattening ram. Supplementing protein-ingredients to ruminants is also not a beneficial, because no more than 40 % of the protein escapes the rumen degradation [1]. Therefore, it is necessary to protect protein-rich ingredients to supply the amino acids for the post rumen tact. Condensed tannins of *Leucaena leucocephala* leaf can be used as natural protectants of protein-rich ingredients as substituents of formaldehyde [2]. Protection can increase the protein supply of the post-rumen gastrointestinal tract, but can reduce the concentration of rumen N-ammonia, which results in a reduction in microbial protein synthesis and rumen microbial activity *in vitro* [3].

12
Urea can be used as a source of crude protein for ruminants, but its rapid degradation to NH₃ is faster than NH₃ utilization by rumen microbial. Slow released urea can be achieved by mixing it with cassava waste through an extrusion process. The mixture of urea-cassava waste can inhibit the hydrolysis of nitrogen-urea in the rumen up to 14 hours in vitro. The combination of condensed tannin-protected soybean meal and slow release urea (urea-cassava waste) can be combined with a ratio of 46:54% DM and supplemented in low-quality control ration to increase microbial protein synthesis and rumen fermentation products in vitro [4]. The *in vivo* study of this supplements has never been done before. Therefore, this study was aimed to find out how they affect dry matter, crude fiber, and protein intake, and the best percentage of supplements based on these variables.

2. Material and methods

The materials used in this study, were \pm 12 months old (23-25 kg) as many as 15 heads, control rations of rice straw and concentrates (30: 70%), concentrate were used in this study consists of 41% cassava waste, 50% pollard, 5% corn flour, 2% soybean meal, 1% mineral mix, and 1% NaCl, urea-cassava waste supplements and protected soybean meal (54: 46%), a set of experimental cages, as well as a set of dry matter, crude fiber, and crude protein analysis tools [5]. The treatments ration (R0,1,2,3,4) contain at least 76,07; 80,40; 84,72; 89,05; 93,38% DM, 27,25; 28,19; 29,13; 30,06; 31,00% CF, 9,56; 11,48; 13,39; 15,30; 17,21% CP, and 58,71; 62,36; 66,02; 69,67; 73,32% TDN respectively.

Urea-cassava waste (slow released urea) is produced by steaming 2 kg of cassava waste with urea solution (200 g of urea dissolved in 750 ml of water) using autoclave for 15 minutes. The steaming process in the autoclave was aimed to process starch gelatinization and starch-nitrogen binding. Soybean meal is protected using *L. leucocephala* crude tannin extract by spraying method. As many as 1.68 g of crude tannin extract was used to protect 100 g of soybean meal. Crude tannin extract is obtained by extracting 20 g of *L. leucocephala* leaf flour with a solvent consisting of 70% ethanol, 30% aquadest, and 1% HCl as much as 600 ml. Extraction was carried out using a rotary evaporator for 2 hours 30 minutes at a temperature of $< 80^{\circ}\text{C}$ to prevent component degradation.

The research method was an *in vivo* experiment using a complete randomized design (CRD) with 5 treatments and 3 replications. The variables observed in this study were dry matter (DM), crude fiber (CF) and crude protein (CP) intake. Data then analyzed using analysis of variance (ANOVA), if the treatment has a significant effect, analysis followed by orthogonal polynomial tests to determine the best or optimum percentage of supplementation [6].

3. Result and Discussion

6
Dry matter, crude fiber, and crude protein intake are shown in Table 1. The result showed that there was no significant effect of supplementation urea-cassava waste and protected soybean meal in DM intake ($P>0,05$). It indicates that the supplementation did not interfere microbial protein synthesis and nutrient digestibility of ram. [2] reported that tannin content in ruminant diet can reduce the concentration of rumen N-ammonia, but in this study, CP in the rumen supplied by urea-cassava waste (slow release urea) to kept N-ammonia concentration in rumen remain stable. This result is in line with the study of [7-8] that stated the usage of slow released urea can provide N-ammonia for microbial protein synthesis, then the microbes increase nutrient digestibility and digestion rate. So that, the tannin content in the diet did not affect DM intake.

Supplementation did not give significant effect on crude fiber intake of ram ($P>0,05$). CF was the highest constituent of ruminant DM diets, so the intake was depending on DM intake. [9] stated that tannin content in diet could reduce fiber digestability, but [10] reported that addition of tannin containing *T. chebula* as much as 10g/kg DM *T. chebula* increase fiber intake and digestibility. The treatment given in this study was supplementation of CP source ingredients, so that the treatment did not affect the CF intake. This is in line with the statement of [11] which states that feed intake is influenced by its nutrient content.

The result showed that the treatment had a very significant effect ($P < 0.01$) on CP intake. The supplementation treatment had an effect on increasing CP intake of rams with R^2 value=56.75%. The

results of further CP intake shown a linear curve followed the equation $Y = 113.005 + 1.86X$ (Figure 1). The higher the percentage of supplements given, the higher the ram CP intake. [12] reported that the supplementation of CP from soybean meal with 0.09% of body weight in castrated cattle significantly increased the CP intake of *Kentucky bluegrass* straw and *Poa pratensis* (C3) based control ration, and *tallgrass prairie*. [13] stated that the higher CP content in the diet increased the CP intake, although DM and OM intake remain stable.

Table 1. Dry matter, crude fiber, and crude protein intake of diet supplemented by urea-cassava waste and protected soybean meal

Treatments	DM IntakeNs	CF IntakeNs	CP Intake**
RO	897,660 ± 81,93	228,03±23,18	104,79±7,54
R1	888,750 ± 201,53	220,68±51,75	129,82±19,07
R2	977,380 ± 57,89	245,01±25,89	140,37±6,88
R3	867,250 ± 44,14	213,50±7,52	134,23±8,50
R4	820,920 ± 103,27	199,29±6,00	149,09±8,87
Average	890,392 ± 97,75	221,30±28,56	131,66±18,07

ns, not significant; **) significant at 1%




Figure 1. The effect of urea-cassava waste and protected soybean meal supplementation on CP intake

4. Conclusion

Supplementation of slow release urea (urea-cassava waste) and *L. leucocephala* leaf condensed tannin-protected soybean meal with a level of 20% DM in ration rams did not interfere with the intake of dry matter and crude fiber, and could optimize the crude protein intake of the rams.

References

- [1] Hristov A N, A Bannink, L A Crompton, P Huhtanen, M Kreuzer, M McGee, P Noziere, C K Reynolds, A R Bayat, D R Yanez-Ruiz, J Dijkstra, E Kebreab, A Schwarm, K J Shingfield, and Z Yu 2018 *J Dairy Sci.* 102 5811-585
- [2] Rimbawanto E A, L M Yusiati, E Baliarti and R Utomo 2015 *J Anim. Product.* 17(2) 83-91
- [3] Cortes J E, B Moreno, M L Pabon, P Avila, M Kreuzer, H D Hess and J E Carulla 2009 *Anim. Feed Sci. Technol.* 151 194-204
- [4] Rimbawanto E A, S Suhermiyati and B Hartoyo 2016 *Efektivitas Penghambatan Hidrolisis Campuran Cassava waste dan Urea, Tepung Tapioka dan Urea, Tepung Gaplek dan Urea dalam Fermentasi Rumen secara In Vitro* In: Prosiding Seminar Nasional: Pengembangan

- Sumber Daya Pedesaan dan Kearifan Lokal Berkelanjutan VI. Purwokerto, November 24-25. LPPM Unsoed. Purwokerto. p. 900-906
- [5] AOAC 2002 *Official Methods of Analysis. Association of Official Agricultural Chemists. Agricultural Chemical; Contaminants; Drugs Vol. 1.* (Virginia: Association of Official Agricultural Chemists, Inc)
- [6] Steel R G D and J H Tome 1980 *Principles and Procedures of Statistics* (New York: Mc. Graw Hill Company Inc)
- [7] Goncalves A P, C F M do Nascimento, F A Ferreira, R C Gomes, M Manella, C T Marino, J J A A Demarchi and P H M Rodrigues 2015 *Braz. Arch. Biol. Technol.* 58(1) 22-30
- [8] Wanapat M, K Boonnop, C Promkot and A Cherdthong 2011 *Maejo Int. J. Sci. Technol.* 5(1) 13-23
- [9] Al-Dobaib S N 2009. *J. Anim. Sci.* 80 532-41
- [10] Patra A K, D N Kamra, N Agarwal and P N Chatterjee 2008 *Indian J. Anim. Nutr.* 24 251-5
- [11] Pech-Cervantes, A A, J Ventura-Cordero, C M Capetillo-Leal, J F J Torres-Acosta and C A Sandoval-Castro 2016 *J BSE.* 68 101-8
- [12] Bohnert D W, T DelCurto, A A Clark, M L Merrill, S J Falck and D L Harmon 2011 *J. Anim. Sci.* 89 3707-17
- [13] Lee C, A N Hristov, K S Heyler, T W Cassidy, M Long, B A Corl and S K R Karnati 2011 *J. Dairy Sci.* 94 5544-57

ORIGINALITY REPORT

10%

SIMILARITY INDEX

7%

INTERNET SOURCES

5%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | Ali Agus, Agussalim Agussalim. "chapter 4 Approach for the Domestication and Propagation of Stingless Bees", IGI Global, 2022
Publication | 1 % |
| 2 | Submitted to Universitas Jenderal Soedirman
Student Paper | 1 % |
| 3 | media.neliti.com
Internet Source | 1 % |
| 4 | repository.unsoed.ac.id
Internet Source | 1 % |
| 5 | www.granthaalayahpublication.org
Internet Source | 1 % |
| 6 | Lies Mira Yusiati, Ismaya, Zaenal Bachruddin, Prastisa Devi Parawansa. "Chapter 39 The Effect of Vitamin E Addition in the Diet on Blood Profile of Female Bligon Goat", Springer Science and Business Media LLC, 2017
Publication | 1 % |
| 7 | garuda.kemdikbud.go.id | |

Internet Source

1 %

8

worldwidescience.org

Internet Source

1 %

9

jurnal.unsyiah.ac.id

Internet Source

1 %

10

www.mdpi.com

Internet Source

1 %

11

www.sciencegate.app

Internet Source

1 %

12

Donielle Pannell, Brou Kouakou, Thomas H. Terrill, Ibukun M. Ogunade et al. "Adding dried distillers grains with solubles influences the rumen microbiome of meat goats fed lespedeza or alfalfa-based diets", Small Ruminant Research, 2022

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On