

Isolation and Characterization of Rhizospheric Bacteria Associated with Canna Plant for Production of Maltooligosaccharide Amylase

by Oedjijono Oedjijono

Submission date: 30-May-2023 09:17AM (UTC+0700)

Submission ID: 2104920256

File name: document.pdf (942.16K)

Word count: 6626

Character count: 36100

Research Article

54

Isolation and Characterization of Rhizospheric Bacteria Associated with Canna Plant for Production of Maltooligosaccharide Amylase

^{1,5} Dwi Agustiani^{1,2}, Oedjijono^{1*}, Nanik Rahmani², Nuraeni Ekowati¹

¹Faculty of Biology, University of Jenderal Soedirman, Jalan dr. Soeparno 63 Purwokerto 53122, Indonesia.

²Center for Applied Microbiology Research, Research Institute for Life and Environmental Sciences, National Research and Innovation Agency (BRIN), Cibinong, Bogor 16911, Indonesia.

³Department of Biology, Faculty of Science and Technology, International Women University, Bandung 40173, Indonesia.

* Corresponding author, email: oedjijono@unsoed.ac.id

Keywords:

Amylolytic bacteria
Canna
Maltooligosaccharides
16S rDNA gene
Submitted:
13 October 2022
Accepted:
30 January 2023
Published:
26 May 2023
Editor:
Miftahul Ilmi

ABSTRACT

The objectives of the study were to isolate amylolytic bacteria from the rhizosphere and plant tissue of *Canna edulis* Ker., as well as litter; to know oligosaccharide compounds produced from starch hydrolyzed by the bacterial enzymes, and to identify the amylolytic bacteria based on phenetic and 16S rRNA gene sequences. From the rhizosphere, Canna plant tissue, and litters obtained thirty-two amylolytic bacterial isolates. Eight isolates (TH6, TH7, T5, T10, D2, D3, A3, S1) produced high clear zone diameters ranging from 18–30 mm; especially an isolate T10, which was consistent in producing a total clear zone diameter of 20 mm. The hydrolysate of starch hydrolysed by the T10 amylase resulted in three oligosaccharide compounds maltotriose, maltotetraose, and maltopentose. The amylase activity of isolate T10 was optimal at a temperature of 40°C and pH at 0.801 U/mL. The isolate T10 was identified as a species member of *Bacillus toyonensis* based on phenotypic characterization and 16S rDNA gene sequencing analysis with a similarity value of 99.93%.

Copyright: © 2023, J. Tropical Biodiversity Biotechnology (CC BY-SA 4.0)

INTRODUCTION

Oligosaccharides are members of an essential group of carbohydrates. Macromolecules with short-chain polysaccharide sugars of 2 to 20 saccharide units. Functional oligosaccharides such as galactooligosaccharide (GOS), fructooligosaccharide (FOS), and maltooligosaccharide (MOS) are well-known prebiotics owing to their ability to selectively stimulate beneficial bacteria in the intestines, particularly bifidobacterial species (Zhao et al. 2017). Developing oligosaccharide products is one of the businesses with high economic value. Plants such as Canna contain much starch, one of the crucial ingredients (substrate) to produce oligosaccharides enzymatically. Canna plants (*Canna edulis* Ker.) contain high levels of carbohydrates, mainly starch (93.3%), which consists of amylose (33.48%) and amylopectin (59.82%) (El-Fakhry et al. 2012). Starch is hydrolysed into smaller oligosaccharides by α -amylase, one of the most important commercial enzymes (Jang et al. 2020).

The starch-processing industry has exploited amylase as a substitute for acid hydrolysis in producing starch hydrolysis. Amylase acts as a biocatalyst for the hydrolysis of starch into simpler carbohydrates, such

as glucose, maltose, and dextrin (Divakaran et al. 2011; Abdalla et al. 2021). Amylolytic bacteria are producers of amylase that can be used as biocatalysts in the starch hydrolysis process (Ding et al. 2021) to produce various maltooligosaccharide products, such as maltotriose, maltotetraose, maltopentaose, and maltohexaose (Pan et al. 2017).

Canna plants and its surrounding, including the rhizosphere and plant tissues, can be sources of isolating amylolytic bacteria. The high starch content in canna tubers makes them a suitable substrate for growing various bacteria, especially amylolytic bacteria. The bacteria isolated from starch-rich sources generally have the potential to produce amylase with high activity (Hellmuth & van den Brink 2013). In addition, the rhizosphere is known as the most diverse microbial habitat concerning species richness and community size. The interaction between plant roots and microorganisms is intensive around the rhizosphere, because the plants secrete exudates containing carbohydrates, amino acids, and other nutrients utilized by bacteria for growth. On the contrary, rhizospheric bacteria can produce proteins and enzymes that are important for the biological function of host plants (Afifah et al. 2018).

Bacteria, fungi, plants, and animals play an important role in utilizing polysaccharides. Members of the genus *Bacillus* were known to produce various enzymes, such as amylase that have been used in many industries, such as fermentation, textiles, paper, medicine, and sugar (Gupta et al. 2003). They are derived mainly from *Bacillus licheniformis* and *B. amyloliquefaciens*. Moradi et al. (2014) found several bacterial isolates producing high amylolytic enzymes, which were subsequently identified as *Bacillus cereus*, *B. amyloliquefaciens*, *B. licheniformis*, and *Paenibacillus lautus*. Luo et al. (2021) isolated *Bacillus toyonensis* P18, a group of Gram-positive bacteria belonging to the *Bacillus cereus* group and often used as probiotics or biocontrol agents. The bacterium has also been known to be treated as a probiotic for preventing microbial diseases in crops or improving the immune response of animals (Santos et al. 2018).

The objectives of the study were to isolate amylolytic bacteria from rhizosphere and plant tissue of Canna, as well as litter; to know oligosaccharide compounds produced from hydrolysate of starch hydrolyzed by the bacterial enzymes; and to identify the selected amylolytic bacteria based on 16S rRNA gene sequences.

MATERIALS AND METHODS

Sample Collection and Location of Sampling

Samples were taken from the rhizosphere and parts of Canna plant (*C. edulis* Ker.) including tubers, stems, leaves, tissue, as well as litter growing in two places, namely in the forest and the community gardens around the Perhutani Forest West Banyumas, Central Java, Indonesia. The coordinates of the former are S 07°20.846 'E 109°06.410 and the latter is S 07°20.812 'E 109°05.92 (Figure 1).

Isolation, Screening, and Morphological Characterization of Amylolytic Bacteria

Plant tissues and litter were cleaned with running water, then cut into 1 cm long pieces and separated according to the plant part. The sample pieces were immersed in 70% alcohol for 1 minute, then in 1% sodium hypochlorite solution for 3 minutes, after which they were soaked again using 70% alcohol for 1 minute, and rinsed with sterile distilled water three times (Duan et al. 2021, with modification). The sterile samples were placed on sterile tissue papers and then crushed using a mortar and one gram of each sample was diluted with 9 mL of sterile distilled water,




Figure 1. A map of Banyumas Regency and sampling sites: (1) the Perhutani Forest KPH West Banyumas, Central Java, (2) Community gardens around the Perhutani Forest KPH West Banyumas, Central Java.

and then serial dilutions were made up to 10^{-7} .

One gram of *Canna* rhizospheric soil was put into a 20 mL of nutrient broth (NB) medium containing 1% soluble starch (Merck) in a 100 mL Erlenmeyer flask. The solution was then homogenized in an agitation speed shaker machine at 150 rpm and incubated for 26 hours at 30 °C. The amount of 1 mL of the solution was diluted with 9 mL of sterile distilled water, and then serial dilutions were made up to 10^{-7} .

One mL from each series of dilutions was inoculated onto nutrient Agar (NB medium containing 1% soluble starch using a pour plate method. The plates were then incubated for 24 hours at 30 °C. Each growing bacterial colony was then inoculated onto an NA medium containing 1% soluble starch and purified using a streak quadrant method.

The number of 0.5L of bacterial cultures aged 24 hours at 30°C growing on NB medium containing 1% soluble starch was spot inoculated onto NA medium containing 1% commercial soluble starch, then incubated for 72 hours at 30°C. The growing colonies were flooded with iodine solution, and the clear zones formed around the colonies were observed and measured in their diameter (Vijayalakshmi et al. 2012). The isolates having high diameter clear zones were selected and tested for their enzyme production. The colonial and cell morphology of the isolated bacteria were characterized using conventional methods (Smibert & Krieg 1981).

Phenetic and Phylogenetic Characterizations of the Selected Bacterium

Phenetic characterizations of the selected bacterium (producing high diameter clear zones and maltoligosaccharide enzyme) including colony morphology, cell morphology, and biochemistry, were conducted by conventional methods (Smibert & Krieg 1981). Biochemical tests were also conducted using the API 50CHB kit.

The 16S rDNA gene was amplified by polymerase chain reaction (PCR) technique using a pair of primers (9F: 5'GAGTTT-GATCCTCCTGGCTCAG-3' 1510R: 5'GGCTACCTTGTTACGA-3')

(Yopi et al. 2017). The obtained bands were stained and visualized by UV Transilluminator. The sequence was confirmed via 1st BASE Sequencing, Mala⁴⁸a. The 16S rDNA nucleotide sequences were analyzed by nucleotide BLA⁴⁵ (Basic Local Alignment Search Tool) search in the Gene Bank of National Center for Biotechnology Inform¹⁹on (NCBI) or BLAST for amino acid analysis (Zhuang et al. 2012). The phylogenetic tree was constructed using a neighbour-joining algorithm in MEGA 6.0 software (De-Moraes-Russo & Selvatti 2018).

Analysis of The Hydrolysis Products by A Thin-Layer Chromatography Method (Rahmani et al. 2013)

An am¹⁰nt of 2 mL of each 24 h old bacterial culture (four selected isolates) was inoculated into a 200 mL flask containing 20 mL of NB medium plus 1% (w/v) starch solution, pH 6 (50 mM acetate buffer) and incubated at 30 °C for 24 h. The culture was sampling every 24 hour and then centrifuged, and the supernatant obtained was tested for its amylolytic activity.

Th⁴⁰ hydrolytic activity of amylase in a substrate solution was carried out at 30°C in 50 mM acetate buffer, pH 6, containing 0.5% of commercial starch. The enzyme-substrate ratio (v/v)⁹ was 1:1 and the reaction times were in hours (0, 1, 2, 3, 4, 24). Reactions were carried out in 2 mL Eppendorf containing 1 mL of reaction mixture in a Deep Well Maximizer (Biosh³⁵er M-BR-022UP, Taitec Japan).

A Thin Layer Chromatography (TLC) of maltooligosaccharide products was carried out on silica gel 60F²⁵⁴ plates (Merck Art 20-20 cm) and eluent using a solvent mixture of n-butanol:acid:water (12:6:6, v/v/v). Spots formed were visualized by spraying the sugar colours (0.5 g α-diphenylamine, 25 mL acetone, 2.5 mL phosphate acid, 0.5 mL aniline). All samples were applied ⁸ equal quantities (4 µL). Glucose (Sigma-Aldrich, U.S.A), maltose (M2), maltotriose (M3), maltotetraose (M4), maltopentaose (M5), maltohexaose (M6), and maltoheptaose (M7) (Megazyme) were used as standards.

Crude Enzyme Production and Amylase Activity at Different Fermentation Time

An amount of 2 mL of the 24 h old bacterial cultures (isolate T10) was inoculated into a 200 mL fla²⁷ containing 20 mL of NB medium plus 1% (w/v) starch solution, pH 6 (50 mM acetate buffer), and incubated at 30 °C for five days. The culture was sampled every 24 hours and then centrifuged and the supernatant obtained was tested for its amylase activity.

The enzyme reaction was conducted as above²⁹ then measuring amylase activity using a DNS method (Miller 1959). The absorbance⁵³ the solution was measured using a spectrophotometer at a wavelength of 540 nm. The enzyme activity (U/mL) was calculated based on the equation:

$$\text{enzyme activity} = \frac{c \times d \times 1000}{t \times mw} \text{ U/mL}$$

Where c: amylase concentration; d: dilution; t: incubation time; mw: molecular weight.

A standard curve used D-Glucose at various concentrations. One unit of amylase activity is defined as the amount of enzyme that liberates one µmol of D-Glucose per minute under the experimental condition given.

Effect of pH and Temperature on Enzyme Activity of the Selected Isolate

The optimal pH of the enzyme activity was done at pH ranges of 3.0-10.0 under standard assay conditions. Various buffers (0.05M) used were sodi-

um acetate (pH 3.0–6.0), sodium phosphate (6.0–8.0), Tris-HCL (pH 7.0–9.0), and Glycine-NaOH (pH 8.0–10.0). The enzyme reactions were incubated at 40°C for 30 min in the presence of 0.5% (w/v) starch solution (Merck). The effect of temperatures on enzyme activity was conducted at temperatures ranging from 30–90 °C in 50 mM acetate buffer at optimum pH for 30 min. Amylase activity was assayed by DNS method (Miller 1959).

RESULTS AND DISCUSSION

Isolation and An Amylolytic Assay of Bacteria Isolated from The Rhizosphere and Plant Tissue of Canna, and Litter

The results of the study found 32 bacterial isolates growing on NA medium supplemented with 1% soluble starch, with details: 11 isolates were from the rhizosphere of the Canna growing in the forest, 12 isolates from the rhizosphere of the Canna growing in the people's gardens around the forest, four isolates from the leaf tissue of the Canna growing in the people's gardens around the forest, three isolates from the roots of the Canna growing in the people's gardens around the forest, and two isolates were from the litters of the Canna in the gardens of the residents around the forest (Table 1).

The ability of the bacteria to grow and produce clear zones in the medium indicates that those bacteria were capable of producing amylase. The more amylase is released, the wider clear zones are produced due to the degradation of amylum in the medium, resulting in enhancing the amylolytic index (Ginting et al. 2021). The research results showed that eight isolates of TH6, TH7, T5, T10, D2, D3, A3, and S1 showed high total clear zone diameters (mm) of 18, 18, 18, 20, 18, 30, 18, and 18, respectively (Table 1). The consistency of the bacterial isolates, resulting in the total clear zone diameter, was shown by the isolates TH6, T10, D3, A3, and S1, while the other isolates tended to reduce or lose their amylolytic activity (Figure 2). Based on the ability of isolates to produce a clear zone diameter ≥ 18 mm and consideration of source representatives, five isolates (TH6, T10, D3, A3, and S1) were selected for further testing, namely their ability to hydrolyze starch. Hasanah et al. (2020) reported that bacterial isolates having an amylolytic index of more than 9 mm were potentials to produce amylase. According to Ochoa-Solano & Olmos-Soto (2006), bacterial isolates produce clear zones two or three times the diameter of the colony are potential enzyme producers.




Figure 2. The amylolytic zones produced by bacterial isolates of: (1) T10, (2) D3, (3) A3, (4) TH6, and (5) S1 on a NA medium + 1% soluble starch.

Table 1. Sources, number, and total clear zone diameter of amylolytic bacteria isolated from the rhizosphere, plant tissues of Canna, and litter.

Source of bacterial isolates	Isolate code	Total clear zone diameter (mm)
Rhizospheres of the Canna plants growing in the forest	TH1	16
	TH2	15
	TH3	17
	TH4	16
	TH5	16
	TH6	18
	TH7	18
	TH8	17
	TH9	17
	TH10	16
Rhizospheres of the Canna plants growing in the people's gardens around the forest	T1	15
	T2	16
	T3	16
	T4	16
	T5	18
	T6	17
	T7	15
	T8	15
	T9	15
	T10	20
	T11	17
	T12	16
Leaves of the Canna plant tissue growing in the people's gardens around the forest	D1	16
	D2	18
	D3	30
	D4	15
Roots of the Canna plant tissue growing in the people's gardens around the forest	A1	17
	A2	16
	A3	18
Litters of the Canna plant from the gardens of the residents around the forest	S1	18
	S2	16

A high number of amylolytic bacteria isolated from the rhizosphere of Canna was by Vaseekaran et al. (2010), who stated that bacteria isolated from starch-rich materials have better potential to produce amylase. Vijayalakshmi et al. (2012) found *Bacillus subtilis* KC3 isolated from the rhizosphere of *Euphorbia hirta* produced a maximum halo zone of 23 mm on a Starch Agar medium. Gebreyohannes (2015) reported that 16 bacterial isolates from soils could produce clear zones of 3–22 mm on starch agar plates. Ginting et al. (2021) found thermophilic bacteria of *Bacillus* sp. L3 and *B. caldotenax* L9 from a marine hydrothermal produces high amylolytic indexes of 3.04 and 3.52, respectively. The clear zone formed results from breaking starch compounds into simple compounds; the wider the clear zone formed, the higher the amylolytic activity (Zubaidah et al. 2019).

The characteristics of colonial morphology of the 34 isolates were rough, dry, bright, and pink; cells were Gram-positive, rod shape, motile, and had endospores. The endospore position of isolates D1–D4 and S1–S2 was in terminal, while isolates of TH1–TH11, T1–T12, and D1–D4 had endospores in the centre. All isolates were able to hydrolyse starch and produce lecithinase (Table 2). Those characteristics indicated that the bacteria were members of the genus *Bacillus*. According to Logan & De

Table 2. Morphological and physiological properties of the bacteria isolated from Canna plants and their surrounding.

Characteristics	Isolate code				
	TH1-TH11	T1-T12	D1-D4	A1-A3	S1-S2
Colonial morphology on NA Agar	Rough, dry, bright, and Pink	Rough, dry, bright and pink	Rough, dry, bright and pink	Rough, dry, bright and pink	Rough, dry, bright, and pink
Gram reaction	+	+	+	+	+
Cell shape	Rod	Rod	Rod	Rod	Rod
Motility	+	+	+	+	+
Presence of spore	+	+	+	+	+
Position of spore	Centre	Centre	Terminal	Centre	Terminal
Starch hydrolysis	+	+	+	+	+
Lecithinase production	+	+	+	+	+

Vos (2009), the main characteristics of the genus *Bacillus* are cells rod-shaped, straight or slightly curved, occurring singly and in pairs, some in chains, form endospores, Gram-positive or Gram-negative, motile, aerobes or facultative anaerobes, and mostly isolated from soil.

Identification of The Selected Isolate of T10 Based on Phenetic and Phylogenetic Characteristics

Based on the ability of the selected isolate to produce malto-oligosachharides of maltotriose, maltotetraose, and maltopentaose (Method 3.3); further characterization of the isolate T10 was conducted. The isolate had colonial morphology of irregular with undulate edges, opaque, cream-coloured, and had a granular texture. The cells formed endospores, facultatively anaerobic, Gram-positive, rod-shaped, motile, and occurring singly or in chains (Table 2, Table 3, Figure 3). These characteristics include biochemistry, physiology, and nutrition, indicating that isolate T10 was similar to those typical of the species *Bacillus cereus*. This species is a species complex within the genus *Bacillus*, with members including *B. anthracis*, *B. thuringiensis*, *B. mycoides*, and *B. toyonensis* (Luo et al. 2021).




Figure 3. The appearance of bacterial cells isolate T10 under a microscope with a magnification of 1000x. The cells appear single or in chains.

Table 3. Phenotypic characterization of the isolate T10.

Characteristics	Isolate T10
Cell length (µm)	3.00 – 4.00
Egg-yolk lecithinase	+
Anaerobic growth	+
Rhizoid colony	-
Parasporal crystal	-
Growth temperature range (°C)	10 – 45
Optimal growth temperature (°C)	35
Salinity tolerance range (%NaCl)	≤ 4
API 50CHB	
Glycerol	-
D-Ribose	+
D-Mannose	+
Methyl-αD-glucopyranoside	+
Amygdalin	+w
Arbutin	+
Salicin	+
Cellobiose	-
D-saccharose	+
D-trehalose	+
Starch	+
Glycogen	+
D-turanose	+

The electrophoresis visualization of the PCR product showed that the DNA of T10 produced a single band with a size of 1500 kb (Figure 4). The results of comparing the 16S rRNA gene sequence of isolate T10 and nucleotide sequences in the GenBank (<http://blast.ncbi.nlm.nih.gov/>) showed that the bacterium is closely related to species members of the genus *Bacillus*. The BLAST analysis showed that isolate T10 had a similarity of 99.3% with either *Bacillus toyonensis* SPa09NA, *B. toyonensis* PZ-48, or *B. toyonensis* SMP1. The phylogenetic tree was constructed using Neighbor-Joining, Model Maximum Composite Likelihood, and 1000x Bootstrapping. A dendrogram resulted from MEGA10 program showed that isolate T10 joined *B. toyonensis* SX04NA, *B. toyonensis* SPa09NA, *B. toyonensis* SMP1, *B. toyonensis* PZ-48, *B. toyonensis* BCT-7112, and *B. toyonensis* l3aM to form a separate cluster (Figure 5). Hence, isolate T10 was identified as the species member of *B. toyonensis* based on the phenetic and phylogenetic characteristics.

B. toyonensis strain BCT-7112^T was first isolated in 1966 in Japan from a survey designed to obtain naturally occurring microorganisms as probiotic cultures in the laboratory for use as probiotics in animal nutrition (Jiménez et al. 2013). The strain was first identified as *B. cereus* var. *toyoi*, and it has been used as the active ingredient of the preparation TOYO-CERIN, an additive for animal nutrition (e.g. swine, poultry, cattle, rabbits and aquaculture). Agamennone et al. (2019) isolated *B. toyonensis* strain VU-DES13 from the gut of the soil-dwelling springtail *Folsomia candida*, which was highly resistant to penicillin and inhibited the growth of a variety of pathogenic microorganisms. Its secondary metabolite clusters produce siderophores, bacteriocins, and nonribosomal peptide synthetases. Wang et al. (2021) reported that *Bacillus toyonensis* XIN-YC13 produce a novel antibiotic, toyoncin, with antimicrobial activity against *B. cereus* and *Listeria monocytogenes*. This antibiotic exerts bactericidal activity and induces cell membrane damage.




Figure 4. An electropherogram of the amplified 16S rRNA gene of isolate T10 with a size of 1500 bp. Marker (M): 1 kb DNA ladder.

Analysis of Hydrolysed Products by The Selected Bacterial Amylases Using a TLC Method

Starch hydrolysis products were assayed by oligosaccharide profile analysis on the amylase-hydrolyzed samples from the fourth isolates (T10, D3, A3, and S1) qualitatively. The results of TLC analysis showed that isolate T10 produced three bands, namely maltotriose (M3), maltotetraose (M4), and maltopentaose (M5), isolate S1 produced two bands, namely maltotriose (M3) and maltotetraose (M4), while two amylases of the isolates D3 and A3 were unable to hydrolyze starch (Figure 6).

Based on the TLC chromatogram, the starch degraded by T10 amylase resulted in malto-oligosaccharides of maltotriose, maltotetraose, and maltopentaose. Amylases can break down starch polymer bonds into shorter oligosaccharides or simple sugar molecules (Putri et al. 2012). The results showed that amylolytic bacteria with high amylolytic indexes (AI) did not correlate with their ability to degrade amyllum. The isolate T10, with its total diameter lower than isolate D3, showed a higher ability to break down starch polymer bonds into shorter or oligosaccharides. The results of this study proved that a high AI value is only sometimes accompanied by the high ability of the amylase to break down starch polymer bonds. The ability of the T10 amylase to produce the maltooligosaccharides was similar to the amylase of *Bacillus circulans* GRS 313 isolated from soil that also produced maltotriose, maltotetraose, and maltopentaose (Dey et al. 2002). On the contrary, Rahmani et al. (2013) found maltose and maltotriose produced by amylase of *Bacterium* sp. using black potato starch as substrate, while amylase of *Bacillus subtilis* strain SDP1 isolated from rhizosphere of *Acacia* produces maltotriose and maltotetraose (Ozturk et al. 2014). Furthermore, Abdul-Manas et al. (2014) reported that amylase of an alkaliphilic *Bacillus lehensis* G66 could degrade oligosaccharides by producing maltooligosaccharides with a higher degree of polymerization than maltoheptaose observed on thin-layer chromatography and high-performance liquid chromatography analyses.

Crude Enzyme Production of a Selected Isolate and Measurement of Amylase Activity at Different Culture Incubations

Based on the ability of the fourth selected amylolytic bacteria to produce different types of hydrolysed product, isolate T10 was further assayed for




Figure 5. A phylogenetic tree showing the relationship between strain T10 isolated from rhizospheres of Canna (*C. edulis*) and several species members of the genus *Bacillus* on the basis of 16S rRNA gene sequence reconstructed based on Neighbor-Joining, Model Maximum Composite Likelihood, and 1000x Bootstrapping. The analysis used a MEGA10 program and *Staphylococcus aureus* ATCC 12600 as an outgroup.




Figure 6. The product profile of starch hydrolyzed by amylase of the amylolytic bacteria (T10, D3, A3, and S1) using a TLC method with reaction times (hours): of 0, 1, 2, 3, 4, and 24 at 30°C. The Standards (STD): monosaccharide (M1), maltose (M2), maltotriose (M3), maltotetraose (M4), maltopentaose (M5), maltohexaose (M6) and maltohepta (M7).

its optimal amylase activity at different incubation times. The results showed that incubation times affected the amylase activity of isolate T10 carried out in a 0.5% starch solution at 30°C in 50 mM acetate buffer of pH 6. The amylase activity of T10 was optimal during incubation 1-3 days ranging from 0.546-0.717 U/mL and the highest amylase activity was found at 24 hr incubation of 0.717 U/mL (Figure 7). The results also showed that amylase activity decreased after 72 h of incubation. The amylase activity value at day 0 is quite high, this might be due to the measurement of the enzyme activity using the DNS method, in which reduc-

ing sugar formed from a carbon source (starch) is used by bacteria for the initial stages of growth; then, the bacteria will use the carbon source for the production of enzymes.




Figure 7. Amylase activity of T10 at different incubation times.

The amylase activity of *Bacillus* *sp.* KN isolated from Ranu Ngebel and incubated for three days was 0.016 U/mL, while strain G20 isolated from Ranu Grati was lower at about 0.0001 U/mL (Nisa et al. 2021). Luang et al. (2019) found *Bacillus* sp. 3.5AL2 isolated from soils of the unexplored Nasinuan Forest, Thailand and incubated for three days exhibiting amylase activity of 1.97 U/mg protein at the optimal conditions of 60°C and pH 7.0 after 30 min incubation with 1% starch in 0.05 M phosphate buffer. Gebreyohan (2015) reported that the amylase activity of *Bacillus* spp. decreased after 48 h incubations due to the suppression and accumulation of other byproducts in the fermentation medium and also depletion of nutrients.

Enzyme Characterization: The Effect of pH and Temperature against Enzyme Activity of The Selected Isolate

The effects of pH's on the amylase activity of isolate T10 showed that optimum conditions were in sodium acetate buffer pH 6 with an amylase activity of 0.262 U/mL and in sodium phosphate buffer pH 7 with an amylase activity of 0.341 U/mL (Figure 8). The optimal pH of isolate T10 was by Naidu et al. (2019) for *Paenibacillus* sp. D9 that its optimal pH for amylase activity is in the neutral range (pH 6-8). The increase in pH beyond these values resulted in a decline in enzyme activity. Any change in pH causes a change in the enzyme's active site (Lim & Oslan 2021). Bajpai et al. (2015) reported that the optimal pH for amylase activity of *Haloferax* sp. HA10 was at pH 7.0. According to Asgher et al. (2007), each enzyme has an optimal pH to work most actively, and the optimal pH of amylase is varied from pH 3.8 to 9.5 depending on the type of enzyme and the source. Behal et al. (2016) reported an amylase produced by *Bacillus* sp. AB04 had optimal activity at pH 8. Moreover, the enzyme is stable in neutral to alkaline (pH 7-10).

The amylase activity of isolate T10 was observed at temperatures ranging from 30-90°C pH 7.0. Amylase activity of the T10 isolate tended to optimum at 40°C with an activity value of 0.801 U/mL (Figure 9). A similar finding was also reported by Sivaramakrishnan et al. (2006) for several species of *Bacillus* sp., *B. subtilis*, *B. stearothermophilus*, *B.*




Figure 8. Amylase activity of T10 at different pH and buffers.

licheniformis, and *B. amiloliquefaciens* have optimum temperatures of 37-60°C. Gebreyohannes (2015) found that the maximum amylase activity of *Bacillus* spp. was 40°C and *Streptomyces* spp. at 37°C, used 4% starch concentration at a neutral pH and an incubated for 48 h. The crude enzyme of *Bacillus* sp. AB04 showed maximum activity at pH 8 with an optimum temperature of 40°C with more than 75% activity in range of 50 - 80°C (Behal et al. 2016). The results showed that either pH or temperature significantly affected the enzyme activity of the T10 amylase which was optimum at pH 7.0 and a temperature of 40°C.




Figure 9. Amylase activity of isolate T10 at different temperatures.

The differences in the pH and temperature characteristics of enzyme activity indicated that enzymes are specific, depending on the species that produces them. A decrease or increase in temperature can affect the secretion of extracellular enzymes by changing the physiology of the cell membrane (Rahmani et al. 2018). The optimum temperature is the temperature that causes chemical reactions at the most incredible speed (Subagiyo et al. 2017). The results showed that after reaching the optimum condition, it was seen that the activity of the T10 amylase de-

creased. High temperatures can cause enzymatic reactions to decrease because enzyme proteins undergo conformational changes so that protein molecules will experience denaturation (Yufinta et al. 2018).

The production of a specific maltooligosaccharide in high yield through the enzymatic hydrolysis of starch is of considerable commercial interest. This has been achieved on an industrial scale after discovering a suitable maltooligosaccharide-forming amylase (MFA_{ses}). Moreover, several studies have tried to improve existing methods by increasing the yields of M3 and M5. These studies have included efforts to find new wild-type strains producing MFA_{ses}, construct novel systems to achieve large-scale MFA_{ses} expression, and immobilize MFA_{ses} for stability and productivity (Ben-Ali et al. 2006). MFA_{ses} from *Bacillus toyonensis*, a novel M5-amylase, seems promising for the manufacture of high M5 syrups from starch and may apply to starch processing technologies due to their particular activity, unique substrate specificity, and endo-type action pattern (Pan et al. 2017).

44 CONCLUSIONS

It can be concluded that amounts of 32 amylolytic bacteria were isolated from rhizosphere and plant tissue of *Canna edulis*, as well as litter; the selected amylolytic bacterial isolate of T10 was capable of hydrolysing starch by producing maltotriose (M3), maltotetraose (M4) and maltopentaose (M5); and the identity of the selected isolate T10 belonged to a species member of *B. toyonensis* based on phenotypic and phylogenetic characterizations.

AUTHORS CONTRIBUTION

RNA designed, collected, and analysed the research data, O, NR and NE supervised all the process, and re-wrote the manuscript.

61

ACKNOWLEDGMENTS

We would like to thank the Research Center for Biotechnology (BRIN) and International Women University (IWU) for funding this research.

CONFLICT OF INTEREST

The author declares that there is no conflict of interest in this research.

REFERENCES

- Abdalla, M. et al., 2021. One-pot production of maltoheptaose (DP7) from starch by sequential addition of cyclodextrin glucotransferase and cyclomaltodextrinase. *Enzyme and Microbial Technology*, 149 (6), 109847. doi: 10.1016/j.enzmtec.2021.109847
- Abdul-Manas, N.H. et al., 2014. The characterization of an alkali-stable maltogenic amylase from *Bacillus lehensis* G1 and improved maltooligosaccharide production by hydrolysis suppression. *PLoS ONE*, 9(9). doi: 10.1371/journal.pone.0106481
- Atifah, N., Putri, D.H. & Irdawati, 2018. Isolation and identification of endophytic bacteria from the Andalas plant stem (*Morus macroura* Miq.). *Bioscience*, 2(1), pp.72-75.
- Agamennone, V. et al., 2019. Genome annotation and antimicrobial properties of *B. toyonensis* VU-DES13, isolated from the *Folsomia candida* gut. *Entomologia Experimentalis et Applicata*, 167, pp:269-285. doi: 10.1111/eea.12763.
- Asgher, M. et al., 2007. A thermostable α -amylase from moderately thermophilic *Bacillus subtilis* strain for starch processing. *J Food Eng.*, 79, pp.950-955.

- Bajpai, B., Chaudhary, M. & Saxena, J., 2015. Production and characterization of α -Amylase from an extremely halophilic archaeon, *Haloferax* sp. HA10. *Food Technol Biotechnol.*, 53(1), pp.11-17, doi: 10.17113/ftb.53.01.15.3824
- Ben-Ali, M. et al., 2006. Thermostability enhancement and change in starch hydrolysis profile of the maltohexaose-forming amylase of *Bacillus stearothermophilus* US100 strain. *Biochemical Journal*, 394(1), pp.51–56. doi: 10.1042/BJ20050726
- Behal, A. et al., 2016. Characterization of alkaline α - amylase from *Bacillus* sp. AB 04. *IJAB*, 8(1), pp.80-83.
- De-Moraes-Russo, C.A. & Selvatti, A.P., 2018. Bootstrap and rogue identification tests for phylogenetic analyses. *Molecular Biology and Evolution*, 35(9), pp.2327–2333. doi: 10.1093/molbev/msy118.
- Dey, G. et al., 2002. Purification and characterization of maltotrioglycosaccharide-forming amylase from *Bacillus circulans* GRS 313. *Journal of Industrial Microbiology and Biotechnology*, 28(4), pp.193-200. DOI: 10.1038/sj/jim/7000220.
- Ding, N. et al., 2021. Carbohydrate-binding module and linker allow cold adaptation and salt tolerance of maltopentaose-forming amylase from marine bacterium *Saccharophagus degradans* 2-40T. *Frontiers in Microbiology*, 12(7), pp.1–14. doi: 10.3389/fmicb.2021.708480.
- Divakaran, D., Chandran, A. & Pratap-Chandran, R., 2011. Comparative study on production of α -amylase from *Bacillus licheniformis* strains. *Brazilian Journal of Microbiology*, 42(4), pp.1397–1404. doi: 10.1590/S1517-83822011000400022.
- Duan, Y. et al., 2021. Isolation, identification, and antibacterial mechanisms of *Bacillus amyloliquefaciens* QSB-6 and its effect on plant roots. *Frontiers in Microbiology*, 12, 746799. doi:10.3389/fmicb.2021.746799.
- El-Fallal, A. et al., 2012. Starch and microbial α -amylases: from concepts to biotechnological applications. In *Carbohydrates-Comprehensive Studies on Glycobiology and Glycotechnology*. Intech Open Science, pp.459-488. doi: 10.5772/51571
- Gebreyohannes, G., 2015. Isolation and optimization of amylase producing bacteria and actinomycetes from soil samples of Maraki and Tewedros campus, University of Gondar, North West Ethiopia. *African Journal of Microbiology Research*, 9(31), pp.1877-1882.
- Ginting, E.L. et al., 2021. Isolation and identification of thermophilic amyolytic bacteria from Likupang Marine Hydrothermal, North Sulawesi, Indonesia. *Biodiversitas*, 22(6), pp.3326-3332. doi: 10.13057/biodiv/d220638.
- Gupta, R. et al., 2003. Microbial α -amylases: A biotechnological perspective. *Process Biochemistry*, 38(11), pp.1599–1616. doi: 10.1016/S0032-9592(03)00053-0.
- Hasanah, U. et al., 2020. Amyolytic activity of bacterial strains isolated from sago pulp of the traditional sago industry in Palopo, South Sulawesi. *AIP Conference Proceedings* 2231, 040073 (2020), <https://doi.org/10.1063/5.0002487>.
- Hellmuth, K. & van-den Brink, J.M., 2013. Microbial production of enzymes used in food applications. In *Microbial Production of Food Ingredients, Enzymes and Nutraceuticals*. Woodhead Publishing Limited. doi: 10.1533/9780857093547.2.262.

- Jang, E.Y. et al., 2020. Amylase-producing maltooligosaccharide provides potential relief in rats with loperamide-induced constipation. *Evidence-Based Complementary and Alternative Medicine*, 2020, 5470268. doi: 10.1155/2020/5470268.
- Jiménez, G. et al., 2013. Description of *Bacillus toyonensis* sp. nov., a novel species of the *Bacillus cereus* group, and pairwise genome comparisons of the species of the group by means of ANI calculations. *Systematic and Applied Microbiology*, 36(6), pp.383-391. doi: 10.1016/j.syapm.2013.04.008.
- Lim, S.J. & Oslan, S.N., 2021. Native to designed: Microbial α -Amylases for industrial applications. *PeerJ*, 9, pp.1-30. doi: 10.7717/peerj.11315.
- Logan, N.A. & De Vos, P., 2009. Genus I. *Bacillus* Cohn 1872. In *Bergey's Manual of Systematic Bacteriology Second Edition Volume Three The Firmicutes*. Springer, Springer Dordrecht Heidelberg London New York. pp.21-127. doi: 10.1007/b92997.
- Luang-In, V. et al., 2019. Isolation and identification of amylase-producing bacteria from soil in Nasinuan community forest, Maha Sarakham, Thailand. *Biomedical & Pharmacology Journal*, 12(3), pp.1061-1068.
- Luo, J.-c. et al., 2021. Characterization of a deep sea *Bacillus toyonensis* isolate: genomic and pathogenic features. *Frontiers in Cellular and Infection Microbiology*, 11, 629116. doi: 10.3389/fcimb.2021.629116.
- Miller, G.L., 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 31(3), pp.426-428. doi: 10.1021/ac60147a030
- Moradi, M. et al., 2014. Screening and isolation of powerful amylolytic bacterial strains. *International Journal of Current Microbiology and Applied Sciences*, 3(2), pp.758-768.
- Naidu, K. et al., 2019. Purification and characterization of α -amylase from *Paenibacillus* sp. D9 and *Escherichia coli* recombinants. *Biocatalysis and Biotransformation*, 38(1), pp.24-34. doi: 10.1080/10242422.2019.1628738.
- Nisa, I.K. et al., 2021. The potential of amylase enzyme activity against bacteria isolated from several lakes in East Java, Indonesia. *Biodiversitas*, 22(1), pp.42-49.
- Ochoa-Solano, J.L. & Olmos-Soto, J., 2006. The functional property of *Bacillus* for Shrimp feeds. *Food Microbiology*, 23(6), pp.519-525, doi: 10.1016/j.fm.2005.10.004.
- Ozturk, H.U. et al., 2014. A maltooligosaccharides producing α -amylase from *Bacillus subtilis* SDP1 isolated from rhizosphere of *Acacia cyanophylla* Lindley. *Food Biotechnology*, 28(4), pp.309-332. doi: 10.1080/08905436.2014.963600.
- Pan, S. et al., 2017. Maltooligosaccharide-forming amylase: Characteristics, preparation, and application. *Biotechnology Advances*, 35(5), pp.619-632. doi: 10.1016/j.biotechadv.2017.04.004.
- Putri, W.D.R. et al., 2012. Isolation and characterization of amylolytic lactic acid bacteria during growol fermentation, an Indonesian traditional food. *Jurnal Teknologi Pertanian*, 13(1), pp.52-60.
- Rahmani, N. et al., 2013. Production of maltooligosaccharides from black potato (*Coleus tuberosus*) starch by α -amylase from a marine bacterium (*Brevibacterium* sp.). *Microbiology Indonesia*, 7(3), pp.129-136. doi: 10.5454/mi.7.3.6.

- Rahmani, N. et al., 2018. Xylanase and feruloyl esterase from actinomycetes cultures could enhance sugarcane bagasse hydrolysis in the production of fermentable sugars. *Bioscience Biotechnology and Biochemistry*, 82(5), pp.904–915. doi: 10.1080/09168451.2018.1438169.
- Santos, F.D.S. et al., 2018. *Bacillus toyonensis* improves immune response in the mice vaccinated with recombinant antigen of bovine herpesvirus type 5. *Benef. Microbes*, 9(1), pp.133–142. doi: 10.3920/BM2017.0021.
- Sivaramakrishnan, S. et al., 2006. α -Amylase from microbial sources: An overview on recent developments. *Food Technol. Biotechnol.*, 44(2), pp.173–184.
- Smibert, R.M. & Krieg, N.R., 1981. General Characterization. In *Manual Methods for General Bacteriology*. American Society for Microbiology, Washington.
- Subagiyo, Djarod, M.S.R. & Setyati, W.A., 2017. Potensi ekosistem mangrove sebagai sumber bakteri untuk produksi protease, amilase, dan selulase. *Jurnal Kelautan Tropis*, 20(2), pp.106–111.
- Vaseekaran, S., Balakumar, S. & Arasaratnam, V., 2010. Isolation and identification of a bacterial strain producing thermostable α - amylase. *Tropical Agricultural Research*, 22(1), pp.1–11. doi: 10.4038/tar.v22i1.2603.
- Vijayalakshmi et al., 2012. Isolation and characterization of *Bacillus subtilis* KC3 for amylolytic activity. *International Journal of Bioscience, Biochemistry and Bioinformatics*, 2(5), pp.336–341. doi: 10.7763/IJBBB.2012.V2.128.
- Wang, J. et al., 2021. Toyocin, a novel leaderless bacteriocin that is produced by *Bacillus toyonensis* XIN-YC13 and specifically targets *B. cereus* and *Listeria monocytogenes*. *Applied and Environmental Microbiology*, 87(12), e00185–21. doi: 10.1128/AEM.00185–21.
- Yopi et al., 2017. Isolation and characterization of mannanase, xylanase, and cellulase from marine bacteria *Bacillus* sp. *Biofarmasi Journal of Natural Product Biochemistry*, 15(1), pp.15–20. doi: 10.13057/biofar/f150103.
- Yufinta, C.P., Julyantoro, P.G.S. & Pratiwi, M.A., 2018. Pengaruh penambahan *Bacillus* sp. terhadap kelulushidupan pasca larva udang Vannamei (*Litopenaeus vannamei*) yang terinfeksi vibriosis. *Current Trends in Aquatic Science*, 1(1), pp.89–95.
- Zhao, C. et al., 2017. Functional properties, structural studies and chemo-enzymatic synthesis of oligosaccharides. *Trends in Food Science and Technology*, 66, pp.135–145. doi: 10.1016/j.tifs.2017.06.008.
- Zhuang, Y., Zhou, X. & Wang, S., 2012. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Plant Systematics and Evolution*, 298(7), pp.3389–3402. doi: 10.2503/ijshs.58.977.
- Zubaidah, A. et al., 2019. Screening bakteri selulolitik dan amilolitik pada rumen sapi sebagai kandidat probiotik pada budidaya ikan secara *in vitro*. *Jurnal Riset Akuakultur*, 14(4), pp.261–271.

Isolation and Characterization of Rhizospheric Bacteria Associated with Canna Plant for Production of Maltooligosaccharide Amylase

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

www.frontiersin.org

Internet Source

1%

2

Paola P. Pereira, Gonzalo A. Torres Tejerizo, Marilina Fernandez, Anicet R. Blanch, Paola S. Gonzalez, Elizabeth Agostini. "Polyphasic characterization and identification of the bioremediation agent *Bacillus* sp. SFC 500-1E", Genomics, 2020

Publication

1%

3

onlinelibrary.wiley.com

Internet Source

1%

4

Kendal, D.. "Plant traits link people's plant preferences to the composition of their gardens", Landscape and Urban Planning, 20120330

Publication

1%

5

www.ncbi.nlm.nih.gov

Internet Source

1%

6	link.springer.com Internet Source	<1 %
7	pu-toyama.repo.nii.ac.jp Internet Source	<1 %
8	Jiang, Ru-wei, Xiao-guang Du, Xuan Zhang, Xin Wang, Ding-yu Hu, Tao Meng, Yue-lei Chen, Mei-yu Geng, and Jing-kang Shen. "Synthesis and bioassay of β -(1,4)-D-mannans as potential agents against Alzheimer's disease", <i>Acta Pharmacologica Sinica</i> , 2013. Publication	<1 %
9	scirp.org Internet Source	<1 %
10	www.j3.jstage.jst.go.jp Internet Source	<1 %
11	Gebreyohannes Gebreselema. "Isolation and optimization of amylase producing bacteria and actinomycetes from soil samples of Maraki and Tewedros campus, University of Gondar, North West Ethiopia", <i>African Journal of Microbiology Research</i> , 2015 Publication	<1 %
12	www.microbiologyresearch.org Internet Source	<1 %
13	ijlpr.com Internet Source	<1 %

14	Thierry K. S. Janssens, Tjalf E. de Boer, Valeria Agamennone, Niels Zaagman, Nico M. van Straalen, Dick Roelofs. " Draft Genome Sequence of VU-DES13, Isolated from (Collembola: Entomobryidae) ", Genome Announcements, 2017 Publication	<1 %
15	journal.unnes.ac.id Internet Source	<1 %
16	www.scielo.br Internet Source	<1 %
17	www.thaiscience.info Internet Source	<1 %
18	biodiversitas.mipa.uns.ac.id Internet Source	<1 %
19	eurosurveillance.org Internet Source	<1 %
20	www.academicjournals.org Internet Source	<1 %
21	www.tkvjerze.com Internet Source	<1 %
22	www.tnsroindia.org.in Internet Source	<1 %
23	F.D.S. Santos, M.R.A. Ferreira, L.R. Maubrigades, V.S. Gonçalves et al. " BCT -	<1 %

7112 transient supplementation improves vaccine efficacy in ewes vaccinated against epsilon toxin ", Journal of Applied Microbiology, 2020

Publication

24

J. C. Oliveira, J. F. Sales, A. Rubio-Neto, C. F. Silva, M. A. Soares, F. G. Silva. "Biological control in the germination of seeds from two species native of the Cerrado region", Brazilian Journal of Biology, 2021

Publication

<1 %

25

Nurachman. "Identification a Novel Raw-Starch-Degrading- α -Amylase from a Tropical Marine Bacterium", American Journal of Biochemistry and Biotechnology, 2010

Publication

<1 %

26

www.sciencegate.app

Internet Source

<1 %

27

M. Fern andez-Gonz lez, J.F.  Beda, T.G. Vasudevan, R.R. Cordero Otero, A.I. Briones. "Evaluation of polygalacturonase activity in *Saccharomyces cerevisiae* wine strains", FEMS Microbiology Letters, 2004

Publication

<1 %

28

Mutia Elida, Agustina Agustina, Ermiati Ermiati, Susi Desminarti. "Isolate Characterization and Amylolytic Properties of Lactic Acid Bacteria from Traditional

<1 %

Fermented Dadih", IOP Conference Series: Earth and Environmental Science, 2022

Publication

29

Sidra Dr, Muhammad Muneeb Zaman,
Zunaira Farooq, Amina Hafeez et al.
"Supplementation of PUFA extracted from
microalgae for the development of chicken
patties", PeerJ, 2023

Publication

<1 %

30

dokumen.pub

Internet Source

<1 %

31

www.science.gov

Internet Source

<1 %

32

www.webnetsystems.com

Internet Source

<1 %

33

"Bacilli in Agrobiotechnology", Springer
Science and Business Media LLC, 2022

Publication

<1 %

34

Chandran Masi, Abel Tebiso, Selva Kumar K V.
"Isolation and characterization of potential
multiple extracellular enzyme-producing
bacteria from waste dumping area in Addis
Ababa", Heliyon, 2023

Publication

<1 %

35

Dubey, K.K.. "Production of demethylated
colchicine through microbial transformation

<1 %

and scale-up process development", Process Biochemistry, 200803

Publication

36

core.ac.uk

Internet Source

<1 %

37

eclipse.nichd.nih.gov

Internet Source

<1 %

38

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

39

Changqing Zhao, Xingxiu Zhao, Jing Zhang, Wei Zou, Yi Zhang, Li Li, Jun Liu. "Screening of Bacillus Strains from Sun Vinegar for Efficient Production of Flavonoid and Phenol", Indian Journal of Microbiology, 2016

Publication

<1 %

40

Mohammed Abdalla, Bo Jiang, Hinawi A.M. Hassanin, Luhua Zheng, Jingjing Chen. "One-pot production of maltoheptaose (DP7) from starch by sequential addition of cyclodextrin glucotransferase and cyclomaltodextrinase", Enzyme and Microbial Technology, 2021

Publication

<1 %

41

Vinod Kumar, Manisha Nanda, Ajay Singh. "Effect of bacterial amylase pretreatment on bioethanol production from starch-based solid waste (SBSW)", Energy Sources, Part A:

<1 %

Recovery, Utilization, and Environmental Effects, 2016

Publication

42	aem.asm.org Internet Source	<1 %
43	docplayer.net Internet Source	<1 %
44	ejournal.undip.ac.id Internet Source	<1 %
45	etheses.saurashtrauniversity.edu Internet Source	<1 %
46	refubium.fu-berlin.de Internet Source	<1 %
47	research.vu.nl Internet Source	<1 %
48	www.nature.com Internet Source	<1 %
49	www.tandfonline.com Internet Source	<1 %
50	"Bioprospecting of Microorganism - Based Industrial Molecules", Wiley, 2021 Publication	<1 %
51	Dipesh Kumar Verma, Gunjan Vasudeva, Chandni Sidhu, Anil K. Pinnaka, Senthil E. Prasad, Krishan Gopal Thakur. "Biochemical	<1 %

and Taxonomic Characterization of Novel Haloarchaeal Strains and Purification of the Recombinant Halotolerant α -Amylase Discovered in the Isolate", *Frontiers in Microbiology*, 2020

Publication

52

Fatma Karray, Manel Ben Abdallah, Najwa Kallel, Manel Hamza, Manel Fakhfakh, Sami Sayadi. "Extracellular hydrolytic enzymes produced by halophilic bacteria and archaea isolated from hypersaline lake", *Molecular Biology Reports*, 2018

Publication

53

I Dewiyanti, D Darmawi, Z A Muchlisin, T Z Helmi, I I Arisa, R Rahmiati, E Destri. "Cellulase enzyme activity of the bacteria isolated from mangrove ecosystem in Aceh Besar and Banda Aceh", *IOP Conference Series: Earth and Environmental Science*, 2022

Publication

54

Pengyu Luan, Yanjie Yi, Yifan Huang, Liuqing Cui, Zhipeng Hou, Lijuan Zhu, Xiujuan Ren, Shao Jia, Yang Liu. "Biocontrol potential and action mechanism of *Bacillus amyloliquefaciens* DB2 on *Bipolaris sorokiniana*", *Frontiers in Microbiology*, 2023

Publication

<1 %

<1 %

<1 %

55	Yawei Wang, Jing Wang, Zhongqiang Zhang, Jiangke Yang, Ossi Turunen, Hairong Xiong. "High-temperature behavior of hyperthermostable Thermotoga maritima xylanase XYN10B after designed and evolved mutations", Applied Microbiology and Biotechnology, 2022 Publication	<1 %
----	--	------

56	archimer.ifremer.fr Internet Source	<1 %
----	---	------

57	bnrc.springeropen.com Internet Source	<1 %
----	---	------

58	jurnal.ugm.ac.id Internet Source	<1 %
----	---	------

59	pakjas.com.pk Internet Source	<1 %
----	---	------

60	repositorio.ipicyt.edu.mx Internet Source	<1 %
----	---	------

61	static.frontiersin.org Internet Source	<1 %
----	---	------

62	watermark.silverchair.com Internet Source	<1 %
----	---	------

63	www.freepatentsonline.com Internet Source	<1 %
----	---	------

64	www.lmaleidykla.lt	
----	--	--

<1 %

65

"Microorganisms in Saline Environments: Strategies and Functions", Springer Science and Business Media LLC, 2019

Publication

<1 %

66

Abdul Manas, Nor Hasmaliana, Samson Pachelles, Nor Muhammad Mahadi, and Rosli Md. Illias. "The Characterisation of an Alkali-Stable Maltogenic Amylase from *Bacillus lehensis* G1 and Improved Maltoligosaccharide Production by Hydrolysis Suppression", PLoS ONE, 2014.

Publication

<1 %

67

Ricardo R. Moraes, Aline M. Pascoal, Samantha S. Caramori, Flavio M. Lopes, Kátia F. Fernandes. " Immobilization of α -Amylase onto Fibers ", Enzyme Research, 2013

Publication

<1 %

68

Sri Wahyuni, Sarinah, Wa Ode Gustiani Purnamasari, Usman Pato, Prima Endang Susilowati, Asnani, Andi Khaeruni. "Identification and Genetic Diversity of Amylase Producing Lactic Acid Bacteria from Brown Rice (*Oryza nivara*) Wakawondu Cultivar Based on 16S rRNA Gene", Fermentation, 2022

Publication

<1 %

69

Ticiane Carvalho Farias, Haroldo Yukio Kawaguti, Maria Gabriela Bello Koblitz. "Microbial amylolytic enzymes in foods: Technological importance of the *Bacillus* genus", Biocatalysis and Agricultural Biotechnology, 2021

Publication

<1 %

70

M. Y. Jung. "Bacillus acidiproducens sp. nov., vineyard soil isolates that produce lactic acid", INTERNATIONAL JOURNAL OF SYSTEMATIC AND EVOLUTIONARY MICROBIOLOGY, 09/01/2009

Publication

<1 %

71

Panpan Wang, Tao Wang, Mohamedelfateih Ismael, Xin Wang, Yanglei Yi, Xin Lü. "Development of an electroporation method and expression patterns of bacteriocin-encoding genes in *Companilactobacillus crustorum* MN047", Food Bioscience, 2021

Publication

<1 %

72

Rugaiyah A. Arfah. "Production Optimization and Characterization of Amylase Enzyme Isolated from Termofil Bacteria *Bacillus* sp RSAll-1b from Lejja Hot Spring South Sulawesi", American Journal of Biomedical and Life Sciences, 2015

Publication

<1 %

73

Valeria Agamennone, Joeri Straalen, Abraham Brouwer, Tjalf E. Boer et al. " Genome annotation and antimicrobial properties of - 13, isolated from the gut ", Entomologia Experimentalis et Applicata, 2019

Publication

<1 %

74

Rosyid Ridlo Al Hakim, Erie Kolya Nasution, Rizaldi, Siti Rukayah, Sri Riani. "Daily behavior of alpha-male compared with subordinate male in long-tailed macaque", AIP Publishing, 2023

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On