

Proteolytic and Clot Lysis Activity Screening of Crude Proteases Extracted from Tissues and Bacterial Isolates of Holothuria Scabra

by Oedjijono Oedjijono

Submission date: 14-Feb-2022 01:51PM (UTC+0700)

Submission ID: 1761963530

File name: 2021_IOP_Conf._Ser._Earth_Environ._Sci._755_012016.pdf (912.19K)

Word count: 4564

Character count: 24839

6

Proteolytic and Clot Lysis Activity Screening of Crude Proteases Extracted from Tissues and Bacterial Isolates of *Holothuria Scabra*

Nur Hidayati¹, Hayatun Fuad¹, Hendra Munandar², Dewi Seswita Zilda³, Nurrahman Nurrahman⁴, Miswar Fattah^{1,5}, Oedjijono Oedjijono⁶, Amin Samiasih⁷, Stalis Norma Ethica^{1*}

¹Magister Study Program of Medical Laboratory Science, Universitas Muhammadiyah Semarang, 50273, Indonesia

²Research and Development Division for Marine Bio Industry, Institute of Sciences Lombok, Nusa Tenggara Barat, 83352, Indonesia

³Research and Development Center for Marine and Fishery Product Processing and Biotechnology, Ministry of Maritime Affairs and Fisheries of Indonesia, Jakarta, 10260, Indonesia

⁴Food Technology Study Program, Faculty of Nursing and Health Sciences, Universitas Muhammadiyah Semarang, 50273 Indonesia

⁵Department of Molecular Biology, Prodia Widyahusada Laboratory, Jakarta, 10430, Indonesia

⁶Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, 53122, Indonesia

⁷Magister Study Program of Nursing Science, Universitas Muhammadiyah Semarang, 50273, Indonesia

*Corresponding author: norma@unimus.ac.id

Abstract. Cardiovascular diseases (CVDs) are still the leading mortality causes in the last decades. The ailments are multifactorial characterized by excessive clot (thrombus) formation in the blood vessels. Thrombus could be degraded through thrombolysis mechanism by plasmin activated by various fibrinolysis agents including, urokinase, nattokinase, or streptokinase. However, the use of these agents is restricted by relatively high cost, short half-life, allergic reaction, and bleeding effects. The search for more economical and safer thrombolytic (clot lysis) agents are essential to address the underlying problem in CVD therapy. Among Holothurians, *H. scabra* has been known to have the highest protein content making it ideal substrate for protease enzymes including fibrinolytic types with clot lysis properties. However, isolation of a proteases with antithrombotic activities either from tissue or from bacteria of *H. scabra* has not been reported. This study aimed to screen proteolytic and clot lysis activities of crude protease extracts from tissue and bacteria isolated from fermented intestine of *H. scabra*. Crude protease of tissue of *H. scabra* was extracted by cold centrifugation, which activity was measured using UV-spectrophotometer. Crude protease of proteolytic bacteria selected by clear zones on skim milk agar (SMA) medium was isolated from nutrient broth (NB). Extracts showing proteolytic activity were subjected to gravimetry-based clot lysis test. As results, crude proteases isolated directly from the *H. scabra*'s tissue showed low proteolytic activities, thus were no proceed to clot lysis activity test. Crude protease extracted bacteria could show both proteolytic and clot lysis activities. In conclusion, based on this screening study, intestine of *H. scabra* is a rich



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

source of proteolytic bacteria, some of them could produce crude enzymes showing competitive thrombolysis activities with Nattokinase. Thus, they appeared to have more potentials to be developed as thrombolysis agent than those directly extracted from the organism's tissue.

1. Introduction

Cardiovascular, venous and arterial thrombosis are multifactorial diseases characterized by unrestrained clot formation in the blood vessels [1]. The occurrence of thrombosis is due to failure of homeostasis. This condition could lead to blockage of the blood vessels and during revery, which could result in blockage of the heart or brain muscle leading to death [2-3]. The world's burden of CVD falls, mainly, on the low and center pay (LMI) nations, representing over 80% of CVD mortality. WHO data says that due to cardiovascular disease, especially stroke and heart disease, about 17 million people are dying each year and predicts that this number will increase to 23.3 million by 2030 [4].

Blood clots have a major protein component, namely fibrin. Blood clots that block blood vessels can be destroyed by fibrinolysis by plasmin. The main process of fibrinolysis is by activating plasminogen into the proteolytic enzyme plasmin. This enzyme changes the shape of the thrombus by destroying the fibrin in the blood clot so that the progression of thrombosis is inhibited [5-6]. Commonly used drugs such as urokinase, tissue-type plasminogen activator (t-PA) and streptokinase, activate plasmin and convert plasminogen to plasmin which degrades fibrin. However, these agents have limitations such as higher cost, shorter half-life, allergic reactions and intravenous administration of these agents are also known to cause bleeding. This has prompted investigators to continue to look for new, safer and less expensive sources of fibrinolytic enzymes for thrombolysis [7-10].

Sea cucumbers are marine invertebrates gaining popularity among researchers in recent decades, not only for their nutritional value, but also for their potential health benefits and therapeutic applications [11]. Sea cucumbers are known organisms producing fibrinolytic proteases with antithrombotic including thrombolysis activity. This is because it is rich in protein, the substrate for the protease enzyme in these organisms, with a content in average between 43.24 - 48.27 % w/w [13]. Of the various species of sea cucumbers found in Indonesia, the sand or white sea cucumbers (*Holothuria scabra*) had been reported to have the highest protein content (79.6% w/w) [14]. However, the isolation of the protease enzyme with fibrinolytic ability from *H. scabra* from both tissue and bacteria of its digestion system has not been published. In this study, blood clot lysis activities of crude protease enzymes from both tissue and bacterial isolates of *Holothuria scabra* digestive tract were evaluated.

4

2. Materials and Method

2.1. Material

Sample used in this study was a sea cucumber *H. scabra*, which was obtained from its captivity located at Lembaga Ilmu Pengetahuan Indonesia Institute for Bio Industry (BBIL LIPI), Kodek Gulf Village, Lombok, West Nusa Tenggara. The size of the 12-mo sea organism was 177 x 4 cm with a weight of 144.01 g. The main reagents used included Nutrient Agar (Oxoid Lot: 2438621), Skim Milk Agar (Oxoid Lot: 4249869), physiological NaCl, casein (Merck, cat. 750344), standard tyrosine (Merck, cat. 1.083710025), Na₂CO₃ (Merck cat. 6392.1000), Folin-reagent Ciocalteau (Merck, cat. 1.0001.0500). The equipment used were Analytical scale (Mettler Toledo), autoclave (Hirayama HVE-50), Laminair Air Flow Cabinet (Dalton), spectrophotometer (Thermo), Micro centrifuge (Bechman Counter), incubator (Mammert), micropipette (Socorex Calibra 822). Vortex mixer (Genie 2TM 12-82), calipers, micro tubes and other tools used in the laboratory according to work procedures.

2.2. Method

2.2.1 Crude Protease Extraction from Tissue of *H. scabra*

To obtain crude protease extract, 3 samples of *H. scabra* "muscle" tissue (from tentacle, ventral and posterior part) were prepared. As much as 2 g each sample was homogenously blended and then added with 2 ml of Tris HCl 0.05 M (pH = 8.22). The mixture was subjected to centrifugation at 10,000 rpm and 4 °C for 15 min, and the supernatant was set aside as crude protease extract. The extract was subjected to protease activity test.

2.2.2 Activity Test of Crude Protease Extracted from Tissue of *H. scabra*

The crude enzyme extracts both from tissues of *H. scabra* and was then determined for its proteolytic activity based on method reported by Cupp-Enyard [15]. A total of 250 µL of crude extract of the enzyme were added to 250 µL of 1% casein (1 g of casein in 100 ml of K-Phosphate buffer 0.05 M pH=7.5). The mixture was incubated for 30 min. The mixture reaction was terminated by addition of 500 µL of 0.4M TCA. A total of 25 µL of supernatant filtrate was added with 1.25 ml of 0.4 M Na₂CO₃ and 125 µL of Folin Ciocalteu reagent and incubated at 40 °C for 30 min. The absorbance of the mixture was measured using a spectrophotometer at λ of 600 nm. One unit of enzyme activity is defined as the amount of enzyme required to release 1 µmol of tyrosine on the casein substrate per minute. The linear tyrosine equation was used as a standard curve to be interpolated with the absorbance value obtained [16].

2.2.3 Proteolytic Bacterial Selection from Intestinal Organ of *H. scabra*

Preparation of bacterial culture from intestine of *H. scabra* was carried out aerobically and anaerobically following to method reported by Zhang et al. [17]. Next, in a sterile test tube, 1 g of the sample was added with 9 mL of sterile sea water, which then homogenized using a vortex. The resulted suspension was diluted and then inoculated on Skim Milk Agar (SMA) media containing casein protein with incubation at 30 °C for 24-48 h to test the ability of bacteria to hydrolyze protein [18]. The bacterial culture that grows and forms a clear zone around the colony is then separated and purified by the quadrant streak method [18-19]. Protease activity was indicated by the presence of a clear circular zone around the colony. The quotient of clear circle diameter and colony diameter is expressed as relative protease activity. This applies also for other common hydrolytic enzyme activities such as lipase, amylase and cellulase from bacteria [20-24]. Each bacterial isolate able to produce clear zones was inoculated on SMA media, one on each and incubated at 30 °C for a week. Hydrolysis Capacity Index calculation was measured on the seventh day of observation on the clear zone to determine the level of ability of each bacterial culture in hydrolyzing the substrate [24].

2.2.4 Crude Protease Extraction from Proteolytic Bacterial Isolates of *H. scabra*

For quantitative protease activity test, bacterial cultures in a liquid medium containing 20% soy milk and 5% lactose as a protein source or substrate were used as a starter in Erlenmeyer. The starter was incubated and shaken on the shaker at a speed of 100 rpm at room temperature for 24 h. A total of 1% starter was then inoculated into 20 ml of the same medium, incubated for 3 days on a shaker at 130 rpm. Next, the culture was centrifuged at 10,000 rpm 4 °C for 15 min to separate the enzyme solution from the substrate particles. The filtrate obtained was regarded as a crude extract of the enzyme [25].

2.2.5 Clot Lysis Activities of Bacterial Crude Proteases

For the gravimetric thrombolysis test, the empty micro-centrifuge tube was weighed and each weight recorded. Blood was taken from healthy volunteers as much as 500 µl, put in a micro centrifugation tube and incubated at 37 °C for 45 min. After clotting, serum was taken so as not to interfere with the blood clot. Weight of the clot was determined as follows: weight of clot = weight of tube with clot - weight of empty tube [26].

Crude extract enzymes from the tentacle, ventral and posterior muscle tissue *H. scabra* tissue were pipetted by 100 µl, each was put into a micro-centrifuge tube containing the blood clot (each was

24
7
labeled). In addition, aquadest was used as a negative control and streptokinase as a positive control. All tubes were then incubated at 37 °C for 90 min and clots lysis was observed [26-27]. Percentage of Clot lysis percentage was calculated based on weight difference of micro-centrifuge tubes obtained before and after clot lysis [26].

20

3. Results and Discussion

3.1. Sample preparation

Sample preparation in this work was intended for two purposes. First was to obtain crude enzyme extract from muscle tissue of *H. scabra* to detect its proteolytic activity. Second was to isolate proteolytic bacteria from intestine part of the sand sea cucumber, which produce protease. For both purposes, a sand sea cucumber or *Holothuria scabra* was freshly taken and used as sample as (Figure 1). To obtain crude enzyme from muscle tissues, the gut part and the muscle tissue was separated aseptically by dissecting set (Figure 2). Each of the muscle parts (tentacle, ventral and posterior) was grounded separately using sterile mortar and then subjected to centrifugation. Each supernatant was later tested for proteolytic activity using spectrophotometer based on absorbance at λ 600 nm.



Figure 1. A sand sea cucumber or *Holothuria scabra* sample freshly captured from its captivity



Figure 2. Muscle part of *H. scabra* (white color)

This work is the first reporting the cultivation of proteolytic bacteria from gut of *H. scabra*, a rich-protein, sand sea cucumber from Indonesia. To obtain culturable bacteria, the gut part was blended using sterile mortar. The anaerobically prepared sample to obtain culturable bacteria is shown in Figure 2 & 3.




Figure 3. Intestine or gut part of *H. scabra* (orange color)




Figure 4. Fermented intestine of *H. scabra* sample as source of culturable bacteria

3.2. Activity Test Results of Crude Protease Extracted from Tissue of *H. scabra*

Based on linear tyrosine equation obtained from a standard curve made following previous method at λ 600 nm [16], protease activity of crude protease could be calculated using the following equation: $Y = 0,0019 X + 0,0029$, where Y= absorbance, while X= activity (in U/mL)

The resulted data of proteolytic activity of crude enzyme extracted from tissues based on absorbance at λ 600 nm were shown in Table 1.

Table 1. Activity of crude protease extracted from *H. scabra* tissues with tyrosine as standard

No.	Treatment	Absorbance at 600 nm* (average)				Relative activity (U/mL)
		Tentacle	Ventral	Posterior	Control (tyrosine)	
1	Vortex	0.0791	0.0759	0.0850	0.0994	-1505.1170
2	No vortex	0.0754	0.0897	0.0881	0.0539	3103.3041

*Measurement was performed in 6 repetitions

In most activity test methods using spectrophotometer, homogenization of crude protease extract using vortex was common in order to homogenize samples prior to absorbance measurement. Data in **Table 1** show that activity test results of samples without vortex were positive. However, it could mean that enzyme having proteolytic activity was present in non-homogenous protease crude extract. So, it was assumed that concentration of crude protease extracted from tissue in this work was not so high. It means, proteolytic activity could not be confirmed. Based on this, the result was not proceeded to next step of clot lysis screening.

3.3 Proteolytic Bacterial Selection from Intestine of *H. scabra*

Unlike result from previous result using intestine of *Holothuria leucospilota* as sample [17], in this study, culturable bacteria could not be obtained from aerobic cultures after 24-h incubation. The gut bacteria appeared to grow well under anaerobic condition resulting 12 unique colonies, which were subsequently purified and named HSFI-1 to HSFI-12 (HSFI refers to *Holothuria scabra* Fermented Intestine). Macroscopic characteristics of HSFI-1 to -12 were shown in Table 2. All of the twelve isolates were then tested for their ability to degrade protein on SMA media. This selection step resulted in 10 proteolytic isolates (Table 3).

Table 2. Macroscopic characteristics of bacteria isolated from fermented intestine of sand sea cucumber (*H. scabra*) on NA media

No.	Isolate Code	Bacterial Colony Morphology					
		Shape	Edge	Diameter size (mm)	Colour	Elevation	Consistency
1	HSFI-1	circular	entire	1	cream	convex	smooth
2	HSFI-2	circular	undulate	8	white	umbonate	smooth
3	HSFI-3	circular	entire	3	cream	raised	smooth
4	HSFI-4	irregular	undulate	5	cream	convex	smooth
5	HSFI-5	filamentous	polyform	8	white	crater-form	rough
6	HSFI-6	irregular	undulate	5	cream	convex	smooth
7	HSFI-7	circular	entire	1	white	convex	smooth
8	HSFI-8	circular	entire	4	cream	convex	smooth
9	HSFI-9	circular	entire	8	cream	umbonate	smooth
10	HSFI-10	circular	entire	2	cream	convex	smooth
11	HSFI-11	circular	entire	4	cream	convex	smooth
12	HSFI-12	circular	entire	4	cream	convex	smooth

Note: HSFI (*Holothuria scabra* fermented intestine)

Table 3. Bacterial colony morphology, proteolytic activity and clot-lysis test result data of bacteria isolated from fermented intestine of *H. scabra*.
























Isolate code	Colony morphology	Proteolytic zone*	Clot lysis (%)
HSFI-1		N.A.	NA
HSFI-2			26.06
HSFI-3			5.24
HSFI-4			19.30
HSFI-5			62.03

Table 4. Continued ...

Sample code	Colony morphology	Proteolytic zone*	Clot lysis (%)
HSFI-6			32.94
HSFI-7		N.A	NA
HSFI-8			84.71
HSFI-9			7.78
HSFI-10			31.28
HSFI-11			46.70
HSFI-12			12.47
Aquadest	 26 N.A.	N.A.	0.29
Nattokinase	N.A.	N.A.	56.05

*After 24-h incubation

19 *Crude Protease Extraction from Proteolytic Bacterial Isolates*

In this study, crude protease extract was retrieved after centrifugation of each of bacterial inoculum in skim milk broth medium. The supernatant was considered as crude protease, thus 10 extracts from HSFI-1 to -6 and HSFI-8 to -12 could be obtained. The 10 bacterial protease extracts were subjected to clot lysis test as described in previous study [23].

3.5 Clot Lysis Activities of Bacterial Crude Proteases

Visual representation of clot lysis test result on 10 bacterial protease extracts is shown in **Figure 3**. When 100 μ l water was added to the negative control (Bl) clot, negligible clot lysis was observed. In other words, no clot lysis was observed in Bl tube. Whereas, tubes to which different crude extracts from different bacterial isolates HFSI-1 to -6 and HFSI-8 to -12 were added, various levels of clot lysis could be visually seen. Percentage of clot lysis obtained after treating clots with water, streptokinase and control group based on gravimetry method is shown in **Table 3**.




Table 5. Clot-lysis activity test results of bacterial crude protease using gravimetry method.

Tube	Isolate code	Initial clot weight	Final clot weight	Lysis percentage (%)
N				
1	HSFI-2	0.2467	0.1824	26.06
2	HSFI-3	0.2326	0.2204	5.24
3	HSFI-4	0.2388	0.1927	19.30
4	HSFI-5	0.4128	0.1567	62.03
5	HSFI-6	0.2486	0.1667	32.94
6	HSFI-8	0.2519	0.2323	84.71
7	HSFI-9	0.5776	0.0883	7.78
8	HSFI-10	0.2439	0.1676	31.28
9	HSFI-11	0.2572	0.1371	46.70
10	HSFI-12	0.2526	0.2211	12.47
11	Aqudest	0.2668	0.2650	0.29
12	Nattokinase	0.2462	0.1082	56.05

As seen on **Table 3**, various percentage of clot lysis was observed from each of crude protease extract from 10 bacterial isolates, HFSI-1 to -6 and HFSI-8 to -12. There were two extracts of crude bacterial protease that could show higher clot lysis activity than positive control, Nattokinase coming from HFSI-5 and -8. To confirm the results however, *in-vitro* validation test should be performed either on the

proteolytic and clot lysis activities of these bacterial protease extracts. Moreover, purification and characterization of bacterial proteases from isolate HFSI-5 and -8 is important to do, especially if the aim is to further develop them into prototypes of thrombolysis agent. Overall, results of this study showed that crude proteases of bacteria isolated from fermented intestine of *H. scabra* have the potential as thrombolysis agent more than those extracted directly from the sea cucumber's muscle tissue.

4. Conclusion

Based on this screening study, intestine of *H. scabra* is a rich source of proteolytic bacteria. Some of them could produce crude enzymes showing competitive thrombolysis activities with Nattokinase. Thus, they appeared to have more potentials to be developed as thrombolysis agent than those directly extracted from the organism's tissue.

4 5. Acknowledgments

This work was financially supported by Ministry of Research and Technology/ National Research and Innovation Agency of the Republic of Indonesia (Kemenristek/BRIN) through Thesis Magister Research Program (Program Penelitian Tesis Magister) 2020 [Grant number 0004/UNIMUS.L/PG/PJ/2020]. *H. scabra* samples were obtained from sea cucumber captivity of Balai Bio Industry Laut, Lembaga Ilmu Pengetahuan Indonesia (BBIL LIPI/ Indonesian Institute of Science), Kodek Gulf Village, Lombok, West Nusa Tenggara.

References

- [1] Ashorobi D, Fernandez R. Thrombosis. InStatPearls [Internet] 2019 May 6. StatPearls Publishing. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538430/>
- [2] Martin K, Ma AD, Key NS. Molecular basis of hemostatic and thrombotic diseases. InMolecular Pathology 2018 Jan 1 (pp. 277-297). Academic Press. <https://doi.org/10.1016/B978-0-12-802761-5.00015-8>
- [3] Sarwar MH, Mughal AR, Mughal S, Sarwar M. Concerns of Heart Diseases and Mediations to Encourage Healthful Actions for Their Deterrence. International Journal of Bioinformatics and Biomedical Engineering. 2015;1(2):70-6. Available from: <http://files.aiscience.org/journal/article/pdf/70070046.pdf>
- [4] McAloon CJ, Boylan LM, Hamborg T, Stallard N, Osman F, Lim PB, Hayat SA. The changing face of cardiovascular disease 2000–2012: An analysis of the world health organisation global health estimates data. International journal of cardiology. 2016 Dec 1;224:256-64. <https://doi.org/10.1016/j.ijcard.2016.09.026>
- [5] Weisel JW, Litvinov RI. Fibrin formation, structure and properties. In Fibrous proteins: structures and mechanisms 2017 (pp. 405-456). Springer, Cham. https://doi.org/10.1007/978-3-319-49674-0_13
- [6] Heissig B, Salama Y, Takahashi S, Osada T, Hattori K. The multifaceted role of plasminogen in inflammation. Cellular Signalling. 2020 Aug 28;109761. <https://doi.org/10.1016/j.cellsig.2020.109761>
- [7] Hassanpour S, Kim HJ, Saadati A, Tebon P, Xue C, van den Dolder FW, Thakor J, Baradaran B, Mosafer J, Baghbanzadeh A, de Barros NR. Thrombolytic Agents: Nanocarriers in Controlled Release. Small. 2020 Aug 12;2001647. <https://doi.org/10.1002/sml.202001647>
- [8] Li T, Yuan D, Yuan J. Antithrombotic Drugs—Pharmacology and Perspectives. In Coronary Artery Disease: Therapeutics and Drug Discovery 2020 (pp. 101-131). Springer, Singapore. https://doi.org/10.1007/978-981-15-2517-9_4
- [9] Kumar SS, Sabu A. Fibrinolytic enzymes for thrombolytic therapy. In Therapeutic Enzymes: Function and Clinical Implications 2019 (pp. 345-381). Springer, Singapore. https://doi.org/10.1007/978-981-13-7709-9_15

- [10] Devaraj Y, Rajender SK, Halami PM. Purification and characterization of fibrinolytic protease from *Bacillus amyloliquefaciens* MCC2606 and analysis of fibrin degradation product by MS/MS. *Preparative Biochemistry and Biotechnology*. 2018 Feb 7;**48**(2):172-80.
<https://doi.org/10.1080/10826068.2017.1421964>
- [11] Pangestuti R, Arifin Z. Medicinal and health benefit effects of functional sea cucumbers. *Journal of traditional and complementary medicine*. 2018 Jul 1;**8**(3):341-51.
<https://doi.org/10.1016/j.jtcme.2017.06.007>
- [12] Carvalhal F, Cristelo RR, Resende DI, Pinto MM, Sousa E, Correia-da-Silva M. Antithrombotics from the sea: polysaccharides and beyond. *Marine Drugs*. 2019 Mar;**17**(3):170.
<https://doi.org/10.3390/md17030170>
- [13] Hidayati N, Fuad H, Munandar H, Zilda DS, Sulistyaningtyas AR, Nurrahman N, Darmawati S, Ethica SN. Potential of Fibrinolytic Protease Enzymes from Tissue of Sand Sea Cucumber (*Holothuria scabra*) as Thrombolysis Agent. In IOP Conference Series: Earth and Environmental Science 2020 Jul (in press).
- [14] Herliany NE, Nofridiansyah E, Sasongko B. Studi pengolahan teripang kering [Study of Dried sea cucumber processing]. *Jurnal Enggano*. 2016 Sep 5;**1**(2):11-9. Available from:
<https://ejournal.unib.ac.id/index.php/jurnalenggano/article/view/1060>
- [15] Cupp-Enyard C. Sigma's non-specific protease activity assay-casein as a substrate. *JoVE* (Journal of Visualized Experiments). 2008 Sep 17(19):e899. <https://doi.org/10.3791/899>
- [16] Umaña R. Reevaluation of the method of Kunitz for the assay of proteolytic activities in liver and brain homogenates. *Analytical biochemistry*. 1968 Dec 1;**26**(3):430-8.
[https://doi.org/10.1016/0003-2697\(68\)90204-2](https://doi.org/10.1016/0003-2697(68)90204-2)
- [17] Zhang X, Nakahara T, Miyazaki M, Nogi Y, Taniyama S, Arakawa O, Inoue T, Kudo T. Diversity and function of aerobic culturable bacteria in the intestine of the sea cucumber *Holothuria leucospilota*. *The Journal of General and Applied Microbiology*. 2012;**58**(6):447-56.
<https://doi.org/10.2323/jgam.58.447>
- [18] Anggorowati DA, Munandar H, Indriana LF. Isolasi dan Penapisan Bakteri Penghasil Enzim Protease, Selulase, dan Amilase dari Sedimen dan Saluran Pencernaan Teripang Hitam *Holothuria atra* [Isolation and Screening of Protease, Cellulase and Amylase-Producing Bacteria from the Sediment and Digestive System of Black Sea Cucumber *Holothuria atra*]. *Jurnal Ilmu dan Teknologi Kelautan Tropis*. 2019 Aug 21;**11**(2):377-86.
<https://doi.org/10.29244/jitkt.v11i2.21353>
- [19] Khiftiyah AM, Hanifah NN, Bachruddin M, Sholichah MA, Istiqomah S, Dewi SR, Rahayu T, Prasetya IA, Marjayandari L, 'Aini N, Tsana I. Isolation and screening of potential proteolytic and amylolytic microbes from Wonorejo Mangrove forest soil, Surabaya, Indonesia. In AIP Conference Proceedings 2018 Oct 10 (Vol. 2019, No. 1, p. 040014). AIP Publishing LLC.
<https://doi.org/10.1063/1.5061884>
- [20] Ethica SN, Muchlissin SI, Saptaningtyas RA, Sabdono A. Protease producers predominate cultivable hydrolytic bacteria isolated from liquid biomedical waste. *Asian Journal of Chemistry*. 2018;**30**(9):2035-8. <https://doi.org/10.14233/ajchem.2018.21400>
- [21] Ethica SN, Saptaningtyas R, Muchlissin SI, Sabdono A. The development method of bioremediation of hospital biomedical waste using hydrolytic bacteria. *Health and Technology*. 2018 Sep 1;**8**(4):239-54. <https://doi.org/10.1007/s12553-018-0232-8>
- [22] Ethica SN, Sabdono A. Bio-remediation potential of hydrolytic bacteria isolated from hospital liquid biomedical waste in Central Java. In *Proceedings of the 3rd World Congress on New Technologies* 2017 Jun 6. <https://doi.org/2010.11159/icbb17.111>
- [23] Ethica SN, Oedijiono O, Semiarti E, Widada J, Raharjo TJ. Genotypic and Phenotypic Characterization of *Alcaligenes javaensis* JG3 Potential as an Effective Biodegrader. *Biotropia-The Southeast Asian Journal of Tropical Biology*. 2018 Apr 26;**25**(1):1-0.
<http://dx.doi.org/10.11598/btb.2018.25.1.583>

- [24] Gupta P, Samant K, Sahu A. Isolation of cellulose-degrading bacteria and determination of their cellulolytic potential. International journal of microbiology. 2012 Oct;2012.
<https://doi.org/10.1155/2012/578925>
- [25] Naiola E, Widhyastuti N. Isolasi, Seleksi dan Optimasi Produksi Protease dari Beberapa Isolat Bakteri (Isolation, selection and optimization of protease production from several bacterial isolates). Berita Biologi. 2002;6(3):467-73. <http://dx.doi.org/10.14203/beritabiologi.v6i3.1219>
- [26] Prasad S, Kashyap RS, Deopujari JY, Purohit HJ, Taori GM, Dagainawala HF. Development of an *in vitro* model to study clot lysis activity of thrombolytic drugs. Thrombosis Journal. 2006 Dec 1;4(1):14. <https://dx.doi.org/10.1186%2F1477-9560-4-14>
- [27] Bi Q, Chu J, Feng Y, Jiang Z, Han B, Liu W. Purification and characterization of a new serine protease with fibrinolytic activity from the marine invertebrate, *Urechis unicinctus*. Applied biochemistry and biotechnology. 2013 Jun 1;170(3):525-40.
<https://link.springer.com/article/10.1007/s12010-013-0168-4>

Proteolytic and Clot Lysis Activity Screening of Crude Proteases Extracted from Tissues and Bacterial Isolates of Holothuria Scabra

ORIGINALITY REPORT

20%
SIMILARITY INDEX

15%
INTERNET SOURCES

18%
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

1 multisitestaticcontent.uts.edu.au **3%**
Internet Source

2 Sweta Prasad, Rajpal S Kashyap, Jayant Y Deopujari, Hemant J Purohit, Girdhar M Taori, Hatim F Daginawala. "Development of an in vitro model to study clot lysis activity of thrombolytic drugs", Thrombosis Journal, 2006 **2%**
Publication

3 www.bioflux.com.ro **2%**
Internet Source

4 S Darmawati, S I Muchlissin, A R Ernanto, A R Sulistyaningtyas, H Fuad, K M Z Rahman, A Sabdono, S N Ethica. "Pathogenicity Scoring System for Selection of Bacterial Consortium Formulated as Bioremediation Agent of Hospital Wastewater in Central Java", IOP Conference Series: Earth and Environmental Science, 2021 **1%**
Publication

5	www.tandfonline.com Internet Source	1 %
6	ouci.dntb.gov.ua Internet Source	1 %
7	Maheshwari Kumari Singh, R. Usha, K. R. Hithayshree, O. S. Bindhu. "Hemostatic potential of latex proteases from <i>Tabernaemontana divaricata</i> (L.) R. Br. ex. Roem. and Schult. and <i>Artocarpus altilis</i> (Parkinson ex. F.A. Zorn) Forsberg", <i>Journal of Thrombosis and Thrombolysis</i> , 2015 Publication	1 %
8	www.mdpi.com Internet Source	1 %
9	Sara Bordbar, Farooq Anwar, Nazamid Saari. "High-Value Components and Bioactives from Sea Cucumbers for Functional Foods—A Review", <i>Marine Drugs</i> , 2011 Publication	1 %
10	Zhongli Cui, Yukie Maruyama, Bunzo Mikami, Wataru Hashimoto, Kousaku Murata. "Crystallization and preliminary crystallographic analysis of the family GH78 α -rhamnosidase RhaB from sp. GL1 ", <i>Acta Crystallographica Section F Structural Biology and Crystallization Communications</i> , 2006 Publication	1 %

11

ijcmas.com

Internet Source

<1 %

12

repozitorij.svkst.unist.hr

Internet Source

<1 %

13

Asma Shah, Kafeel Ahmad, Ali Talha Khalil, Farhat Amin, Ghosia Lutfullah, Khalid Khan, Ghazala Shah, Aftab Ahmad. " mediated synthesis of gold nanoparticles and its diverse biomedical applications ", Materials Research Express, 2019

Publication

<1 %

14

Varadharajan, Venkatramanan, Sethupathi Shanmugam, and Arulvel Ramaswamy. "Thrombolytic activity of the ethanolic extract of *Vitis vinifera* seeds", Bangladesh Journal of Pharmacology, 2015.

Publication

<1 %

15

link.springer.com

Internet Source

<1 %

16

S Hamdani, N Asstiyani, D Astriany, M Singgih, S Ibrahim. "Isolation and identification of proteolytic bacteria from pig sludge and protease activity determination", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

krishikosh.egranth.ac.in

17

Internet Source

<1 %

18

www.pubfacts.com

Internet Source

<1 %

19

"Encyclopedia of Marine Biotechnology",
Wiley, 2020

Publication

<1 %

20

onlinelibrary.wiley.com

Internet Source

<1 %

21

Martín S. Iglesias, Cynthia Sequeiros,
Sebastián García, Nelda L. Olivera. "Eco-
friendly anti-felting treatment of wool top
based on biosurfactant and enzymes", Journal
of Cleaner Production, 2019

Publication

<1 %

22

www.ijesd.org

Internet Source

<1 %

23

Zhaojun Peng, Yuhong Jin. "Purification,
Identification, and Characterization of an
Endo-1,4- β -Xylanase from Wheat Malt",
Molecules, 2020

Publication

<1 %

24

aip.scitation.org

Internet Source

<1 %

25

hdl.handle.net

Internet Source

<1 %

26

ia904503.us.archive.org

Internet Source

<1 %

27

Xiaolong Zhang, Zeping Liu, Bo Kang, Yao Huang, Caixia Fu, Wei Li, Qian Wu, Zhijie Liu, Dongsheng Li, Chao Wang, Ning Xu. " Effect of or as co - inoculants with in koji making on the physicochemical properties of soy sauce ", Journal of Food Science, 2022

Publication

<1 %

28

daneshyari.com

Internet Source

<1 %

29

Fu, X.y.. "Characterization of proteases from the digestive tract of sea cucumber (Stichopus japonicus): High alkaline protease activity", Aquaculture, 20050518

Publication

<1 %

30

Hirobumi Yamamoto, Mayumi Yamaguchi, Kenichiro Inone. "Absorption and increase in the production of prenylated flavanones in Sophora flavescens cell suspension cultures by cork pieces", Phytochemistry, 1996

Publication

<1 %

31

Mushafau Adebayo Oke, Mohamad Suffian Mohamad Annuar, Khanom Simarani. "Mixed Lignocellulosic Biomass Degradation and Utilization for Bacterial Cellulase Production", Waste and Biomass Valorization, 2016

Publication

<1 %

32

Sercan Özbek Yazici, İsmail Özmen. "Effect of the Crude Saponin Extract from *Gypsophila pilulifera* Boiss. & Heldr. on Protease from *Bacillus subtilis* ATCC 6633 and Antioxidant Properties of the Extract", Iranian Journal of Science and Technology, Transactions A: Science, 2017

Publication

<1 %

33

Ana Mariatul Khiftiyah, Nabela Nur Hanifah, Muhammad Bachruddin, Mar'atus Sholichah et al. "Isolation and screening of potential proteolytic and amylolytic microbes from Wonorejo Mangrove forest soil, Surabaya, Indonesia", AIP Publishing, 2018

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On