

KnE Caribu 11822-Article Text- 58265-1-10-20220914

by Agus Susanto

Submission date: 01-Apr-2023 11:06AM (UTC+0700)

Submission ID: 2052679316

File name: KnE_Caribu_11822-Article_Text-58265-1-10-20220914.pdf (273.77K)

Word count: 2971

Character count: 13829

Conference Paper

Performance of Dairy Cows in Late Lactation Period Recieving Heit Chrose Feed Supplement

4
C H Prayitno*, T Widiyastuti, A N Fitri, and A Susanto

1
Faculty of Animal Science, University of Jenderal Soedirman

ORCID

<https://orcid.org/0000-0002-0920-657X>

Abstract.

The objective was to evaluate Heit Chrose (HC) as a feed supplement for dairy cows, which was measured through digestibility, milk production, milk components and body conditions scores. The research was conducted at UPTD Pagerkukuh Wonosobo using 21 dairy cows in late lactation period (12 cows in 2nd lactation and 9 cows in 4th lactation). The diet provided contained 10.1% crude protein and 60.5% total digestible nutrients. The experimental study was conducted with a nested design. The treatments given were R0: control feed + 0 HC, R1: control feed + 1% HC and R2: control feed + 2% HC. The results showed that HC supplementation did not affect feed consumption but increased the dry matter (DM) and organic matter (OM) digestibility. The HC supplement increased milk production in both 2nd and 4th lactations and increased total solid in 2nd lactation. However, the total solid in 4th lactation decreased due to HC supplementation. In other milk components (fat, protein and SNF) HC supplementation created no significant effect. The conclusion of this research is that supplementing HC at the level of 1% in dairy cows feed during late lactation period can increase total solid in 2nd lactations and tend milk production.

Keywords: Late lactation, Heit Chrose, Dairy Cows

1. Introduction

Dairy cattle nutrient intake in Indonesia tends to be insufficient the nutrient needs of livestock, whoever the average milk production is low. Therefore it is necessary to add additional feed supplements that can increase feed consumption and milk production performance. The Heit-Chrose is one of feed supplement products derived from garlic extract which contains allisin, saponins and organic minerals (Se, Cr and Zn). Saponins and alliscins contained in Heit-Chrose are able to increase propionate production and reduce methane gas production which is a result of carbohydrate fermentation [4]. The addition of Cr can increase the use of glucose in the body, while the Se given in feed will be able to cell protected of oxidative damage during metabolism and Zn-lysinant will increase metabolism proces. Based on this, it is necessary to study how much influence and the optimal level of feeding Heit-Chrose supplements on feed intake and

Corresponding Author: C H
Prayitno; email:
caribuunsoed@gmail.com

Published: 13 Sepetmber 2022

Publishing services provided by
Knowledge E

© C H Prayitno et al. This article

distributed under the terms of
the Creative Commons

Attribution License, which
permits unrestricted use and
redistribution provided that the
original author and source are
credited.

8
Selection and Peer-review under
the responsibility of the ICASI
Conference Committee.

OPEN ACCESS

4
How to cite this article: C H Prayitno*, T Widiyastuti, A N Fitri, and A Susanto, (2022), "Performance of Dairy Cows in Late Lactation Period Recieving Heit Chrose Feed Supplement" in 3rd International Conference on Advance & Scientific Innovation ICASI – Life Sciences Chapter, KnE Life Sciences, Page 383 pages 383–390. DOI 10.18502/kls.v0i0.11822

digestibility as well as milk production and quality. The average peak of milk production in Indonesia occurs in the 5-6th week after parturition, after which milk production will gradually decline. In this study, feed supplements were able to stimulate milk production after the peak production was passed.

2. Material and Methods

The materials were used for the study were twentyone lactating Frisien Holstein dairy cows at 2nd and 4th lactation of month 5th. The treatment were forage (elephant grass) and concentrate with a ratio of 60: 40% and feed supplement with Heit-Chrose levels, each treatment, namely 0%, 1% and 2% (Shown in Table 1 and 2). Feed was given 3 percent of the cow's body weight.

All procedures were performed according to the guiding principles for the care and use of research

animals and were approved by Animal Science Faculty of Jenderal Soedirman University.

TABLE 1: Chemical Composition of Forages, Concentrates and Heit-Chrose feed supplements.

Feedstuff	% DM	Ash	CP	EE	CF	BeTN	TDN
				---- %DM ----			
Elephant grass	22	15.9	11.5	3.2	29.3	40.1	64.24
Concentrats	85.14	13.26	13.77	3.78	17.41	37.95	61.63
Heit-Chrose	93.23	48.75	3.71	0.88	0.56	39.32	44.71

TABLE 2: Nutrient composition of feed treatment.

Treatments	%DM	Ash	CP	EE	CF	BETN	TDN
				---- % DM ----			
R0	47.25	14.24	9.34	2.21	23.65	31.40	63.20
R1	48.19	14.72	9.37	2.22	23.66	31.79	63.65
R2	49.12	15.21	9.41	2.23	23.66	32.19	64.09

The variables observed were feed consumption, feed digestibility, milk production and quality. Milk production was recorded every day for 30 days after the preliminary period. The feed was given based on dry matter intake data obtained in the preliminary period (30 days). The feed was given was weighed, if there was refusal it was also weighed in order to obtain feed consumption data. The refusal was weighed the next

day before feeding and samples were taken (approximately 10%) every day and dried in an oven at 105 ° C for 8 hours for 5 consecutive days.

At the last of the study, the feed samples and the refusal samples were proportionally composited per head. The research method used was experimental using a nested classification design with four replications in group A (lactation 2) and three replications in group B (lactation 4). Research location: Satker Pager Kukuh Wonosobo, belonging to the Livestock Service Office of Central Java Province.

3. Results and Discussion

3.1. Daily feed intake

The results showed that adding of feed supplement feed in dairy cattle feed had no significant effect ($P \geq 0.05$) on feed intake. The average feed intake shown in table 3.

TABLE 3: Average of feed intake dan feed digestibility.

Treatment	Group	Digestibility			
		Feed intake (kg/d)	DMD (%)	OMD(%)	
R0 R1 R2	A (lactation 2 nd)	13.52 + 1.26 13.65 + 1.14 13.50 + 2.43	61.85±5.45 ^a 66.24±6.79 ^b 71.23±4.81 ^c	59.90 ± 6.94 ^a 66.64 ^b 68.50 ± 5.37 ^c	63.04 ±
R0 R1 R2	B (lactation 4 th)	12.18 + 1.83 + 0.95 11.55 0.88	62.63±6.78 ^a 64.54±2.46 ^b 67.44±2.50 ^c	60.32 ± 7.15 ^a 3.86 ^a 66.15 ± 1.77 ^b	61. 60 ±

DMD : Digestibility of dry matter, OMD : organic matter of digestibility

The average of feed intake of group A was 13.56 ± 1.61 kg and the average DM consumption of group B was 11.75 ± 1.22 kg. the DM needs of the lactation group A were R0 with average body weight 464 kg was 13.9 kg, R1 with average body weight 506.6 kg was 15.2 kg, R2 with average body weight 524.5 kg was 15.7 kg, and from lactation group B, R0 with average body weight 464.8 kg was 13 , 9 kg, R1 3% with average body weight 389.9 kg ± 11.7 kg, R2 with average body weight 407.4 kg was 12.2 kg. All treatments received feed according to the NRC standard [3]. DM consumption was influenced by several factors, namely body weight, level of milk production, and quality of feed ingredients [4, 5], it is further explained that, one of the factors affecting consumption is the quality of feed, good quality feed has a relatively high level of consumption compared to quality feed low. From the three treatments, it showed that the consumption of DM in A and B in repetitions of R0, R1 and R2 decreased, in repetitions

of R0 the consumption of DM was 12.95 kg in replications R1 the consumption of DM was 12.74 kg and in tests R2 consumption of DM was 12.67 kg.

3.2. The digestibility of dry matter

The process of digestion of feed in ruminants is carried out by microbes. The rumen requires an optimum condition where the number of bacteria and protozoa is balanced so that fermentation in the rumen is efficient. Research on the addition of Heit-Chrose feed supplements to dairy cattle feed on dry matter digestibility presented in Table 3. The dry matter digestibility in lactation 2nd, respectively, namely R0, R1 and R2 were 59.90%, 63.04% and 68.50%. In lactation 4th, the dry matter digestibility values were 60.32%, 61.60% and 66.15%, respectively. The results of this study inline with other studies [3]. The highest dry matter digestibility value in lactating dairy cows 2 and lactating dairy cows 4 was treatment R2. These results indicated that HC supplementation was able to increase rumen microbial activity. The addition of Heit-Chrose feed supplements showed a significant effect on the digestibility of dry matter and organic matter, this is thought to be the presence of saponins and allisins in Heit-Chrose. Saponins as defaunation agents are able to suppress the growth of rumen protozoa optimally, while allisin compounds are able to suppress methanogenic populations [4].

3.3. The digestibility of organic matter

The digestibility values of the 2 lactating dairy cows, namely R0, R1 and R2 were 61.85%, 66.24% and 71.23%, respectively. In lactation 4, the digestibility values of organic matter, namely R0, R1 and R2, were 62.63%, 64.54% and 67.44%, respectively. The highest organic matter digestibility value in lactating dairy cows 2 and lactating dairy cows 4 was treatment R2.

3.4. Milk Production

The results showed that the average production of FH cow's milk based on period 2 / A in R0 was 5.94 ± 1.166 kg, R1 was 6.80 ± 0.460 kg and R2 was 6.73 ± 0.388 kg and period 4, R0 was 6.52 ± 1.113 kg, R1 was 6.87 ± 0.988 kg and R2 was 7.29 ± 0.867 kg/day/head. The results of this study indicate that the basal feed insufficient of feed its need of dairy cows both in quality and quantity milk, therefor milk production is low. In the treated feed there was an increase in milk production but did not increase optimally,

it is suspected that the supplement feed was unable to stimulate the development of mammary gland cells at the end of the lactation period. The results of this study showed that limited feed energy and protein greatly affect milk production, both in lactation of 2nd or 4th, resulting in low milk production [10]

TABLE 4: Milk production and components.

Variables	Laction- 2 nd			Lactation- 4 th		
	R0	R1	R2	R0	R1	R2
Milk Production (kg/day)	5.94±1.17	6.52±1.11	6.80±0.46	6.87±0.99	6.73±0.39	7.09±0.87
Milk Component (%)						
Fat	5.20±0.22	4.19±0.19	5.29±0.21	5.05±0.23	4.66±0.18	4.62±0.20
Protein	2.53±0.13	2.55±0.17	2.49±0.14	2.49±0.17	2.47±0.10	2.50±0.11
Lactose	4.20±0.20 ^{ab}	4.27±0.24 ^{ab}	4.26±0.12 ^{ab}	4.28±0.27 ^{ab}	3.87±0.15 ^{ac}	3.95±0.16 ^{bc}
Total solids	11.91±0.40 ^{ab}	10.81±0.61 ^{ab}	12.16±1.15 ^{ac}	11.13±0.31 ^{bc}	12.09±0.85 ^{ab}	11.23±0.5 ^{ab}
Solid Non Fat	6.86±0.15	6.90±0.24	6.70±0.13	6.70±0.22	6.51±0.14	6.70±0.17

3.5. Total Solid

The mean of Total Solid (Table 4) the results of this study are considered high in accordance with SNI (2011) which implies that the standard of Total Solid for cow's milk in Indonesia is 11.3%. Total solid is a component of milk which consists of fat content and solid non fat. Fat content and solid non fat are influenced by crude fiber consumed by livestock. The results showed that the feed supplement heit chose had a very significant effect on the levels of total solid ($P < 0.01$). The total solid is influenced by the lactation period. In addition, the mineral Cr also plays a role in protein metabolism, especially the incorporation of amino acids [1]. Therefore, the addition of Heit Chrose can increase milk protein synthesis so that the total solid was increases, because the total solid consists of protein, fat, lactose, vitamins and minerals.

3.6. Solid Non Fat

The results of the average value of solid non-fat research with treatment R0, R1 and R2 are listed in table 3. The average solid non fat (Table 3) the results of this study are low and below the SNI standard (2011) of 7.8%. This is presumably due to the small milk protein content and large fat during the study. The high milk protein content will

increase the non-fat solid value, while the high fat will decrease the non-fat solid value [6].

3.7. Milk Protein

Protein content of milk is one of the parameters determining the quality of fresh milk for dairy cows. The results showed that the addition of HC had no effect on milk protein content. The lowest average protein content was found in group B which received R1 treatment (1% addition of Heit-Chrose feed supplement), while the highest average was in group A who received R1 treatment (1% addition of Heit-Chrose feed supplement). The milk protein content was carried out using a Lactoscan, resulting in the average milk protein content for group A was $2.527 \pm 0.101\%$ and the protein content for group B was $2.476 \pm 0.164\%$. When viewed from the average value of each group, it can be categorized as low quality because the value is below the standard set by SNI (2011), namely milk protein content, namely 2.80%.

The results showed that the average group A (2nd lactation) for R1 (addition of 1% Heit-Chrose feed supplement) was $2.56 \pm 0.101\%$, this figure was greater than R2 (addition of Heit-Chrose feed supplement), namely $2.49 \pm 0.101\%$. Whereas in group B (4th lactation) the R2 was $2.48 \pm 0.164\%$, this figure was greater than R1 which was $2.48 \pm 0.164\%$.

The difference in lactation group is thought not to affect milk protein content, milk protein content is influenced by the month of lactation. This is in accordance with the opinion of [9], that the percentage of milk fat clearly decreases during the first 2-3 months of lactation, then increases in line with the decrease in milk production. The protein content of milk regularly increases according to the month of lactation. From this mechanism, it can be concluded that the protein content is not influenced by the level of lactation but there is an increase in protein levels during the month of lactation [9].

3.8. Milk fat

The results showed that the addition of feed supplements had no significant effect ($P > 0.05$) on milk fat content. The addition of feed supplements in group A (lactation 2) resulted in an increase in the value of milk fat content from 5.05% (R0) to 5.39% (R2), while for group B resulted in a decrease in milk fat from 5.68% (R0) to 4.52% (R2). Feed supplement is given as much as 1-2% of the total dry matter requirement. The levels of fat, protein and lactose in this study were lower than that of other researchers [7, 8]

4. Conclusions

The conclusion of this research is that supplementing HC at the level of 1% in dairy cows feed at late lactation period can increase total solid in 2nd lactations and tend milk production.

References

- [1] Larson BL. Lactation. Ames: The IOWA State University Press; 1985.
- [2] Chen P, Ji P, Cao Z, Li SL. Effects of processing whole cottonseed on nutrient digestibility and ruminal fermentation in dairy cows. *J Anim Feed Sci.* 2007;16(2):537–42.
- [3] NRC (Natl A, editor). Nutrient requirements of dairy cattle, 7th. Washigton: Rev. Sci; 2001.
- [4] Prayitno CH, Yusuf Subagyo A. Nutrient utilization and milk yield of dairy cows fed a diet containing monensin, garlic peel and organic minerals during the lactation period. *Adv Anim Vet Sci.* 2019;7(5):412–6.
- [5] Auldist MJ, O'Brien G, Cole D, Macmillan KL, Grainger C. Effects of varying lactation length on milk production capacity of cows in pasture-based dairying systems. *J Dairy Sci.* 2007;90(7):3234–41.
- [6] McDonald P, Edwards RA, Greenhalgh JF, Morgan CA. *Animal Nutrition* 6th Edition. London (GB): Pearson Education; 2002.
- [7] Iwaniuk ME, Erdman RA. Intake, milk production, ruminal, and feed efficiency responses to dietary cation-anion difference by lactating dairy cows. *J Dairy Sci.* 2015;98(12):8973–85.
- [8] Useni BA, Muller CJC, Cruywagen CW. Milk production of dairy cows as affected by the length of the preceding dry period. *South African Journal of Animal Science.* 2014;44(5).
- [9] Jóźwik A, Strzałkowska N, Bagnicka E, Grzybek W, Krzyżewski J, Poławska E, Kołataj A, Horbańczuk JO. Relationship between milk yield, stage of lactation, and some blood serum metabolic parameters of dairy cows. *Czech J. Anim. Sci.* 2012;57(8): 353–360.
- [10] Rémond B, Ollier A, Miranda G. Milking of cows in late pregnancy: milk production during this period and during the succeeding lactation. *J Dairy Res.* 1992;59(3):233–41.

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

9%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	S Suyadi, A Furqon, W A Septian, R F Putri, C D Nugraha. "The Reproduction Performance in Domestic Animals Controlled by Genetic Mechanism: A Review", KnE Life Sciences, 2022 Publication	2%
2	Submitted to Universitas Jenderal Soedirman Student Paper	1%
3	Submitted to Universitas Diponegoro Student Paper	1%
4	www.knepublishing.com Internet Source	1%
5	sciendo.com Internet Source	1%
6	www.faunajournal.com Internet Source	1%
7	www.ijpab.com Internet Source	1%

8

Nurnaningsih ., Ramadhan Tosepu, Devi Savitri Effendy. "Epidemiology of Ca Mammae Using Surveillance Data in Southeast Sulawesi, Indonesia", KnE Life Sciences, 2022

Publication

<1 %

9

repository.usu.ac.id

Internet Source

<1 %

10

"Proceedings of the XIIIth International Symposium on Ruminant Physiology (ISRP 2019)", Advances in Animal Biosciences, 2019

Publication

<1 %

11

D. J. Minson. "THE INTAKE AND NUTRITIVE VALUE OF FRESH, FROZEN AND DRIED SORGHUM ALMUM, DIGITARIA DECUMBENS AND PANICUM MAXIMUM", Grass and Forage Science, 6/1966

Publication

<1 %

12

www.researcherslinks.com

Internet Source

<1 %

13

Diash Agie Permata, Lucky Radita Alma, Moch Yunus. "The Relationship between Complementary Food After Exclusive Breastfeeding, Immunization and Family Income with Stunting in Kedungrejo, Malang Regency", KnE Life Sciences, 2021

Publication

<1 %

garuda.ristekbrin.go.id

14

Internet Source

<1 %

15

pphouse.org

Internet Source

<1 %

16

Andrea Brenes-Soto, Ellen S. Dierenfeld, Geert P. J. Janssens. "The interplay between voluntary food intake, dietary carbohydrate-lipid ratio and nutrient metabolism in an amphibian, (*Xenopus laevis*)", PLOS ONE, 2018

Publication

<1 %

17

animal.ifas.ufl.edu

Internet Source

<1 %

18

docplayer.hu

Internet Source

<1 %

19

it.scribd.com

Internet Source

<1 %

20

www.animbiosci.org

Internet Source

<1 %

21

www.mdpi.com

Internet Source

<1 %

22

Leman Esra Dolgun, Gülser Köksal. "Effective use of quality function deployment and Kansei engineering for product planning with sensory customer requirements: A plain yogurt case", Quality Engineering, 2017

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On