

20

by Ismoyowati Ismoyowati

Submission date: 01-Oct-2022 09:05PM (UTC+0700)

Submission ID: 1913727896

File name: 20_Egg_Production,_Egg_Quality,_and_Fatty_Acid_Profile.pdf (384.06K)

Word count: 7142

Character count: 38316



Egg Production, Egg Quality, and Fatty Acid Profile of Indonesian Local Ducks Fed with Turmeric, Curcuma, and Probiotic Supplementation

I. Ismoyowati^{a*}, D. Idrasanti^b, A. Ratriyanto^c, & Sumiati^d

^aLaboratory of Poultry Production, Faculty of Animal Science, Jenderal Soedirman University

^bLaboratory of Animal Veterinary, Faculty of Animal Science, Jenderal Soedirman University

Jalan Soeparno Karangwangkal Purwokerto 53122, Indonesia

^cDepartment of Animal Science, Faculty of Agriculture, Sebelas Maret University

Jalan Ir Sutami No. 36A, Jebres, Surakarta 57126, Indonesia

^dDepartment of Nutrition and Feed Technology, Faculty of Animal Science, IPB University

Jalan Agatis, Kampus IPB Dramaga, Bogor, West Java 16680, Indonesia

*Corresponding author: ismoyowati@unsod.ac.id

(Received 05-10-2021; Revised 14-11-2021; Accepted 30-11-2021)

ABSTRACT

Indonesian local ducks are commonly raised for egg production purposes. However, the performances of these ducks are still variable and must be improved. This study investigated the effects of turmeric, curcuma, and probiotic supplementations on the egg production and quality of Indonesian local ducks, emphasizing the eggs' fatty acid profile. Two hundred female local ducks aged 16 weeks were randomly allotted to four dietary treatments with five replicates of 10 birds. The ducks were fed a corn and rice bran-based diet containing different supplements, i.e., a diet without supplementation as the control diet, a diet supplemented with turmeric at the level of 4%, a diet supplemented with curcuma at the level of 4%, and a diet supplemented with starbio probiotics at the level of 2%. The measured data were analyzed using analysis of variance using the 13 Systat program and continued with Duncan's Multiple Range Test. Turmeric supplementation increased egg production compared with the control, and the duck fed probiotics consumed more feed than the control. Curcuma supplementation generated the lowest feed consumption, egg production, and physical egg quality than the other treatments ($p < 0.05$). The probiotics supplementation enhanced the blood high-density lipoprotein concentration ($p < 0.05$). Turmeric, curcuma, and probiotics supplementations generate variable responses in egg production and egg quality, including the fatty acid profile in the eggs. Turmeric and probiotics supplementations positively impact egg production, egg quality, or unsaturated fatty acid profile in the egg. However, curcuma supplementation decreased egg production and egg quality of local ducks. Furthermore, the fatty acid profile was not influenced by these supplements. It is concluded that supplementation of turmeric at the level of 4% and probiotics at the level of 2% in the diet can increase egg production and egg quality of local duck.

Keywords: local duck; egg production; egg quality; unsaturated fatty acids

INTRODUCTION

In Indonesia, duck (*Anas platyrhynchos*) farming is commonly intended for egg production. The population of ducks in Indonesia reached about 50.31 million, with total egg production approximately 329.565.69 tons in 2021 (Directorate General of Animal Husbandry and Health, 2021). Many duck breeds in Indonesia have varying performances, both between breeds and within the same breed. The production rate of laying ducks in Indonesia ranges between 60.2% and 72.9%. It is important to note that duck eggs contain higher lipid concentrations than chicken eggs (Ismoyowati & Sumarmono, 2019). Various efforts can be made to increase duck productivity, including selection, improved maintenance management, and the provision of adequate and balanced nutrients. The inclusion of feed supplements,

such as turmeric, curcuma, and probiotics, can also increase the production and quality of duck eggs.

Turmeric (*Curcuma domestica* Val.) contains some bioactive components, such as curcumin (7.798%), which has antibacterial effects (Prakasita *et al.*, 2019; Basavaraj *et al.*, 2011) and acts as an antioxidant to restore body tissues by activating the cytochrome enzyme P-450 as the catalysator of the oxidation reaction in releasing free radicals. Inactivated cytochrome P-450 can damage tissue due to the depletion of free radicals (Masubuchi & Horie, 2007). Turmeric contains phytoestrogen that can interact with the endocrine system and affect the hypothalamic-hypophyseal-ovarian axis (Sirotkin *et al.*, 2018). In laying ducks, estrogen has a vital role in follicle development. When many follicles are developed, the yolk components, such as cholesterol, are widely distributed throughout the follicles, eventu-

ally decreasing the cholesterol concentrations in the eggs produced (Etches, 1996). Estrogen also stimulates vitellogenin biosynthesis in the liver. As a yolk component, vitellogenin is secreted into the bloodstream and transported into the growing oocytes to be used to synthesize egg yolk (Lu & Zou, 2006).

Curcuma (*Curcuma xanthorrhiza* Roxb.) is a medicinal plant of the rhizome family (Zingiberaceae), a popular ingredient used for seasoning and medicine. Curcuma contains curcuminoid fraction yield at the level of 10.06% from the ethanol extract (Atun *et al.*, 2020), which generates a yellow color in the rhizome and exhibits several properties, such as antibacterial, anticancer, antitumor, and antiinflammation, as well as antioxidant and hypocholesterolemia (Lee *et al.*, 2008).

Some microbial species, including *Bacillus*, *Bifidobacterium*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Streptococcus*, and others, have been used as probiotics. *Lactobacillus* and *Bifidobacterium* have been extensively used in humans, while *Bacillus*, *Enterococcus*, and *Saccharomyces* are mostly used in livestock (Mikulski *et al.*, 2012). Probiotics for laying poultry are intended to increase feed efficiency, egg production, egg quality, intestinal microflora modulation, and inhibit the growth of pathogenic bacteria (Zhu *et al.*, 2015).

Adding feed additives to the diets of laying poultry has been shown to improve feed efficiency, animal health, and productivity (Ismoyowati & Sumarmono, 2019). For example, the inclusion of several Phyto biotics improved the performances of growing ducks (Ismoyowati *et al.*, 2015). Herbal additives (phytogenic and medicinal compounds derived from plants) could act as a substitute for the growth-promoting antibiotic (Masoud-Moghaddam *et al.*, 2021). While the addition of probiotics enhances the laying performance of chickens (Zhang *et al.*, 2017), there are limited studies on the supplementations of turmeric, curcuma, and probiotics in the laying ducks, particularly with an emphasis on egg production, egg quality, and the fatty acid profile of the eggs. It is hypothesized that supplementations of turmeric, curcuma, or probiotics in the diet of laying ducks can increase egg production and egg quality. Therefore, this study aimed to investigate the effect of turmeric, curcuma, and probiotics supplementations in the diet of laying ducks on the egg production and egg quality of laying Indonesian local ducks, emphasizing the fatty acid contents of the eggs.

49

MATERIALS AND METHODS

Ethical Approval

All procedures involving animals were performed by following the Guiding principles Research Animals and were approved by the Institute of Research and Community Service, Jenderal Soedirman University No. 123/UN23.18/PT.01.05/2020.

Experimental Design and Diet

This experiment used 200 Tegal ducks aged 16 weeks with an average body weight of 1.520 ± 86 g. This type of duck is primarily intended for egg production, with the ability to produce 200 eggs per year. The experimental

ducks were allotted to four dietary treatments with five replicates of 10 ducks following a completely randomized design. The four treatments were basal diet without supplementation (control), a basal diet supplemented with 4% turmeric powder (Turmeric), a basal diet supplemented with 4% curcuma powder (Curcuma), and a basal diet supplemented with 2% probiotic (starbio). The herbal additives (turmeric and curcuma) could substitute for the growth-promoting antibiotic, while the addition of probiotics enhanced the laying performances of experimental ducks. The levels of turmeric, curcuma, and probiotic supplementations were based on the most applied levels in the previous studies. The turmeric (*Curcuma domestica* Val.) and curcuma (*Curcuma xanthorrhiza*) powders were made from fresh turmeric and curcuma. They were washed in running water, then thinly sliced (0.5 cm), oven-dried at 55 °C, and ground to powder. The probiotics used in this experiment was Starbio, produced by Lembah Hijau Multifarm Ltd., Solo, Indonesia. Based on information of product specification, Starbio contains various microbes, namely proteolytic (6×10^9 cfu/g *Nitrosococcus*), lignolytic (6×10^9 cfu/g *Clavaria dendroidea*), non-symbiotic nitrogen fixation (4×10^8 cfu/g *Azotobacter spp.*), cellulolytic (8×10^8 cfu/g *Trichoderma polysporum*), and lipolytic (8×10^8 cfu/g *Spirillum liporerum*). The feed ingredients and nutrient composition of the basal diet are presented in Table 1.

The experimental ducks were kept in litter cages with rice husk as the litter. The cage dimension was 2 m × 2 m × 1 m (length × width × height). During the experiment, the cage temperature was 27-29 °C, and the relative humidity was 60%-80%. Lighting was provided for 12 hours. The feed and drinking water were offered *ad libitum*. The experimental ducks were allowed to adapt to the cage and experimental diet for one week, and they were then maintained for 12 weeks during the production period.

Performance Measurements

The feed consumption, egg production (as a percentage of hen day production), and egg weights were measured daily. Feed conversion was calculated from the total feed consumed divided by the egg mass (Mousavi *et al.*, 2016). In total, 400 eggs (20 eggs per replicate) were used for the physical egg quality measurement. The physical egg quality, including the yolk and albumen, yolk color score, and eggshell thickness, and the Haugh Unit were measured according to Stadelman & Cotteril (1995). Meanwhile, four eggs from each treatment unit (in total 80 eggs) were used to analyze the fat and fatty acids contents using the Association of Official Agricultural Chemists (AOAC) method (AOAC, 2012a; AOAC, 2012b) and the cholesterol content using the HPLC method. Fatty acid analysis was performed by composited egg fat extraction from each treatment (from 5 replicates to become one sample), homogenized, and then analyzed. Each treatment was analyzed for fatty acids in duplicate.

Blood Lipid Profile Analyses

Blood samples for lipid profile analyses were collected from two ducks per replicate (in total 40 ducks) after

Table 1. Feed ingredients and nutrient compositions of experimental diet without supplementation (control), supplemented with turmeric powder, curcuma powder, and probiotics

Feed ingredients (%)	Treatments			
	Control	Turmeric	Curcuma	Probiotics
Corn meal	40	38	38	39
Rice bran	32	30	30	31
Soybean waste	15	15	15	15
Fishmeal	10	10	10	10
Turmeric	0	4	0	0
Curcuma	0	0	4	0
Probiotic	0	0	0	2
Palm oil	1	1	1	1
Premix vitamin and mineral	1	1	1	1
Lysine	0.5	0.5	0.5	0.5
47 Methionine	0.5	0.5	0.5	0.5
Nutrient content				
Crude protein (%)	17.62	17.45	17.83	17.92
Metabolizable energy (kcal/kg)	2870	2833	2894	2870
Crude fiber (%)	6.09	6.31	6.11	6.09
Crude fat (%)	4.81	5.11	5.08	4.81
Calcium (%)	2.85	2.83	2.81	2.85
Phosphorus (%)	1.52	1.52	1.52	1.52
Lysine (%)	1.02	1.02	1.02	1.02
Methionine (%)	0.69	0.70	0.71	0.71
Methionine + Cystine (%)	0.90	0.92	0.93	0.93

eight weeks of treatment. The blood sample (3 mL) was collected from the vena axillaries and stored in a sterile mini-tube containing an anticoagulant. Whole blood was centrifuged, and plasma was placed in a minitube and analyzed for lipid profile. The blood lipid profile consisted of triglycerides, total cholesterol, and high-density lipoprotein (HDL) concentrations, were determined by the spectrophotometric method. The contents of low-density lipoprotein (LDL) in the plasma were calculated based on the Friedewald equation (Friedewald *et al.*, 1972).

35

Statistical Analysis

The collected experimental data were subjected to an analysis of variance in a completely randomized design using the Systat 13 program (Systat Software, Inc., San Jose, CA). The Duncan test was employed to measure the difference between treatments at $\alpha = 0.05$ (Steel *et al.*, 1996). The data concerning the fatty acid profile in the egg were presented descriptively since the chemical analysis was performed in a composite manner.

RESULTS

Productive Performance and Physical Egg Quality

Turmeric powder supplementation increased egg production without affecting the egg weight, feed consumption, and feed conversion compared with the control ($p < 0.01$; Table 2). The experimental ducks fed ration supplemented with probiotics consumed more feed than the control (129.17 g vs. 128.15 g; $p < 0.01$). However, this increased feed consumption did not significantly affect the productive performance of the experimental

ducks. On the contrary, supplementation of 4% curcuma powder in the basal diet reduced feed consumption and egg production compared with the other treatments. Consequently, the feed conversion for the experimental ducks fed basal ration supplemented with curcuma powder (Curcuma group) was higher than that of the other treatments ($p < 0.01$).

Supplementation of the basal diet with turmeric powder, curcuma powder, and probiotics did not change the yolk weight and yolk-color score. However, all dietary supplementations generated a lower albumen weight than non-supplemented diet ($p < 0.01$; Table 3). Furthermore, turmeric powder and probiotics supplementations produced a similar Haugh Unit, higher than that in experimental ducks supplemented with curcuma powder ($p < 0.01$). In addition, feeding experimental ducks with basal diets supplemented with turmeric powder and probiotics produced a thicker eggshell, but experimental ducks fed the basal diet supplemented with curcuma powder produced a thinner eggshell than the control ($p < 0.01$).

Blood Lipid Profile and Yolk Fatty Acids Contents

Turmeric powder supplementation in the basal diet decreased blood HDL concentration, while probiotic supplementation increased the blood HDL concentrations compared with the experimental control ducks without supplementation ($p < 0.01$; Table 4). However, total blood cholesterol tended to decrease following turmeric powder, curcuma powder, and probiotic supplementations ($p = 0.098$). In addition, these supplements did not influence the other blood-lipid parameters, such as LDL, triglycerides, yolk fat, and yolk cholesterol.

Table 2. Productive performance of ducks fed basal diet without supplementation (control), supplemented with turmeric powder, curcuma powder, and probiotics

Variables	Treatments				SEM	p Value
	Control	Turmeric	Curcuma	Probiotics		
Feed consumption (g)	128.50 ^b	128.88 ^{ab}	127.48 ^c	129.17 ^a	0.175	<0.001
Egg production (%)	70.71 ^b	79.01 ^a	49.43 ^c	73.86 ^b	4.028	0.002
Egg weight (g)	62.06	63.00	61.49	62.29	0.246	0.182
Feed conversion	2.99 ^b	2.65 ^b	4.28 ^a	2.87 ^b	0.177	<0.001

Note: ^{abc} Means in the same row with different superscripts differ significantly ($p < 0.01$).

Table 3. Physical quality of eggs ducks fed basal diet without supplementation (control), supplemented with turmeric powder, curcuma powder, and probiotics

Variables	Treatments				SEM	p Value
	Control	Turmeric	Curcuma	Probiotics		
Albumen weight (g)	34.40 ^a	32.83 ^b	31.44 ^c	33.08 ^b	0.257	<0.001
Yolk weight (g)	18.41	18.35	18.00	18.39	0.153	0.787
Yolk color	7.86	7.63	7.71	7.40	0.073	0.139
Yolk haugh unit	88.66 ^a	86.52 ^a	81.62 ^b	88.45 ^a	0.860	0.003
Eggshell thickness (mm)	0.43 ^b	0.51 ^a	0.32 ^c	0.54 ^a	0.199	<0.001

Note: ^{abc} Means in the same row with different superscripts differ significantly ($p < 0.01$).

Table 4. Profiles of blood and yolk lipid content of ducks fed basal diet without supplementation (control), supplemented with turmeric powder, curcuma powder, and probiotics

Variables	Treatments				SEM	p Value
	Control	Turmeric	Curcuma	Probiotics		
Total blood cholesterol (mg/dl)	253.70	203.70	209.26	229.63	7.960	0.098
Blood HDL (mg/dl)	65.27 ^b	53.17 ^b	64.53 ^b	91.67 ^a	3.944	<0.001
Blood LDL (mg/dl)	188.44	150.54	144.73	137.96	8.095	0.111
Blood triglyceride (mg/dl)	235.71	226.19	238.09	245.24	15.285	0.981
Yolk fat (%)	33.88	35.21	34.88	35.47	0.374	0.485
Yolk cholesterol (%)	13.66	13.48	12.91	13.99	0.334	0.743

Note: ^{abc} Means in the same row with different superscripts differ significantly ($p < 0.01$).

Furthermore, the fatty acid profiles of the yolk were changed by these supplements (Table 5). It seemed that curcuma powder supplementation generated the lowest. In contrast, probiotic supplementation generated the highest saturated fatty acid (SFA), mono-unsaturated fatty acid (MUFA), and poly-unsaturated fatty acid (PUFA) concentrations, as well as the highest ratios of MUFA/SFA, PUFA/SFA, and (MUFA+PUFA)/SFA. SFA consists of fatty acids that do not contain double bonds, including myristic, pentadecanoic, palmitic, heptadecanoic, stearic, arachidic, behenic, and lignoceric acids. MUFA consists of fatty acids that contain one double bond, including myristoleic, palmitoleic, elaidic, oleic, and eicosenoic acids. PUFA consists of fatty acids that have two or more double bonds, including linoleic, linolenic, g-linolenic, arachidonic, eicosadienoic, eicosatrienoic, eicosapentaenoic, and docosahexaenoic acids.

DISCUSSION

Productive Performance and Physical Egg Quality

Probiotics supplementation resulted in the highest feed consumption, whereas curcuma powder supplementation generated the lowest feed consumption,

although of small magnitudes. Turmeric powder supplementation positively affected duck performance, as indicated by the higher egg production than the other treatments. On the contrary, feeding with curcuma powder supplementation adversely affected the productive performance, as indicated by the lower feed consumption and egg production and the higher feed conversion ratio. Based on the results, turmeric powder supplementation showed the best response on egg production and eggshell thickness of the experimental ducks. A previous study showed that turmeric powder supplementation did not negatively affect the physiological condition and growth of experimental ducks (Zhang *et al.*, 2017; Ismoyowati *et al.*, 2019). In laying quails, turmeric supplementation improved egg production and quality (Putri *et al.*, 2020). Turmeric rhizome contains 2.5%–6.0% essential oil composed of turmerone, alpha and beta turmerone, alpha atlanton, beta caryophyllene, linalool, 1.8 cineol, zingiberene, dd-phellandrene, d-sabinene, and bomeol. It also contains 3%–5% curcuminoid and its derivatives, namely, desmethoxycurcumin and bisdemethoxycurcumin (Hayakawa *et al.*, 2011). Turmeric powder exhibits a beneficial effect on the digestive system by improving mucin secretion, and it acts as a gastro-protectant (Malekizadeh *et al.*,

Table 5. The profiles of egg yolk fatty acids of ducks fed basal diet without supplementation (control), supplemented with turmeric powder, curcuma powder, and probiotics

Fatty acids (%)	Treatments			
	Control	Turmeric	Curcuma	Probiotics
38 Myristic acid (C14:0)	0.31	0.27	0.29	0.33
Myristoleic acid (C14:1)	0.03	0.02	0.03	0.04
Pentadecanoic acid (C15:0)	0.05	0.04	0.06	0.05
Palmitic acid (C16:0)	21.56	19.91	20.97	23.46
23 Mitoleic acid (C16:1)	1.54	1.49	1.42	1.79
Heptadecanoic acid (C17:0)	0.18	0.12	0.18	0.16
Stearic acid (C18:0)	4.90	3.79	4.28	4.81
Elaidic acid (C18:1n9t)	0.18	0.14	0.16	0.18
Oleic acid (C18:1n9c)	41.87	38.97	36.73	43.57
Linoleic acid (C18:2n6c)	7.85	7.88	5.54	10.08
Linolenic acid (C18:3n3)	0.15	0.19	0.08	0.23
33 Linolenic acid (C18:3n6)	0.12	0.14	0.05	0.19
Arachidic acid (C20:0)	0.03	0.02	0.03	0.04
Arachidonic acid (C20:4n6)	2.40	2.95	0.45	3.58
Eicosenoic acid (C20:1)	0.25	0.21	0.17	0.22
Eicosadienoic acid (C20:2)	0.26	0.22	0.22	0.24
Eicosatrienoic acid (C20:3n6)	0.21	0.27	0.07	0.27
23 Psapentaenoic acid (C20:5n3)	0.05	0.03	0.05	0.03
Behenic acid (C22:0)	0.05	0.06	0.04	0.07
Docosahexaenoic acid (C22:6n3)	0.18	0.27	0.05	0.31
Lignoceric acid (C24:0)	0.09	0.04	0.13	0.08
Nervonic acid (C24:1)	0.02	0.02	0.02	0.03
Total fatty acids identified	82.29	77.04	71.00	89.74
SFA	27.17	24.25	25.98	29.00
MUFA	43.86	40.83	38.5	45.79
64 PUFA	11.22	11.95	6.51	14.93
PUFA n-3	0.38	0.49	0.18	0.57
PUFA n-6	10.58	11.24	6.11	14.12
PUFA/SFA	0.41	0.49	0.25	0.51
MUFA/SFA	1.61	1.68	1.48	1.58
(MUFA+PUFA)/SFA	2.03	2.18	1.73	2.09
50 PUFA n-6/n-3	27.84	22.94	33.94	24.77

Note: SFA= saturated fatty acids; MUFA= monounsaturated fatty acids; PUFA= polyunsaturated.

2012). Supplementation of turmeric powder in the basal diet of experimental ducks may improve the concentration of vitellogenin in the serum as a precursor of egg yolk formation, leading to an increase in the number of developing egg follicles (Saraswati *et al.*, 2013).

Moreover, a decrease in egg production following 4% curcuma powder supplementation observed in this study was in contrast with the previous results reported in laying hens (Frita *et al.*, 2017). This decrease can be attributed to the excessive alkaloid content in curcuma powder. Curcuma powder contains an alkaloid, which may disrupt the function of intestinal villi during nutrient absorption, resulting in lower egg productions in the experimental ducks fed a diet supplemented with curcuma powder (Siahaan *et al.*, 2013). The other possible explanation for the decrease in egg production in experimental ducks supplemented with curcuma powder is that the active compounds of alkaloids in the curcuma powder inhibit the rate of vitellogenesis in the liver cells (Van den Berge *et al.*, 2012). The liver functions as a vitellogenesis organ to produce vitellogenin containing fatty acids, which are secreted into the blood vessel

and distributed and deposited into the ovarium as the precursor of yolk follicle formation. Any disorder in the liver function could decrease vitellogenin synthesis and deposition in the ovary, decreasing follicle recruitment, growth, and development and eventually decreasing egg production.

The microbes contained in the probiotics grow in the digestive system and form a bacterial ecosystem that exhibits a positive effect, such as improving the immune system. Probiotics also stimulate physical changes in the intestinal structure, particularly the villus development or the ratio of villus height to crypt depth in the ileum due to the increase in nutrient digestion, leading to the improved availability of protein and fat for egg formation (Peralta-Sánchez *et al.*, 2019). However, the probiotics dosage applied in this study did not positively affect egg production.

Supplementation of the basal diet with curcuma powder reduced the albumen weight because the alkaloid in the curcuma powder reduces the absorption of amino acid from the diet, leading to lower availabilities of amino acids for albumen synthesis (Dei Cas &

Ghidoni, 2019). The egg weight is affected by the weight of the albumen and egg yolk, which is mainly composed of protein (Ratriyanto *et al.*, 2018). The depletion of protein and amino acid in feed during the growth phase could inhibit sex maturity and decrease egg size (Siahaan *et al.*, 2013).

The yolk weight in this study was not affected by the treatments. Similarly, a previous study reported that supplementation of amino acids in feed did not significantly affect the yolk weight of chicken eggs. Apparently, the genetic factor has a more significant effect on yolk weight (Mori *et al.*, 2020). In addition, the contributing factors that affect egg yolk weight are nutrient content, including dietary supplementation and lipoprotein, including phosphatidylcholine and lipovitellin (Ezzat *et al.*, 2011; Hafeez *et al.*, 2016). Again, as was mentioned previously, the main organ producing precursors of vitellogenin is liver. The hepatoprotector effects of turmeric powder will improve liver function to produce vitellogenin as a precursor of egg yolk synthesis in the growing follicles, as was observed in the experimental ducks supplemented with turmeric powder.

Furthermore, turmeric powder, curcuma powder, and probiotics supplementations in feed did not change the score of yolk color (Table 3). This finding suggests that supplementations of experimental ducks with turmeric powder, curcuma powder, and probiotics did not contribute to the pigment, such as carotenoid, which can change the yolk color (Simanjuntak *et al.*, 2013). The yolk's carotenoid is the hydroxy component called xanthophyll, lutein, and zeaxanthin (Jacob & Miles, 2011; Rodriguez-Sanchez *et al.*, 2019). The finding of this study indicated that the feed used in this study contained few coloring substances.

The increased eggshell thickness following turmeric powder and probiotic supplementations was in line with the previous study. The probiotics supplementation containing *Bacillus subtilis* in laying chickens produced 7.5%–8.4% thicker eggshells than the non-supplemented group (Abdelqader *et al.*, 2013). This improvement may be attributed to the increase in the health status of ducks, which is linear to the improved condition in the intestines. The positive impact of probiotics supplementation on eggshell quality is associated with the enhanced availability and digestibility of calcium, as observed by a previous study (Mikulski *et al.*, 2012). Turmeric powder also improves the oviductal environment, particularly the isthmus and uterus, where calcification occurs, leading to improvements in eggshell thickness.

Blood and Yolk Lipid Profile

The tendency to decrease blood cholesterol concentration in this study might be associated with the presence of curcumin in turmeric powder and curcuma powder and the beneficial bacteria in probiotics. Furthermore, probiotics showed the best response in enhancing the HDL level in the blood, which is needed to absorb cholesterol, carry it back to the liver, and flush it from the body. Probiotics supplementation can maintain the balanced composition of microorganisms in the digestive system, improving nutrient digestibility and maintaining

animal health. Curcumin can regulate lipid metabolism in poultry. Furthermore, a study in broilers showed that curcumin alters the lipid profile and decreases the gene expression levels of fatty acid synthase and acetyl CoA carboxylase related to lipogenesis and lipolysis because these genes might be directly responsible for the effect of curcumin on the fatty acid synthesis and lipogenesis (Xie *et al.*, 2019).

Ester cholesterol from feed undergoes hydrolysis, is converted to cholesterol, and is absorbed by the intestines with the non-esterified cholesterol and other lipids (Levy *et al.*, 2015). The blood cholesterol tended to decrease because it is used to synthesize other steroid compounds, such as bile acid or hormones (Gunadi *et al.*, 2021). For instance, dietary supplementation of turmeric powder (Sirotkin *et al.*, 2018), curcuma powder (Liu *et al.*, 2020), or probiotics (Zhang *et al.*, 2019) have also been shown to increase the synthesis of progesterone, luteinizing hormone, estradiol, and follicle-stimulating hormone. The lower cholesterol concentration was due partly to bile salt deconjugation by bacteria, which produces hydrolase enzyme. Probiotics can increase bile acid excretion through feces, leading to a reduced cholesterol concentration in the serum, as cholesterol is a precursor of bile acid (Huang *et al.*, 2013; Rukayadi & Hwang, 2013). Previous studies reported that supplementation of turmeric powder at the level of 3% lowered total blood cholesterol concentrations of experimental ducks (Ismoyowati *et al.*, 2019). Turmeric powder plays a role in decreasing cholesterol levels by improving bile acid production (Xie *et al.*, 2019). The curcumin increases one of the enzymes in the liver, namely, cholesterol-7- α -hydroxylase, which converts cholesterol to bile acid and eventually reduces cholesterol (He *et al.*, 2018).

The tendency to decrease cholesterol concentration may be correlated with the depleting thymoquinone and MUFA in cholesterol synthesis due to hepatocyte or fractional reabsorption in the small intestine (Khalaji *et al.*, 2011). Acetyl CoA is less available in poultry receiving feed with curcumin supplementation, which may decrease the triacylglycerol concentration in the liver. Curcumin directly stimulates beta-oxidation in the hepatocyte, increases the absorption of non-esterified fatty acid, and catalyzes triacylglycerol into fatty acid and glycerol (Xie *et al.*, 2019). However, the present study showed that turmeric powder, curcuma powder, and probiotics supplementations did not affect the fat and cholesterol content in the yolk of duck eggs, indicating that these supplements did not interfere with the fat deposition during yolk formation. The cholesterol level in the blood is less correlated with the level in the egg since it is also a precursor of other substances, such as bile acid or hormones (Liu *et al.*, 2020; Gunadi *et al.*, 2021). Furthermore, curcumin in sufficient quantities can interfere with the reabsorption of cholesterol in the intestinal tract, thereby reducing cholesterol reabsorption. Niemann-Pick protein C1-Like 1 (NPC1L1) is a specific transporter for cholesterol absorption on the surface of the plasma membrane (Feng *et al.*, 2010).

The highest percentage of fatty acid in the ducks' egg yolk is MUFA, followed by SFA, and PUFA. SFA consisted of eight fatty acids. Palmitic acid (C16: 0) showed

a relatively higher concentration than the other SFAs. Probiotics play a role in changing lipid metabolism. The study was in accordance with Milkulski *et al.* (2012), who reported the effect of probiotics on fatty acid composition in chicken egg yolks. Supplementation of 8.0×10^8 cfu/kg *Pediococcus acidilactici* to the diet increased the proportion of PUFAs, including linoleic and linolenic acids. However, changes in the fatty acid profile did not change the ratio of PUFA n-6/PUFA n-3. Lipoprotein metabolism in laying hens may differ, but the information is still limited. The concentration of MUFA was relatively low, except oleic acid (C18:1n9c), which showed a higher concentration than the other fatty acids. The PUFA identified in this study consisted of three n-3 and four n-6 fatty acids. The PUFA ratio of n-6 to n-3 (between 22.94 and 33.94, Table 5) obtained in this study is higher than that observed in a previous study, 7.39 (Sinanoglou *et al.*, 2011).

CONCLUSION

Turmeric powder, curcuma powder, and probiotics supplementations generate variable responses in egg production and egg quality, as well as the fatty acid profile in eggs. Turmeric supplementation increased egg production and eggshell thickness, while probiotics supplementation increased feed consumption, eggshells thickness, and the concentration of MUFA and PUFA. Curcuma powder supplementation decreased egg production and the egg quality of local duck. Finally, supplementation of the basal diet with turmeric powder at the level of 4% or probiotics at the level of 2% can increase egg production and quality of local ducks.

CONFLICT OF INTEREST

We certify no conflict of interest with any financial, personal, or other relationships with other people or organizations related to the material discussed in the manuscript.

ACKNOWLEDGEMENT

Our sincerest gratitude is expressed to the Rector of Universitas Jenderal Soedirman for the financial support through Professor Facilitation Grant No. P No:T/339/UN23.18/PT.01.03/2020.

REFERENCES

- Abdelqader, A., R. Irshaid, & A. R. Al-Fataftah. 2013. Effects of dietary probiotic inclusion on performance, egg shell quality, cecal microflora composition, and tibial traits of laying hens in the late phase of production. *Trop. Anim. Health Prod.* 45: 1017-1024. <https://doi.org/10.1007/s12100-012-0326-7>
- AOAC. 2012a. Analysis of fatty acids. https://www.cfs.gov.hk/english/programme/programme_nifl/files/Analysis_of_Fatty_acids.pdf
- AOAC. 2012b. Analysis of total fat. https://www.cfs.gov.hk/english/programme/programme_nifl/files/Analysis_of_Total_Fat.pdf
- Atun, S., N. Aznam, R. Arianingrum, Senan, B. I. An Naila, A. Lestari, N. A. Pumamaningsih. 2020. Characterization of curcuminoid from *Curcuma xanthorrhiza* and its activity test as antioxidant and antibacterial. *Molekul* 15:79-87. <https://doi.org/10.20884/1.jm.2020.15.2.540>
- Basavaraj, M., V. Nagabhanu, N. Prakash, M. M. Appannavar, P. Wagnare, & S. Mallikarjunappa. 2011. Effect of dietary supplementation of *Curcuma longa* on the biochemical profile and meat characteristics of broiler rabbits under summer stress. *Vet. World*. 4:15-18. <https://doi.org/10.5455/vetworld.2011.15-18>
- Dei Cas, M. & R. Ghidoni. 2019. Dietary curcumin: Correlation between bioavailability and health potential. *Nutrients* 11:2147. <https://doi.org/10.3390/nu11092147>
- Directorate General of Animal Husbandry and Health. 2021. Livestock and Animal Health Statistics. Ministry of Agriculture, Jakarta.
- Etches, R. J. 1996. Reproduction in Poultry. CAB International, Wallingford, UK.
- Ezzat, W. & M. S. Shoeib. 2011. Impact of betaine, vitamin C and folic acid supplementations to the diet on productive and reproductive performance of matrouh poultry strain under Egyptian summer condition. *Egyptian Poultry Science Journal* 31:521-537.
- Feng, D., L. Ohlsson, & R. D. Duan. 2010. Curcumin inhibits cholesterol uptake in Caco-2 cells by downregulation of NPC1L1 expression. *Lipids Health Dis.* 9: 40. <https://doi.org/10.1186/1476-511X-9-40>
- Friedewald, W. T., R. I. Levy, & D. S. Fredrickson. 1972. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clinical Chemistry* 18:499-502. <https://doi.org/10.1093/dinchem/18.6.499>
- Frita, Y., H. Chang, M. J. Lin, & E. Widodo. 2017. Effect of curcuma domestica stock solution on layer performance, egg quality, and antioxidant activity. *International Seminar on Tropical Animal Production* 2017:309-312.
- Gunadi, D., Rukmiasih, & W. Manalu. 2021. Potential Uses of Curcumin and PMSG Hormones to increase egg production of muscovy ducks through increasing estradiol concentrations and shortening laying rest period. *Anim. Prod.* 23:84-94. <https://doi.org/10.20884/1.jap.2021.23.2.98>
- Hafeez, A., A. Mader, I. Ruhnke, K. Manner, & J. Zentek. 2016. Effect of feed grinding methods with and without expansion on prececal and total tract mineral digestibility as well as on interior and exterior egg quality in laying hens. *Poult. Sci.* 95:62-69. <https://doi.org/10.3382/ps/pev316>
- Hayashi, H., Y. Minaniya, K. Ito, Y. Yamamoto, & T. Fukuda. 2011. Difference of curcumin content in *Curcuma longa* L. (*Zingiberaceae*) caused by hybridization with other curcuma species. *Am. J. Plant Sci.* 2:111-119. <https://doi.org/10.4236/ajps.2011.22013>
- He, J., H. Zheng, D. Pan, T. Liu, Y. Sun, J. Cao, Z. Wu, & X. Zeng. 2018. Effects of aging on fat deposition and meat quality in Sheldrake duck. *Poult. Sci.* 97:2005-2010. <https://doi.org/10.3382/ps/pey077>
- Huang, Y., X. Wang, J. Wang, F. Wu, Y. Sui, L. Yang, & Z. Wang. 2013. *Lactobacillus plantarum* strains as potential probiotic cultures with cholesterol-lowering activity. *J. Dairy Sci.* 96:2746-2753. <https://doi.org/10.3168/jds.2012-6123>
- Ismoyowati, I. & J. Sumarmono. 2019. Duck production for food security. *IOP Conf. Ser. Earth Environ. Sci.* 372:012070. <https://doi.org/10.1088/1755-1315/372/1/012070>
- Ismoyowati, I., D. Indrasanti, M. Mufti, & A. S. Farjam. 2015. Phytobiotic properties of garlic, red ginger, turmeric and kencur in growing ducks. *Anim. Prod.* 17:49-55. <https://doi.org/10.20884/1.anprod.2015.17.1.484>
- Ismoyowati, I., D. Indrasanti, S. Mugiyono, & M. Pangestu. 2019. Phytochemical compounds do not interfere physiological parameters and growth performances on two Indonesian local breeds of ducks. *Vet. World*. 12:1689-1697. <https://doi.org/10.14202/vetworld.2019.1689-1697>

- Jacob, J. & R. Miles. 2011. Designer and specialty eggs. The Institute of Food and Agricultural Sciences PS51:1-4.
- Khalaji, S., M. Zaghari, K. H. Hatami, S. Hedari-Dastjerdi, L. Lotfi, & H. Nazarian. 2011. Black cumin seeds, artemisia leaves (*Artemisia sieberi*), and *Camellia* L. plant extract as phyto-genic products in broiler diets and their effects on performance, blood constituents, immunity, and cecal microbial population. *Poult. Sci.* 90:2500-2510. <https://doi.org/10.3382/ps.2011-01393>
- Lee, Y. L., J. S. Shim, Y. Rukayadi, & J. K. Hwang. 2008. Antibacterial activity of xanthorrhizol isolated from *Curcuma xanthorrhiza* Roxb. against foodborne pathogens. *J. Food Prot.* 71:1926-1930. <https://doi.org/10.4315/0362-028X-71.9.1926>
- Levy, A. W., J. W. Kessler, L. Fuller, S. Williams, G. F. Mathis, B. Lumpkins, & F. Valdez. 2015. Effect of feeding an encapsulated source of butyric acid (ButiPEARL) on the performance of male Cobb broilers reared to 42 d of age. *Poult. Sci.* 94:1864-1870. <https://doi.org/10.3382/ps/pev130>
- Liu, M., Y. Lu, P. Gao, X. Xie, D. Li, D. Yu, & M. Yu. 2020. Effect of curcumin on laying performance, egg quality, endocrine hormones, and immune activity in heat-stressed hens. *Poult. Sci.* 99:2196-2202. <https://doi.org/10.1016/j.psj.2019.12.001>
- Lu, J. & X. Zou. 2006. Effects of adding betaine on laying performance and contents of serum yolk precursors VLDL and VTG in laying hen. *J. Zhejiang Univ.* 32:287-291.
- Malekizadeh, M., M. M. Moeini, & S. Ghazi, S. 2012. The effects of different levels of ginger (*Zingiber officinale* Rose) and turmeric (*Curcuma longa* Linn) rhizomes powder on some blood metabolites and production performance characteristics of laying hens. *J. Agric. Sci. Technol.* 14:127-134.
- Masoud-Moghaddama, S., J. Mehrzada, A. H. Alizadeh-amsarib, R. Bahari Kashanic, & J. Saiedi. 2021. Comparison of different herbal additives on immune response and growth performance of broiler chickens. *Trop. Anim. Sci. J.* 44:327-335. <https://doi.org/10.5398/tasj.2021.44.3.327>
- Masubuchi, Y. & T. Horie. 2007. Toxicological significance of mechanism-based inactivation of cytochrome P450 enzymes by drugs. *Crit. Rev. Toxicol.* 37: 389-412. <https://doi.org/10.1080/10408440701215233>
- Mikulski, D., J. Jankowski, J. Naczemski, M. Mikulska, & V. Demey. 2012. Effects of dietary probiotic (*Pediococcus acidilactici*) supplementation on performance, nutrient digestibility, egg traits, egg yolk cholesterol, and fatty acid profile in laying hens. *Poult. Sci.* 91:2691-2700. <https://doi.org/10.3382/ps.2012-02370>
- Mori, H., M. Takaya, K. Nishimura, & T. Goto. 2020. Breed and feed affect amino acid contents of egg yolk and egg-shell color in chickens. *Poult. Sci.* 99:172-178. <https://doi.org/10.3382/ps/pez557>
- Mousavi, S. N., E. Fahimi, & R. Taherkhani. 2016. Effects of different feed forms and cage densities on laying hen performance and stress. *Eur. Poult. Sci.* 80:1-9.
- Peralta-Sánchez, J. M., A. M. Martín-Platero, J. J. Ariza-Romero, M. Rabelo-Ruiz, M. J. Zurita-González, A. Baños, S. M. Rodríguez-Ruano, M. Maqueda, E. Valdivia, & M. Martínez-Bueno. 2019. Egg production in poultry farming is improved by probiotic bacteria. *Front. Microbiol.* 10:1-13. <https://doi.org/10.3389/fmicb.2019.01042>
- Prakasa, V. C., W. Asmara, S. Widyarini, & A. E. T. H. Wahyuni. 2019. Combinations of herbs and probiotics as an alternative growth promoter: An in vitro study. *Vet. World.* 12:614-620. <https://doi.org/10.14202/vetworld.2019.614-620>
- Putri, A. A., A. Widodo, R. Damayanti, & T. W. Suprayogi. 2020. The potency of giving turmeric (*Curcuma domestica* Val.) to the quality of quail (*Coturnix coturnix japonica*) eggs. *Journal of Applied Veterinary Science and Technology* 1:1-5. <https://doi.org/10.20473/javest.V1.I1.2020.1-5>
- Ratriyanto, A., R. Indreswari, R. Dewanti, & S. Wahyuningsih. 2018. Egg quality of quails fed low methionine diet supplemented with betaine. *IOP Conf. Ser. Earth Environ. Sci.* 142:012002. <https://doi.org/10.1088/1755-1315/142/1/012002>
- Rodriguez-Sanchez, R., A. Tres, R. Sala, C. Garcés-Narro, F. Guardiola, J. Gasa, & A. C. Barroeta. 2019. Effects of dietary free fatty-acid content and saturation degree on lipid-class composition and fatty-acid digestibility along the gastrointestinal tract in broiler starter chickens. *Poult. Sci.* 98:4929-4941. <https://doi.org/10.3382/ps/pez253>
- Rukayadi, Y. & J. K. Hwang. 2013. In vitro activity of xanthorrhizol isolated from the rhizome of javanese turmeric (*Curcuma xanthorrhiza* Roxb.) against *Candida albicans* Biofilms. *Phytotherapy Research* 27:1061-1066. <https://doi.org/10.1002/ptr.4834>
- Saraswati, R., W. Manalu, D. R. Ekastuti, & N. Kusumorini. 2013. Increased egg production of Japanese quail (*Coturnix japonica*) by improving liver function through turmeric powder supplementation. *Int. J. Poult. Sci.* 12:601-614. <https://doi.org/10.3923/ijps.2013.601.614>
- Siahaan, N. B., E. Suprijatna, & L. D. Mahfudz. 2013. Effect of red ginger meal (*Zingiber officinale* var. Rubrum) in Kampung Chicken's diet on body weight rate and egg production. *Animal Agriculture Journal* 2:478-488.
- Simanjuntak, R., U. Santoso, & T. Akbarillah. 2013. Pengaruh pemberian tepung daun katuk (*Sauropus androgynus*) dalam ransum terhadap kualitas telur Itik Merah (Anas javanica). *Jurnal Sain Peternakan Indonesia*. 8:65-76. <https://doi.org/10.31186/jspi.id.8.1.65-76>
- Sinaglou, V. J., L. F. Strati, & S. Miniadis-Meimaroglou. 2011. Lipid, fatty acid and carotenoid content of edible egg yolks from 58 species: A comparative study. *Food Chem.* 124:971-977. <https://doi.org/10.1016/j.foodchem.2010.07.037>
- Sirotkin, A. V., A. Kadasi, A. Stochmalova, A. Balazi, M. Földvári, P. Makovicky, P. Chrenek, & A. H. Harrath. 2018. Effect of turmeric on the viability, ovarian folliculogenesis, fecundity, ovarian hormones and response to luteinizing hormone of rabbits. *Animal* 12:242-249. <https://doi.org/10.1017/S175173111700235X>
- Stadelman, W. J. & O. J. Cotteril. 1995. *Egg Science and Technology*. The Haworth Press, Oxon.
- Steel, R. G. D., J. H. Torrie, & D. A. Dickey. 1996. *Principles and Procedures of Statistics: A Biometrical Approach*. McGraw-Hill, New York.
- Van den Berge, V., E. Delezie, P. Delahaut, G. Pierret, P. de Backer, E. Daeseleire, & S. Croubels. 2012. Transfer of flubendazole and tylosin at cross contamination levels in the feed to egg matrices and distribution between egg yolk and egg white. *Poult. Sci.* 91:1248-1255. <https://doi.org/10.3382/ps.2011-02071>
- Xie, Z., G. Shen, Y. Wang, & C. Wu. 2019. Curcumin supplementation regulates lipid metabolism in broiler chickens. *Poult. Sci.* 98:422-429. <https://doi.org/10.3382/ps/pey315>
- Zhang, P., T. Yan, X. Wang, S. Kuang, Y. Xiao, W. Lu, & D. Bi. 2017. Probiotic mixture ameliorates heat stress of laying hens by enhancing intestinal barrier function and improving gut microbiota. *Ital. J. Anim. Sci.* 16:292-300. <https://doi.org/10.1080/1828051X.2016.1264261>
- Zhang, J., Z. Sun, S. Jiang, X. Bai, C. M. Q. Peng, K. Chen, H. Chang, T. Fang, & H. Zhang. 2019. Probiotic *Bifidobacterium lactis* V9 regulates the secretion of sex hormones in polycystic ovary syndrome patients through the gut-brain axis. *mSystems*. 4:e00017-e00019. <https://doi.org/10.1128/mSystems.00017-19>
- Zhu, Y. Z., J. L. Cheng, M. Ren, L. Yin, & X. S. Piao. 2015. Effect of γ -aminobutyric acid-producing *Lactobacillus* strain on laying performance, egg quality and serum enzyme activity in Hy-line brown hens under heat stress. *Asian-Australas. J. Anim. Sci.* 28:1006-1013. <https://doi.org/10.5713/ajas.15.0119>

ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

aac.asm.org

Internet Source

<1 %

2

ejurnal.litbang.pertanian.go.id

Internet Source

<1 %

3

jgeb.springeropen.com

Internet Source

<1 %

4

jwpr.science-line.com

Internet Source

<1 %

5

microbiologyjournal.org

Internet Source

<1 %

6

ojs.jmolekul.com

Internet Source

<1 %

7

Borbon, I.A.. "Lack of efficacy of curcumin on neurodegeneration in the mouse model of Niemann-Pick C1", Pharmacology, Biochemistry and Behavior, 201203

Publication

<1 %

8

Khan, Sohail H.. "The use of green tea (*Camellia sinensis*) as a phytogetic substance

<1 %

in poultry diets", Onderstepoort Journal of Veterinary Research, 2014.

Publication

9

M. Onderci, N. Sahin, K. Sahin, T.A. Balci, M.F. Gursu, V. Juturu, O. Kucuk. "Dietary arginine silicate inositol complex during the late laying period of quail at different environmental temperatures", British Poultry Science, 2007

Publication

<1 %

10

Submitted to Queensland University of Technology

Student Paper

<1 %

11

V. Guyonnet, H. Jing. "23. The eggs and cholesterol: a liability or an opportunity?", Wageningen Academic Publishers, 2016

Publication

<1 %

12

edepot.wur.nl

Internet Source

<1 %

13

www.epa.gov

Internet Source

<1 %

14

www.uaiasi.ro

Internet Source

<1 %

15

Mu-Yeol Cho, Si-Mook Kang, Eun-Song Lee, Baek-II Kim. " Antimicrobial activity of nanoemulsions on biofilms ", Biofouling, 2020

Publication

<1 %

- | | | |
|----|---|------|
| 16 | Muhammad Saeed, Daryoush Babazadeh, Muhammad Naveed, Muhammad Asif Arain, Faiz Ul Hassan, Sun Chao. "Reconsidering betaine as a natural anti-heat stress agent in poultry industry: a review", Tropical Animal Health and Production, 2017
Publication | <1 % |
| 17 | Sissi Wachtel-galor, Yim-tong Szeto, Brian Tomlinson, Iris Ff Benzie. " ('Lingzhi'); acute and short-term biomarker response to supplementation ", International Journal of Food Sciences and Nutrition, 2009
Publication | <1 % |
| 18 | jurnal.unmer.ac.id
Internet Source | <1 % |
| 19 | veterinaryresearch.biomedcentral.com
Internet Source | <1 % |
| 20 | uad.portalgaruda.org
Internet Source | <1 % |
| 21 | www.cfs.gov.hk
Internet Source | <1 % |
| 22 | Damiana D. Rosa, Manoela M. S. Dias, Łukasz M. Grześkowiak, Sandra A. Reis, Lisiane L. Conceição, Maria do Carmo G. Peluzio. " Milk : nutritional, microbiological and health benefits ", Nutrition Research Reviews, 2017
Publication | <1 % |

23	Ong, Y.K.. "The current status and perspectives of biofuel production via catalytic cracking of edible and non-edible oils", Energy, 201001 Publication	<1 %
24	Paul Dent, Laurence Booth, Jane L. Roberts, Andrew Poklepovic, John F. Hancock. "(Curcumin+sildenafil) enhances the efficacy of 5FU and anti - PD1 therapies in vivo", Journal of Cellular Physiology, 2020 Publication	<1 %
25	onesearch.id Internet Source	<1 %
26	K.G. Wiryawan. "Front-matter", Tropical Animal Science Journal, 2019 Publication	<1 %
27	Stelzleni, A.M.. "Benchmarking sensory off-flavor score, off-flavor descriptor and fatty acid profiles for muscles from commercially available beef and dairy cull cow carcasses", Livestock Science, 201006 Publication	<1 %
28	personel.trakya.edu.tr Internet Source	<1 %
29	sinta3.ristekdikti.go.id Internet Source	<1 %

30

Xiaojun Feng, Lei Zhang, Suowen Xu, Ai-zong Shen. "ATP-citrate lyase (ACLY) in lipid metabolism and atherosclerosis: An updated review", Progress in Lipid Research, 2019

Publication

<1 %

31

ejournal.almaata.ac.id

Internet Source

<1 %

32

Aan Andri Yano, Ning Iriyanti, Rosidi. "Effect of supplemental fermeherbafit on total blood lipid and cholesterol in broiler chickens", Journal of Physics: Conference Series, 2018

Publication

<1 %

33

M.Pilar Vaquero, Marcel Veldhuizen, Beatriz Sarriá. "Consumption of an infant formula supplemented with long chain polyunsaturated fatty acids and iron metabolism in rats", Innovative Food Science & Emerging Technologies, 2001

Publication

<1 %

34

Setyawan Setyawan, Junaidi Pangeran Saputra. "Kajian Penambahan Konsentrat Ampas Tahu Terhadap Pertambahan Berat Badan Sapi Peranakan Ongole", Jurnal Pengembangan Penyuluhan Pertanian, 2021

Publication

<1 %

35

repositorio.unesp.br

Internet Source

<1 %

36	www.slideshare.net Internet Source	<1 %
37	Guangliang Gao, Dengfeng Gao, Xianzhi Zhao, Songsong Xu et al. "Genome-Wide Association Study to Identify SNPs and Haplotypes Associated with Reproductive Performance and Egg Quality Traits in Geese", Research Square, 2020 Publication	<1 %
38	Submitted to School of Business and Management ITB Student Paper	<1 %
39	journal2.unusa.ac.id Internet Source	<1 %
40	oalib.perpustakaan.upi.edu Internet Source	<1 %
41	Submitted to University of Adelaide Student Paper	<1 %
42	Submitted to University of Science and Technology Student Paper	<1 %
43	Ayodele Akinyemi, Gustavo Thomé, Vera Morsch, Nathieli Bottari et al. "Effect of Ginger and Turmeric Rhizomes on Inflammatory Cytokines Levels and Enzyme Activities of	<1 %

Cholinergic and Purinergic Systems in Hypertensive Rats", Planta Medica, 2016

Publication

44

Claire d'Andre Hirwa, Jules Mutabazi, Jean De Dieu Nsabimana, Leonidas Dusengemungu et al. "Challenges and opportunities of smallholder pig production systems in Rwanda", Tropical Animal Health and Production, 2022

Publication

<1 %

45

Submitted to Middle Tennessee State University

Student Paper

<1 %

46

Muhammad Risman Wahid, Andhika Yudha Prawira, Chairun Nisa', Srihadi Agungpriyono, Mokhamad Fakhru Ulum. " Sonoanatomy of female reproductive organ of Sunda porcupine () ", Anatomia, Histologia, Embryologia, 2020

Publication

<1 %

47

docplayer.net

Internet Source

<1 %

48

escholarship.org

Internet Source

<1 %

49

journal-cdn.frontiersin.org

Internet Source

<1 %

ijhst.ut.ac.ir

50

Internet Source

<1 %

51

www.e3s-conferences.org

Internet Source

<1 %

52

Submitted to Ana G. Méndez University

Student Paper

<1 %

53

Nialiana Endah Endriastuti, Masita Wulandari Suryoputri, Dewi Latifatul Ilma. "THE EVALUATION OF ANTI-EPILEPTIC DRUGS DOSAGE IN MALNOURISHED CHILDREN: CASES IN INDONESIAN COMMUNITY OF EPILEPSY", Jurnal Farmasi Sains dan Praktis, 2022

Publication

<1 %

54

brain.unboundmedicine.com

Internet Source

<1 %

55

ejournal.forda-mof.org

Internet Source

<1 %

56

M Trenggono, R R Hidayat, T N Cahyo, M D Mahardiono, A D Destrianty. "An assessment Indonesia's Ocean Thermal Energy Conversion (OTEC) as an electrical energy resource", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

57	Ting Ren, Yuming You, Yuanli Luo, Yuanwei Wang, Xiong Liu. " Hypocholesterolemic Effects of Capsaicinoids and Swun5815 Combined by Inhibiting Cholesterol Synthesis and Increasing Bile Acid and Sterols Excretion on Ovariectomized Rats ", Journal of Food Science, 2018 Publication	<1 %
58	cris.vtt.fi Internet Source	<1 %
59	mospi.nic.in Internet Source	<1 %
60	ojs.stiperkutim.ac.id Internet Source	<1 %
61	revistabiomedica.org Internet Source	<1 %
62	www.gcirc.org Internet Source	<1 %
63	M. Rodehutsord, P. Wendt, E. Strobel. "Reducing the phosphorus concentration in diets for turkeys between 10 and 22 weeks of age", British Poultry Science, 2003 Publication	<1 %
64	Pugliese, C., F. Sirtori, A. Acciaioli, R. Bozzi, G. Campodoni, and O. Franci. "Quality of fresh and seasoned fat of Cinta Senese pigs as	<1 %

affected by fattening with chestnut", Meat Science, 2013.

Publication

65

Stéphanie Grizard, Francisco Dini-Andreote, B. Irene Tieleman, Joana F. Salles. "Dynamics of bacterial and fungal communities associated with eggshells during incubation", Ecology and Evolution, 2014

Publication

<1 %

66

Submitted to Thammasat University

Student Paper

<1 %

67

download.garuda.kemdikbud.go.id

Internet Source

<1 %

68

krishikosh.egranth.ac.in

Internet Source

<1 %

69

oasis.postech.ac.kr

Internet Source

<1 %

70

opac.giga-hamburg.de

Internet Source

<1 %

71

www.journals.uchicago.edu

Internet Source

<1 %

72

T D Nova, Y Rizal, S Y Sari. "The Effect Levels of Cassava Leaf and Palm Kernel Meal Fermented Used *Bacillus amylolificans* in Laying Duck Feed", IOP Conference Series: Earth and Environmental Science, 2021

<1 %

73

Wendy A. Mustaqim, Reza Raihandhany,
Muhammad Badrut Tamam. "Chapter 168-1
Phaleria macrocarpa (Scheff.) Boerl.
Thymelaeaceae", Springer Science and
Business Media LLC, 2020

Publication

<1 %

74

Yasuhiro Masubuchi, Toshiharu Horie.
"Toxicological Significance of Mechanism-
Based Inactivation of Cytochrome P450
Enzymes by Drugs", Critical Reviews in
Toxicology, 2008

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off