

# Genetically continuous populations of Striped Snakehead (*Channa striata*) in the Cingcingguling River fragmented by Sempor Reservoir, Central Java, Indonesia

*by Agus Nuryanto*

---

**Submission date:** 20-Mar-2023 05:40PM (UTC+0700)

**Submission ID:** 2041588483

**File name:** 50.\_Nuning\_Setyaningrum\_Januari\_2022.pdf (623.23K)

**Word count:** 7710

**Character count:** 41073

## Genetically continuous populations of Striped Snakehead (*Channa striata*) in the Cingcingguling River fragmented by Sempor Reservoir, Central Java, Indonesia

NUNIN<sup>1,4</sup>, SETYANINGRUM<sup>1</sup>, W. LESTARI<sup>1</sup>, KRISMONO<sup>2</sup>, AGUS NURYANTO<sup>1,\*</sup>

<sup>1</sup>Faculty of Biology, Universitas Jenderal Soedirman, Jl. Dr. Soeparno 63, Purwokerto Utara, Banyumas 53122, Central Java, Indonesia.

Tel.: +62-281-638794, Fax.: +62-281-631700, \*email: agus.nuryanto@unsoed.ac.id

<sup>2</sup>Research Institute for Fisheries Enhancement, Ministry of Marine Affairs and Fisheries, Jl. Cilalawi No. 1, Purwakarta 41152, West Java, Indonesia

Manuscript received: 20 November 2021. Revision accepted: 23 December 2021.

**Abstract.** Setyaningrum N, Lestari W, Nuryanto A. 2022. Genetically continuous populations of Striped Snakehead (*Channa striata*) in the Cingcingguling River fragmented by Sempor Reservoir, Central Java, Indonesia. *Biodiversitas* 23: 222-230. Cingcingguling River, located in Kebumen District, Central Java Province, Indonesia. The Sempor Reservoir fragments it. The previous study proved the negative impact of the reservoir on positive rheotaxis fish, mainly in genetic constituents between the reservoir and river populations. However, research has not been conducted on the negative rheotaxis fish, such as *Channa striata*. Assessing population genetic and taxonomic validity study of Striped Snakehead in the Cingcingguling River is an essential effort. Both studies could be done using the cytochrome c oxidase I gene. Therefore, this research aims to determine taxonomic status and evaluate the population genetic of *C. striata* in the Cingcingguling River. The samples were collected at eight sites, inside and outside the reservoir. The used marker was sequenced from 53 individuals, and all specimens showed high identity (98.67% to 100%) and low genetic distances (0.00 to 0.01) to *C. striata* (KU692421, KU852443, and MG438366). Those values proved that all samples were genetically identified as *Channa striata*. The vertical genetic distribution analysis demonstrated that *C. striata* populations are genetically not different along the river. Unlike rheotaxis positive fish phenomena, the reservoir's existence does not cause genetic fragmentation and leads to continuous striped snakehead populations.

**Keywords:** Genetic diversity, reservoir, rheotaxis, striped snakehead

### INTRODUCTION

Striped snakehead (*Channa striata*) is an important freshwater fish species in several Asian countries. In Indonesia, it is mainly found in the main islands of the Sunda Sh<sup>24</sup> including Sumatra, Java, and Borneo (Adamson et al. 2010; Lakra et al. 2010; Bezinger et al. 2011; Coad et al. 2016). Currently, this species is also found in the lesser Sunda Island, such as Bali (Yudha et al. 2018), and introduced to the Wallacea Regions (Irmawati et al. 2017). *C. striata* is primarily discovered in stagnant or swampy water ecosystems (Amilhat and Lorenzen 2005; Muflikhah 2007; Listyanto and Andriyanto 2009). This species was also found both inside and outside Sempor Reservoir in the Cingcingguling River, Central Java, Indonesia (Setyaningrum et al. 2020; 2021).

Aquatic organisms are able to move in response to water currents, known as rheotaxis (Baker and Montgomery 1999; Kanter and Coombs 2006; Enders et al. 2009). Fish that actively swim against the water current is referred to as positive rheotaxis fish (Suli et al. 2012; Back-Coleman et al. 2015; Oteiza et al. 2017). In the case of *C. striata*, previous studies had reported that *C. striata* also lives in the river, but it could only be found in the parts of the river with stagnant water, river flood plain, and reservoir. It seems that *C. striata* tended to avoid water current (Iskandar and Dahiyat 2012; Nuryanto et al. 2012;

Roesma 2013; Nuryanto et al. 2015). Therefore, *C. striata* could be grouped into negative rheotaxis fish. Fish species that tend to avoid water current are negative rheotaxis fish (Enders et al. 2009; Febrina 2016).

Sempor Reservoir was built approximately 51 years ago and has caused the Cingcingguling River to be fragmented into two extremely different habitats. These include entirely stagnant and running water bodies located underneath the reservoir (Hedianto et al. 2014). The reservoir is a physical barrier for gene flow and causes significant genetic differences among its populations (Heggenes and Roed 2006). However, the available data concerning the reservoir's negative impact on river populations was only available for the positive rheotaxis fish species (Wibowo et al. 2012; Bahiyah et al. 2013; Barasa et al. 2014; Plavova et al. 2017). Meanwhile, there is no recorded information about the reservoir's genetic effect on negative rheotaxis fish species. Therefore, it is essential to research the genetic impact of Sempor Reservoir on the *C. striata* population in the Cingcingguling River.

The genetic impact of a reservoir on the fish population could be assessed with a molecular tool, such as the cytochrome c oxidase I (COI) gene (Nuryanto et al. 2019). Previous research reported that it was used as a robust marker for population genetic analysis of *C. striata* in Perak State situated in Malaysia (Jammaluddin et al. 2011;

Tan et al. 2012; 2015). Nevertheless, population genetic research tends to be carried out when the taxonomic status of the analyzed organisms is valid. In the case of *C. striata*, it was reported that the morphological identification of samples obtained from different regions showed inconsistent diagnostic characters (Zhu et al. 2013; Arma et al. 2014; Khan et al. 2019; Muslimin et al. 2020). Taxonomic validity could also be determined using the cytochrome c oxidase 1 (COI) gene (Ko et al. 2013; Nuryanto et al. 2017; 2019; 2020). Furthermore, it was reliable for species delineation of *C. striata* from Sumatra (Muchlisin et al. 2013; Dahruddin et al. 2016; Irmawati et al. 2017; Syaifudin et al. 2020) and is used to validate morphological identification (Nuryanto et al. 2021). Therefore, this research was aimed to validate and assess the taxonomic status and genetic population of *C. striata* in Sempor Reservoir Central Java using cytochrome c oxidase 1 gene.

## MATERIALS AND METHODS

### Research location and sampling sites


Striped snakehead specimens were collected from eight different sites, four of them were situated inside the reservoir, while the remaining were located downstream (Figure 1). The specimens were collected using traps and lines with the help of fishers. Tissue samples were

chopped from the pectoral fin of each specimen and preserved in ethanol 96%.

### Procedures

#### Genomic DNA extraction and marker polymerization

The total genome was extracted from the pectoral fin tissue using the Quick-DNA™ Miniprep Plus kit adopted from Zymo's research. Extraction procedures were carried out based on the company's manual, and its success was tested using 1% agarose electrophoresis. Subsequently, the COI gene target fragments were reproduced using FishF2 and FishR2 primers (Ward et al. 2005) in Primus 25 Peqlab Thermocycler. Meanwhile, 11 µL of amplified reactions consisted of 1x buffer PCR, 2 mM MgCl<sub>2</sub>, 0.2 mM of each primer, 0.2 mM dNTP mix, 1 U Taq polymerase, and 2.0 ng/µL template DNA. Furthermore, the final volume of 50 µL was adjusted by adding DNA-RNA-free water. Thermal cycles were pre-denatured at 95°C for 4 minutes and were repeated 35 times. The denaturation steps lasted for 30 seconds at 95°C, 2 minutes at 53°C, and 1 minute at 72°C for primer annealing and chain elongation. Additionally, a final extension terminated the cycles after 5 minutes, at 72°C. The PCR products were stained using ethidium bromide and 1.5% agarose gel and placed under ultraviolet light. Gel documentation was further performed using the GelDoc apparatus (BioRad).



**Figure 1.** Research location with five sampling sites along the river in Cingcingguling River, Central Java, Indonesia. Four subsampling sites are located inside the reservoir

### Ma<sup>23</sup> sequencing and editing

The PCR products of the marker were shipped to 1st BASE Malaysia for sequencing, while that the sequencing process was performed using the Sanger method. Consensus and multiple sequences alignment were obtained by assembling the forward and reverse sequences using ClustalW ver.1.4 in Bioedit (Hall 2011). Haplotype data was obtained from its generating process in DnaSP 5 (Rozas et al. 2017).

### Data analysis

The striped snakehead specimens' taxonomic status was validate<sup>8</sup> through a sequence identity test carried out using a basic local alignment search tool (BLAST) closest to the taxa in GenBank. Genetic distance<sup>33</sup> also used to support the identity data. Haplotypes (h) and nucleotide ( $\pi$ )<sup>4</sup> diversities were calculated using Arlequin 3.5, while neutral evolution of the COI marker was estimated using Fu's Fs and Tajima D test (Excoffier and Lischer 2010). Population differentiation was<sup>31</sup> calculated using Fst and variance analysis (AMOVA) in Arlequin 3.5 (Excoffier and Lischer 2010). It was also estimated us<sup>15</sup> a shared haplotype observed in its network. The network was reconstructed using the median-joining method in NETWORK software (Bandelt et al. 1999). The phylogenetic relationships<sup>10</sup> of *C. striata* in Cingcingguling River was estimated using Neighbor-Joining (NJ) and Maximum Likelihood algorithms in MEGA X (Kumar et al. 2018) with 1000 bootstraps replications. Also, the topological stability tree was obtained from the out-group comparison (*Channa gacua* MK599522; *Channa micropeltes* JN024962; *Channa Lucius* KJ937433).

## RESULTS AND DISCUSSION

### Taxonomic status

The COI gene of 53 *Channa* specimens was successfully sequenced, resulting in fragments within the range of 596 bp to 689 bp lengths. Sequence identity test showed that the samples were genetically similar to the top 10 hits closest taxa in the GenBank, all identified as *C. striata* (KU692421, KU852443, and MG438366). However, their percentages were between 98.67% and 100%, with the expected value being 0.0. The samples showed varied genetic distances following Kimura 2 parameter (K2P) from 0.000 % to 1.019, indicating low genetic distances to their closest related taxa in GenBank (Table 1).

This research delineated the samples as *C. striata* because their high genetic identities (above 97%) and genetic distance were less than 3%, respectively, to their conspecific. According to Ratnasingham and Hebert (2013), this is the standard identity threshold for animal species determination. Simultaneously, a distance of 3% is acceptable for threshold species determination in fish barcoding (Ratnasingham and Hebert 2007; Hubert et al. 2010; Candek and Kuntner 2015). Even though a higher threshold of approximately 4% and 5% is allowed, other

factors need to be considered (Higashi et al. 2011; Jeffrey et al. 2011; Candek and Kuntner 2015).

**Table 1.** Sample code, expect value, percent identity, genetic distances, and closest taxa in GenBank

Sample code	E-value	Percent identity (%)	Genetic distance (%)	Closest taxa in GenBank
KW 1	0.0	100.0	0.000	Channa striata KU692421
KW 2	0.0	100.0	0.000	Channa striata KU692421
KW 3	0.0	98.67	1.019	Channa striata KU852443
KW 4	0.0	100.0	0.000	Channa striata KU692421
KW 5	0.0	100.0	0.000	Channa striata KU692421
KW 6	0.0	100.0	0.000	Channa striata KU692421
KW 7	0.0	100.0	0.000	Channa striata KU692421
KW 8	0.0	100.0	0.000	Channa striata KU692421
KW 9	0.0	100.0	0.000	Channa striata KU692421
KW 10	0.0	100.0	0.000	Channa striata KU692421
BK 1	0.0	100.0	0.000	Channa striata KU692421
BK 2	0.0	100.0	0.000	Channa striata KU692421
BK 3	0.0	100.0	0.000	Channa striata KU692421
BK 4	0.0	100.0	0.000	Channa striata KU692421
BK 5	0.0	100.0	0.000	Channa striata KU692421
BK 6	0.0	100.0	0.000	Channa striata KU692421
KA 1	0.0	100.0	0.000	Channa striata KU692421
KA 2	0.0	100.0	0.000	Channa striata KU692421
KA 3	0.0	100.0	0.000	Channa striata KU692421
KA 4	0.0	100.0	0.000	Channa striata KU692421
KA 5	0.0	100.0	0.000	Channa striata MG438366
KA 6	0.0	99.20	0.169	Channa striata KU692421
KA 7	0.0	99.54	0.000	Channa striata KU692421
WO 1	0.0	99.68	0.508	Channa striata KU692421
WO 2	0.0	99.84	0.000	Channa striata KU692421
WO 3	0.0	99.35	0.848	Channa striata MG438366
WO 4	0.0	99.69	0.000	Channa striata KU692421
KS 1	0.0	99.07	0.678	Channa striata KU692421
KS 2	0.0	100.0	0.000	Channa striata KU692421
KS 3	0.0	100.0	0.000	Channa striata KU692421
KS 4	0.0	100.0	0.000	Channa striata KU692421
KS 5	0.0	100.0	0.000	Channa striata KU692421
KS 6	0.0	100.0	0.000	Channa striata KU692421
KS 7	0.0	98.77	0.849	Channa striata KU692421
KS 8	0.0	100.0	0.000	Channa striata KU692421
KS 9	0.0	100.0	0.000	Channa striata KU692421
KS 10	0.0	100.0	0.000	Channa striata KU692421
PW 1	0.0	99.84	0.000	Channa striata KU692421
PW 2	0.0	100.0	0.000	Channa striata KU692421
KB 1	0.0	100.0	0.000	Channa striata KU692421
KB 2	0.0	100.0	0.000	Channa striata KU692421
KB 3	0.0	100.0	0.000	Channa striata KU692421
KB 4	0.0	100.0	0.000	Channa striata KU692421
KB 5	0.0	100.0	0.000	Channa striata KU692421
KB 6	0.0	100.0	0.000	Channa striata KU692421
KB 7	0.0	100.0	0.000	Channa striata KU692421
KB 8	0.0	100.0	0.000	Channa striata KU692421
KB 9	0.0	99.84	0.000	Channa striata KU692421
KB 10	0.0	99.84	0.000	Channa striata KU692421
BA 1	0.0	100.0	0.000	Channa striata KU692421
BA 2	0.0	100.0	0.000	Channa striata KU692421
BA 3	0.0	100.0	0.000	Channa striata KU692421
5-4	0.0	100.0	0.000	Channa striata KU692421

Note: KW: Kedungwringin, BK: Bangkong, KA: Klianget, WO: Waduk outlet, KS: Sempor, PW: Purbowangi, BA: Buayan, KB: Karang Bolong



The low genetic distance among individuals of *C. striata* was reportedly occurred in the wild population found in Lake Towuti, South Sulawesi, with values between 0.043 and 0.309% (Irmawati et al. 2017). Similar values were reported in China (Zhu et al. 2013), using 5 *C. striata* populations, which showed that the intraspecific genetic distances were ranged from 0.002% to 0.027%. In contrast, it was approximately 8 to 45 fold higher than among the species (0.091% to 0.219%). As observed in this research, the minimum (98%) and maximum (1.019%) values of genetic identity and distance, respectively, were reliable to determine the species status of the striped snakehead samples from Cingcingguling Rivers. The present result is consistent with previous research conducted by Aquilino et al. (2011) and Irmawati et al. (2017) that DNA barcoding is a powerful technique for species-level identification of snakehead fish.

Furthermore, the K2P phylogenetic tree was reconstructed by considering neighbor-join and maximum likelihood (Figure 2). Both algorithms produced a similar topology supported by high bootstrap values (ML=100; NJ=100). *C. striata* samples formed a monophyletic clade with their conspecific reference (Figure 2). According to Xu et al. (2015) and Kusbiyanto et al. (2021), monophyly is also reliable data for species determination. Figure 2 shows that the striped snakehead samples and their conspecific were had a smaller branch scale than the predetermined scale of 0.02. This information strongly indicates that the samples belong to the same species as their closest related taxa (*C. striata*). Monophyly of *C. striata* was also detected between the natural and cultivated population in Vietnam (Nguyen and Duong 2015).

This research also indicates that the COI gene is a reliable marker for species identification. Its reliability serves as a barcode because this gene varies among species due to its high mutation rate (Sachithanandam et al. 2012). Due to its variability, the COI gene is a suitable marker for unambiguous species identification (Balkhis et al. 2011; Winarni et al. 2021). This result is congruent with previous research in several locations in Indonesia (Muchlisin et al. 2013; Irmawati et al. 2017; Pramono et al. 2017) and other countries (Aquilino et al. 2011; Triantafyllidis et al. 2011), including Lake Greece.

### Historical demography and genetic diversity

Overall, Tajima's D value was -2.564; meanwhile, this significant result proved that the neutral hypothesis of marker evolution was rejected, thereby leading to selection pressure. However, the negative sign rejected the assumption on selection pressure and indicated a recent population bottleneck (Tajima 1989; Jong et al. 2011). The negative signs and insignificant  $F_{ST}$  values supported the neutral marker and population bottleneck assumption, as shown in Table 2. According to Jong et al. (2011) and Mohammed et al. (2021), Tajima's D and  $F_{ST}$  values are calculated based on haplotype and nucleotide variations, respectively. This difference in the data used simply signifies that  $F_{ST}$  values are more sensitive than Tajima's D in terms of using it for neutral theory testing of the marker.




Figure 2. Phylogenetic tree showing monophyly between samples and their conspecific references. left: ML value, right: NJ value

This research analyzed a 593 bp COI gene fragment length of 53 individual *C. striata* collected from eight sampling sites. Furthermore, it was reported that 17 out of 593 bp were polymorphic, resulting in 6 haplotypes. Overall haplotype and nucleotide diversities were  $0.181 \pm 0.071$ , and  $0.108\% \pm 0.095\%$ , respectively. The data indicate that *C. striata* populations in the Cingcingguling River have low genetic diversity, and this is due to 2 reasons. First, it is caused by small population sizes due to the recent bottleneck. Besides, this has been proven by negative and positive insignificant Tajima's D and  $F_{ST}$  values, respectively, as shown in Table 2. According to

Zanella et al. (2016) and Doublet et al. (2019), a small population shows low genetic diversity due to inbreeding depression. Second, it is caused by limited ancestors. This condition was proven by the haplotype network, which showed that the *C. striata* population in the Cingcingguling River had evolved from a common ancestor, as indicated in Figure 3. It was previously stated that limited maternal ancestors lead to the low genetic diversity of the offspring population because of the drift effect (Zanella et al. 2016). Besides, this attribute in *C. striata* populations was also observed in Malaysia (Jamaluddin et al. 2011).

The present result is inconsistent with the previous research carried out in India (Baisvar et al. 2018), stating that *C. striata* populations exhibited a complex pattern of genetic diversity. However, this implies that it is a common phenomenon. Meanwhile, several fish species have reported high and low haplotype genetic diversity (Sukmanomon et al. 2012; Song et al. 2013; Barasa et al. 2014; Nuryanto et al. 2020). This complex pattern of genetic diversity of *C. striata* populations indicates that environmental factors have exhibited different evolutionary forces on their populations, which needs further analysis.

The within-population evaluation indicates that the haplotype diversity of the *C. striata* population in the Cingcingguling River ranges from 0.000±0.000 to 0.378±0.181. The values prove that the striped snakehead population had low genetic diversity. The majority of the populations underneath the reservoir (PW, KB, and BA) are genetically homogenous. Moreover, two subpopulations (KA and KP) had low genetic diversity. This data indicates that river subpopulations show a complex genetic diversity pattern. The obtained values were lower than the previously reported results (Boonkusol and Tongbai 2016; Baisvar et al. 2018; 2019). The exploitation of *C. Striata* causes low genetic diversity, as indicated by the bottleneck effect shown by Tajimas' D and Fu's Fs values in Table 2, which caused minor population size and an opportunity for inbreeding to occur. According to Hauser et al. (2002) and Tan et al. (2012), fishing pressure reduces genetic diversity

in fish species. Meanwhile, low genetic diversity caused by exploitation also occurs in various aquatic organisms and several regions (Kochzius and Nuryanto 2008; Wibowo 2012; Barasa et al. 2014; Tan et al. 2015; Baisvar et al. 2019).

Table 2 shows that the nucleotide diversity ranges from 0.00±0.000% to 14.568±9.703%, and these values indicate that *C. striata* in the Cingcingguling River has both low and high nucleotide diversities. According to Kochzius and Nuryanto (2008) and Nuryanto et al. (2019), when this attribute is greater than 1%, it is highly diverse. Moreover, high nucleotide diversity was detected in reservoir populations of fish species in Victoria Lake (Barasa et al. 2014).

#### Population connectivity

Per the reservoir population, there was no genetic difference between the four subpopulations. Therefore, this research focused on differentiating the reservoir and river populations. The amova results demonstrated that genetic variations were mainly observed within the population (104.27%, Table 3). It was assumed that no genetic differences occurred between reservoir and river populations along the Cingcingguling River, and this was supported by a negative fixation index (-0.043) and p-values of 0.115. The data proved that *C. striata* populations in the Cingcingguling River formed a genetically continuous population. An interesting finding was that the Sempor Reservoir did not lead to fragmentation, as proven by the genetic similarities among the river populations. The phenomenon is related to the ecological characteristics of striped snakehead as negative rheotaxis fish, which prefer stagnant water ecosystems. Alteration of running water into a static ecosystem due to Sempor Reservoir did not significantly affect the genetics of *C. striata* both inside and outside. Previous studies reported that *C. striata* typically lives in a swampy ecosystem with stagnant water (Amilhat and Lorenzen 2005; Muflikhah 2007; Listyanto and Andriyanto 2009).

**Table 2.** Genetic diversity value and neutrality test for the used marker

Population	N	Genetic diversity			Neutrality test			
		nbp	h (x±SD)	π (x±SD%)	D	P	Fs	P
Overall	53	6	0.181±0.071	0.108±0.095	-2.564***	0.000	-2.360ns	0.070
SR	27	4	0.214±0.103	4.195±3.617	-2.226***	0.001	-0.913ns	0.220
KS	10	3	0.378±0.181	14.568±9.702	-1.901ns	0.006	1.726ns	0.831
PW	2	1	0.000±0.000	0.000±0.000	0.000ns	1.000	-	-
KB	10	1	0.000±0.000	0.000±0.000	0.000	1.000	-	-
BA	22	1	0.000±0.000	0.000±0.000	0.000	1.000	-	-

Note: p> 0.05 = ns, 0.05>p>0.01= significant, p<0.01= highly significant, ns= non-significant, \*\*\*= highly significant

**Table 3.** Variance and Fst analysis among *Channa striata* subpopulation

Source of variation	d.f.	Sum of square	Variance components	Percentage of variation	Fixation index (FST)	p-value
Among subpopulations	4	0.239	-0.004 Va <sup>ns</sup>	-4.27	-0.043 <sup>ns</sup>	0.115±0.003
Within subpopulations	48	4.478	0.093 Vb	104.27		
Total	52	4.717	0.089			

Note: p> 0.05 = non-significant (ns), 0.05>p>0.01= significant, p<0.01= highly significant

The present result is inconsistent with previous research carried out by Song et al. (2013), that significant genetic structure was found among rivers of the *C. striata* in Malaysia. However, this is due to differences in the research locations. This research examined *C. striata* in only one river but fragmented by the reservoir. In contrast, Song et al. (2013) researched different Malaysian river systems. It was previously reported that the river is a closed ecosystem and the freshwater populations tend to show significant genetic differences (Hughes 2009). Therefore, it was reasonable that Song et al. (2013) observed significant genetic differences among rivers. Even Kano et al. (2011) stated solid genetic structures could be observed among tributaries within a river system without physical barriers. However, the different research locations caused an imbalance comparison about genetic differentiation between the present research and that carried out by Kano et al. (2011).

These findings were also inconsistent with the research on *Barbonymus balleroides* in the Serayu River conducted by Bahiyah et al. (2013). According to this research, the significant genetic structures between the reservoir and river population in the Serayu were observed. However, this research evaluated the population genetics of positive rheotaxis species (*B. balleroides*) whose primary habitat is running water. Therefore, the presence of reservoirs in the Serayu River altered the habitat of *B. balleroides* from running to stagnant water, which became an evolutionary factor causing genetic changes in its reservoir populations. Therefore, a significant genetic structure was observed in between *B. balleroides* population inside and outside the reservoir.

It was previously reported, the presence of the Sempor Reservoir has altered upstream areas of Cingcingguling River to become flooded ecosystems or stagnant water ecosystems (Hedianto et al. 2014). Nevertheless, ecosystems alteration in the Cingcingguling River did not change the typical habitat of *C. striata*. It means that *C. striata* collected inside and outside the reservoir live in similar habitat types. Therefore, identical habitat types inside and outside the Sempor Reservoir did not cause genetic fragmentation of *C. striata*.

A detailed analysis of the within-population showed that WO and KS subpopulations showed slightly higher genetic variability than downstream (Table 2). This difference in genetic diversity level might indicate that both subpopulations are less exploited than the lowland river regions. The assumption arises because both subpopulations reside close to reservoirs' outlets with strong outflow. The fishers are prohibited from collecting fish near the outlet because it is hazardous (Setyaningrum et al. 2020). However, due to the age of the reservoir, which is approximately 51 years old (Hedianto et al. 2014), higher genetic variability in the upper stream subpopulations did not cause significant genetic differentiation. Similar phenomena were reported in *Chitala lopis* (Wibowo et al. 2012) and African catfish (Barasa et al. 2014).

Meanwhile, 6 COI haplotypes were observed in the *Channa striata* populations in the Cingcingguling River Central Java, Indonesia. Median-joining analysis proved that haplotype 1 was dominant and found in all subpopulations (Figure 3). The phenomenon strengthens the result of AMOVA that genetic homogeneity occurred along the Cingcingguling River, and the reservoir did not cause genetic fragmentation in the *C. striata* population.




Figure 3. Haplotype network indicates the genetic homogeneity of *Channa striata* in Cingcingguling River, Central Java, Indonesia



The star-like haplotype network (Figure 3) showed that *C. striata* populations in the Cingcingguling River evolved from a single maternal ancestor (H1). The network proved that H1 is the most primitive haplotype, characterized by high abundance and wide distribution in most populations. Similar phenomena were reported in preliminary research on the *Channa* in several regions (Balkhis et al. 2011; Song et al. 2012; Adamson et al. 2012; Baisvar et al. 2018; 2019) and other fish groups (Barasa et al. 2014; Abila et al. 2004).

The snakehead fish in the Cingcingguling River was genetically identified as *Channa striata* and had low genetic diversity. The upper-stream subpopulations had higher genetic diversity than the downstream subpopulations. Also, there were no genetic differences between the reservoir and river populations, which simply means that *C. striata* formed a genetically homogeneous population. These data indicate that *C. striata* need to be treated as a single genetic conservation unit.

### ACKNOWLEDGEMENTS

The authors are grateful to DRPM BRIN for funding this research and to the Faculty of Biology, Universitas Jenderal Soediman (Unsoed), Indonesia for allowing the use of their facilities.

### REFERENCES

- Abila R, Barluenga M, Engelken J, Meyer A, Salzburger W. 2004. Population structure and genetic diversity in a haplochromine fish cichlid of a satellite lake of Lake Victoria. *Mol Ecol* 13 (9): 2589-2602. DOI: 10.1111/j.1365-294X.2004.02270.x.
- Adamson EAS, Hurwood DA, Mather PB. 2010. A reappraisal of the evolution of Asian snakehead fishes (Pisces, Channidae) using molecular data from multiple genes and fossil calibration. *Mol Phylogenet Evol* 56: 707-717. DOI: 10.1016/j.ympev.2010.03.027.
- Adamson EAS, Hurwood DA, Mather PB. 2012. Insights into historical drainage evolution based on the phylogeography of the chevron snakehead fish (*Channa striata*) in the Mekong Basin. *Freshwater Biol* 57 (11): 1-19. DOI: 10.1111/j.1365-2427.2012.02864.x.
- Amilhat E, Lorenzen K. 2005. Habitat use, migration pattern and population dynamics of chevron snakehead *Channa striata* in a rainfed rice farming landscape. *J Fish Biol* 67: 23-34. DOI: 10.1111/j.0022-1112.2005.00927.x.
- Aquilino SVL, Tango JM, Fontanilla IKC, Pagulayan RC, Basiao ZU, Ong PS, Quilang JP. 2011. DNA barcoding of the ichthyofauna of Taal Lake, Philippines. *Mol Ecol Resour* 11 (4): 612-619. DOI: 10.1111/j.1755-0998.2011.03000.x.
- Arma NR, Illijas MA, Irmawati MA. 2014. Morphometric and meristic characteristics of snakehead fish for hatchery production. Proceeding of the 3rd International and National Seminar on Fisheries and Marine Science: Strengthening Science and Technology towards the Development of Blue Economy, Faculty of Fisheries and Marine Science University of Riau, Riau, Indonesia.
- Astuti P. 2018. Dampak sosial ekonomi pembangunan waduk Sempur bagi masyarakat Kebumahan tahun 1958-1978. *Jurnal Prodi Ilmu Sejarah* 3 (5): 1-15. [Indonesian]
- Back-Coleman J, Smith D, Coombs S. 2015. Going with, then against the flow: Evidence against the optomotor hypothesis of fish rheotaxis. *Animal Behaviour* 107: 7-17. DOI: 10.1016/j.anbehav.2015.06.007.
- Bahiyah, Solihin DD, Affandi R. 2013. Variasi genetik populasi ikan brek (*Barbonymus balleroides* Val. 1842) sebagai dampak fragmentasi habitat di Sungai Serayu. *Jurnal Iktiologi Indonesia* 13 (2): 175-186. [Indonesian]
- Baisvar VS, Kumar R, Singh M, Kushwaha B. 2019. Cytochrome c oxidase I gene-based genetic divergence and molecular phylogeny among the species of fish Genus *Channa*. *Proceedings of the National Academy of Sciences India Section B-Biological Sciences* 89 (4): 1455-1463. DOI: 10.1007/s40011-018-01070-w.
- Baisvar VS, Singh M, Kumar R. 2018. Population structuring of *Channa striata* from Indian waters using control region of mtDNA. *Mitochondrial DNA Part A* 30 (3): 414-423. DOI: 10.1080/24701394.2018.1532416.
- Baker, Montgomery 1999. The sensory basis of rheotaxis in the blind Mexican cave fish, *Astyanax fasciatus*. *J Comparative Physiol A* 184: 519-527. DOI: 10.1007/s003590050351.
- Balkhis ABS, Firdaus A, Jamsari J, Hwai TS, Yasin Z. 2011. Evidence of geographical structuring in the Malaysian Snakehead, *Channa striata* based on a partial segment of the CO1 gene. *Genet Mol Biol* 34 (3): 520-523. DOI: 10.1590/S1415-47522011005000016.
- Bandelt HJ, Foster P, Rohlf A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* 16 (1): 37-48. DOI: 10.1093/oxfordjournals.molbev.a026036.
- Barasa JE, Abila R, Grobler JP, Dangasuk OG, Njahira MN. 2014. Genetic diversity and gene flow in *Clarias gariepinus* from Lakes Victoria and Kanyaboli, Kenya. *Afr J Aquatic Sci* 39 (3): 287-293. DOI: 10.2989/16085914.2014.933734.
- Benziger A, Philip S, Raghavan R, Ali PHA, Sukumaran M, Tharian JC, Dahanukar N, Baby F, Peter R, Devi KR, Radhakrishnan KV, Haniffa MA, Britz R, Antunes A. 2011. Unraveling a 146 years old taxonomic puzzle: Validation of malabar snakehead, species-status and its relevance for channid systematics and evolution. *PLoS One* 6 (6): e21272. DOI: 10.1371/journal.pone.0021272.
- Boonkusol D, Tonghai W. 2016. Genetic variation of striped snakehead fish (*Channa striata*) in the river basin of central Thailand inferred from mtDNA COI gene sequences analysis. *J Biol Sci* 16 (1): 37-43. DOI: 10.3923/jbs.2016.37.43.
- Candek K, Kuntner M. 2015. DNA barcoding gap: Reliable species identification over morphological and geographical scales. *Mol Ecol Resour* 15 (2): 268-277. DOI: 10.1111/1755-0998.12304.
- Cia WOC, Asriyana, Halili. 2018. Mortality and exploitation rate of striped snakehead (*Channa striata*) in Aopa Watumohai Swamp, District of Anga. *Jurnal Manajemen Sumber Daya Perairan* 3 (3): 223-231. [Indonesian]
- Coad BW. 2016. Contribution to the knowledge of the snakeheads of Iran (Family Channidae). *Iranian J Ichthyol* 3 (1): 65-72.
- Dahruddin H, Hutama A, Busson F, Sauri S, Hanner R, Keith P, Hadiaty R, Hubert N. 2016. Revisiting the ichthyodiversity of Java and Bali through DNA barcodes: Taxonomic coverage, identification accuracy, cryptic diversity and identification of exotic species. *Mol Ecol Resour* 17 (2): 288-299. DOI: 10.1111/1755-0998.12528.
- Doublet AC, Croiseau P, Fritz S, Michenet A, Hoze C, Danchin-Burge C, Laloe D, Restoux G. 2019. The impact of genomic selection on genetic diversity and genetic gain in three French dairy cattle breeds. *Genet Sel Evol* 51: 52. DOI: 10.1186/s12711-019-0495-1.
- Enders EC, Gessel MH, Williams JG. 2009. Development of successful fish passage structures for downstream migrants requires knowledge of their behavioural response to accelerating flow. *Can J Fish Aquat Sci* 66: 2109-2117. DOI: 10.1139/F09-141.
- Estradivari, Handayani CNN, Daniel D, Mustofa A. 2017. Marine protected area network design: A case study of Maluku Province. *Coastal Ocean J* 1 (2): 135-146. DOI: 10.29244/COJ.1.2.135-146.
- Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 10 (3): 564-567. DOI: 10.1111/j.1755-0998.2010.02847.x.
- Fahmi Z, Nurdawati S, Supriyadi F. 2013. Growth and exploitation status (*Channa striata* Bloch, 1793) in Lubuk Lampam Floodplains, South Sumatera. *Ind Fish Res J* 19 (1): 1-7. DOI: 10.15578/iffj.19.1.2013.1-7.
- Febrina R. 2016. Modeling rheotaxis based on preference to predict fish migration behavior in a river. *Media Komunikasi Teknik Sipil* 22 (1): 23-34. [Indonesian]
- Hall T. 2011. BioEdit: An important software for molecular biology. *GERF Bull Biosci* 2 (1): 60-61. DOI: 10.1017/S0317167100012865.
- Hanif MA, Siddik MAB, Chaklader MR, Nahar A, Mahmud S. 2015. Raznolikost ihtiofaune južnih obalnih voda bangladeša: Postojeće stanje, prijetnje i mogućnosti očuvanja. *Ribarstvo, Croatian J Fisheries* 73 (4): 148-161. DOI: 10.14798/73.4.848.



- Hauser L, Adcock GJ, Smith PJ, Ramirez JHB, Carvalho GR. 2002. Loss of microsatellite diversity and low effective population size in an overexploited population of New Zealand snapper (*Pagrus auratus*). *Proc Nat Acad Sci USA* 99 (18): 11742-11747. DOI: 10.1073/pnas.172242899
- Hedianto DA, Purnomo K, Kartamihardja ES, Warsa A. 2014. Population parameters of three spot cichlid (*Cichlasoma trimaculatum*, Günther 1867) in Sempor Reservoir, Central Java. *Jurnal Penelitian Perikanan Indonesia* 20 (2): 81-88. [Indonesian]
- Heggenes J, Røed KH. 2006. Do dams increase genetic diversity in brown trout (*Salmo trutta*)? Microgeographic differentiation in a fragmented river. *Ecol Freshw Fish* 15 (4): 366-375. DOI: 10.1111/j.1600-0633.2006.00146.x.
- Higashi R, Tsukagoshi A, Kimura H, Kato K. 2011. Male dimorphism in a new interstitial species of the genus *Microloxoconcha* (Podocopa: Ostracoda). *J Crustacean Biol* 31 (1): 142-152. DOI: 10.1651/09-3234.1.
- Hughes JM, Schmidt DJ, Finn DS. 2009. Genes in streams: Using DNA to understand the movement of freshwater fauna and their riverine habitat. *Bio Sci* 59: 573-583. DOI: 10.1525/bio.2009.59.7.8.
- Hubert N, Delieu-Trottin E, Irissou JO, Meyer C, Planes S. 2010. Identifying coral reef fish larvae through DNA barcoding: A test case with the families Acanthuridae and Holocentridae. *Mol Phylogenet Evol* 55: 1195-1203. DOI: 10.1016/j.ympev.2010.02.023.
- Irmawati Tresnati J, Nadianti, Fachrudin L, Arma NR. 2017. Identification of wild stock and the first generation (F1) of domesticated snakehead fish, *Channa striata*. *Jurnal Iktiologi Indonesia* 17 (2): 165-173. [Indonesian]
- Iskandar J, Dahiyat Y. 2012. Keanekaragaman ikan di Sungai Siak Riau. *Bionatura* 14 (1): 51-58. [Indonesian]
- Jamaluddin JAF, Pau TM, Siti-Azizah MN. 2011. Genetic structure of the snakehead murrel, *Channa striata* (Channidae) based on the cytochrome c oxidase subunit I gene: Influence of historical and geomorphological factors. *Genet Mol Biol* 34 (1): 152-160. DOI: 10.1590/S1415-47572011000100026.
- Jeffery NW, Elias-Gutierrez M, Adamowicz SJ. 2011. Species diversity and phylogeographical affinities of the Branchiopoda (Crustacea) of Churchill, Manitoba, Canada. *PLoS One* 6 (5): e18364. DOI: 10.1371/journal.pone.0018364.
- Joadder MAR, Galib SM, Haque SMM, Chaki N. (2015). Fishes of the river Padma, Bangladesh: Current trend and conservation status. *J Fisheries* 3 (2): 259. DOI: 10.17017/jfish.v3i2.2015.111.
- Jong M, Wahlberg N, van Eijk M, Brakefield PM, Zwaan BJ. 2011. Mitochondrial DNA signature for range-wide populations of *Bicyclus anynana* suggests a rapid expansion from recent refugia. *PLoS One* 6 (6): e21385. DOI: 10.1371/journal.pone.0021385.
- Kano Y, Vokoun JC, Letcher BH. 2011. Fine-scale population structure and riverscape genetics of brook trout (*Salvelinus fontinalis*) distributed continuously along headwater channel networks. *Mol Ecol* 20: 3711-3729. DOI: 10.1111/j.1365-294X.2011.05210.x.
- Kanter MJ, Coombs S. 2006. Rheotaxis and prey detection in uniform currents by Lake Michigan mottled sculpin (*Cottus bairdi*). *J Exp Biol* 206: 59-70. DOI: 10.1242/jeb.00056.
- Khan AM, Khan S, Miyan K. 2019. Stock identification of the *Channa striata* inhabiting the Gangetic River system using truss morphometry. *Russian J Ecol* 50 (4): 391-396. DOI: 10.1134/S1067413619040106.
- Ko HL, Wang YT, Chiu TS, Lee MA, Leu MY, Chang KZ, Shao KT. 2013. Evaluating the accuracy of morphological identification of larval fishes by applying DNA barcoding. *PLoS One* 8(1): e053451. DOI: 10.1371/journal.pone.0053451.
- Kochzius M, Nuryanto A. 2008. Strong genetic population structure in the boring giant clam, *Tridacna crocea*, across the Indo-Malay Archipelago: Implications related to evolutionary processes and connectivity. *Mol Ecol* 17: 3775-3787. DOI: 10.1111/j.1365-294X.2008.03803.x.
- Kottelat M, Whitten AJ, Kartikasari SN, Wirjoatmodjo S. 1993. *Freshwater Fishes of Western Indonesia and Sulawesi*. Periplus Editions (HK) Ltd., Hong Kong.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35 (6): 1547-1549. DOI: 10.1093/molbev/msy096.
- Kusbiyanto, Bhagawati D, Nuryanto A. 2020. DNA barcoding of crustacean larvae in Segara Anakan, Cilacap, Central Java, Indonesia using cytochrome c oxidase gene. *Biodiversitas* 21 (10): 4878-4887. DOI: 10.13057/biodiv/d211054.
- Lakra WS, Goswami M, Gopalakrishnan A, Singh DP, Singh A, Nagpure NS. 2010. Genetic relatedness among fish species of Genus *Channa* using mitochondrial DNA genes. *Biochem Syst Ecol* 38: 1212-1219. DOI: 10.1016/j.bse.2010.12.012.
- Listyanto N, Andriyanto S. 2009. Ikan gabus (*Channa striata*) manfaat pengembangan dan alternatif teknik budidayanya. *Media Akuakultur* 4 (1): 18-25. [Indonesian]
- Mohammed MA, Nuryanto A, Kusmintarsih ES. 2021. Genetic differentiation of dengue vector *Aedes aegypti* in the small geographical scale of Banyumas District, Indonesia based on Cytochrome Oxidase I. *Biodiversitas* 22 (2): 675-683. DOI: 10.13057/biodiv/d220219.
- Muchlisin ZA, Thomy Z, Fadli N, Sarong MA, Siti-Azizah MN. 2013. DNA barcoding of freshwater fishes from Lake Laut Tawar, Aceh Province, Indonesia. *Acta Ichthyologica et Piscatoria* 43 (1): 21-29. DOI: 10.3750/AIP.2013.43.1.04.
- Muslimin B, Rustadi R, Hardaningsih H, Retnoaji B. 2020. Morphometric variation of cork fish (*Channa striata* Bloch, 1793) from nine populations in Sumatra Island, Indonesia. *Iranian J Ichthyol* 7 (3): 209-221. DOI: 10.22034/iji.v7i3.480.
- Nguyen T, Duong Y. 2016. Morphological and genetic differences between cultured and wild populations of *Channa striata* in Viet Nam and its phylogenetic relationship with other *Channa* species. *Songklanakarin J Sci Technol* 38 (4): 427-434.
- Muflikhah N. 2007. Domestikasi ikan gabus (*Channa striata*). *Bawal* 1 (5): 169-175. [Indonesian]
- Nurdawati S, Fahmi Z, Supriyadi F. 2019. Parameter populasi ikan betok (*Anabas testudineus* Bloch, 1792) di ekosistem paparan banjir Sungai Musi, Lubuk Lapam. *Berita Biologi Jurnal Ilmu-Ilmu Hayati* 18 (1): 25-35. [Indonesian]
- Nuryanto A, Kusbiyanto K, Bhagawati D. 2021. Molecular barcoding of marine ornamental fish from the southern coast of West Java validates conventional identification. *E3S Web of Conferences* 322: 01004. DOI: 10.1051/e3sconf/202132201004.
- Nuryanto A, Bhagawati D, Husein MOH, Rachmawati FNUR. 2020. Molecular characterization of *Anguilla* from Cibereum and Sapuregel Rivers Segara Anakan Watersheds Cilacap, Central Java. *Biogenesis* 8 (2): 145-156. DOI: 10.24252/bio.v8i2.15532.
- Nuryanto A, Komalawati N, Sugiharto. 2019. Genetic diversity assessment of *Hemibagrus nemurus* from rivers in Java Island, Indonesia using COI gene. *Biodiversitas* 20 (9): 2707-2717. DOI: 10.13057/biodiv/d200936.
- Nuryanto A, Pramono H, Sastranegara MH. 2017. Molecular identification of fish larvae from East Plawangan of Segara Anakan, Cilacap, Central Java, Indonesia. *Biosaintifika* 9 (1): 33-40. DOI: 10.15294/biosaintifika.v9i1.9191.
- Nuryanto A, Bhagawati D, Abulias MA, Indarmawan. 2015. Fauna ikan di Sungai Cikawung Kabupaten Cilacap Jawa Tengah. *Jurnal Iktiologi Indonesia* 15 (1): 25-37. [Indonesian]
- Nuryanto A, Bhagawati D, Abulias MA, Indarmawan. 2012. Fish diversity at Cileumeuh River in District of Majenang, Cilacap District, Central Java. *Jurnal Iktiologi Indonesia* 12 (2): 147-153.
- Oteiza P, Odstreil I, Lauder G, Portugues R, Engert F. 2017. A novel mechanism for mechanosensory-based rheotaxis in larval zebrafish. *Nature* 547 (7664): 445-448. DOI: 10.1038/nature23014.
- Pavlova A, Beheregaray LB, Coleman R, Gilligan D, Harrison KA, Ingram BA, Kearns J, Lamb AM, Lintermans M, Lyon J, Nguyen TTT, Sasai M, Tonkin Z, Yen JDL, Sunnucks P. 2017. Severe consequences of habitat fragmentation on genetic diversity of an endangered Australian freshwater fish: A call for assisted gene flow. *Evol Appl* 10: 531-550. DOI: 10.1111/eva.12484.
- Pramono TB, Arfiati D, Widodo MS, Yanuar U. 2017. Identifikasi ikan Genus *Mystus* dengan pendekatan genetik. *Jurnal Sumberdaya Akuatik Indopasifik* 1 (2): 123-132. DOI: 10.30862/jsai-fpik-unipa.2017.Vol.1.No.2.34. [Indonesian]
- Firlianty, Pratasik SB. 2018. The Potential of *Channa micropeltes* and *C. striata* pudding for healing acceleration of the white rat. *Jurnal Agribisnis Perikanan* 11 (2): 65-69. DOI: 10.29239/j.agrikan.11.2.65-69. [Indonesian]
- Ratnasingham S, Hebert PDN. 2013. A DNA-based registry for all animal species: The barcode index number (BIN) system. *PLoS One* 8 (8): e66213. DOI: 10.1371/journal.pone.0066213.
- Ratnasingham R, Hebert PDN. 2007. BOLD: The barcode of life data system. *Mol Ecol Notes* 7: 355-364. DOI: 10.1111/j.1471-8286.2007.01678.x.

- Roesma DI. 2013. Evaluasi keanekaragaman spesies ikan danau Maninjau. Prosiding Semirata FMIPA Universitas Lampung 197-204. [Indonesian]
- Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Osins SE, Sanchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Mol Biol Evol* 34 (12): 3299-3302. DOI: 10.1093/molbev/msx248.
- Sachithanandam V, Mohan PM, Muruganandam N, Chaaithanya IK, Dhivya P, Baskaaran R. 2012. DNA barcoding, phylogenetic study of *Epinephelus* spp. from Andaman coastal region, India. *Indian J Geo-Marine Sci* 41 (4): 203-211.
- Setyaningrum N, Sugiharto S, Susatyo P. 2020. Species richness and status guilds of fish in Sempor Reservoir Central Java. *Depok* 9 (3): 411-420. DOI: 10.13170/depik.9.3.15094. [Indonesian]
- Setyaningrum, Nuryanto A, Lestari W, Krismono. 2021. Spatial distribution and abundance of *Channa striata* Bloch1793 in Sempor Reservoir, Kebumen Central Java. *E3S Web of Conferences* 322: 01029. DOI: 10.1051/e3sconf/202132201029.
- Sofarini D, Mahmudi M, Hertika AMS, Herawati EY. 2018. Population dynamic of snakehead fish (*Channa striata*) at Danau Panggang Swamp, South Kalimantan. *Enviro Scientiae* 14 (1): 16-20. DOI: 10.20527/es.v14i1.4890.
- Song LM, Munian K, Rashid ZA, Bhassu S. 2013. Characterisation of Asian snakehead murrel *Channa striata* (Channidae) in Malaysia: An insight into molecular data and morphological approach. *Sci World J* 2013: 917506. DOI: 10.1155/2013/917506.
- Sukmanomon S, Senanan W, Kapuscinski AR. 2012. Genetic diversity of feral populations of Nile Tilapia (*Oreochromis niloticus*) in Thailand and evidence of genetic introgression. *Kasetsart J Nat Sci* 46 (2): 200-216.
- Suli A, Watson GM, Rubel EW, Raible DW. 2012. Rheotaxis in larval zebrafish is mediated by lateral line mechanosensory hair cells. *PLoS One* 7 (2): e29727. DOI: 10.1371/journal.pone.0029727.
- Syaifudin M, Wijayanti M, Dwianti SH, Muslim, Mahendra M, Marlina S. 2020. DNA barcodes and phylogenetic of striped snakehead and ocellated snakehead fish from South Sumatra, Indonesia. *Biodiversitas* 21 (3): 1227-1235. DOI: 10.13057/biodiv/d210350.
- Tajima F. 1989. Statistical method for testing then mutation hypothesis by DNA polymorphism. *Genetics* 123: 585-595. DOI: 10.1093/genetics/123.3.585.
- Tan MP, Jamsari AFJ, Muchlisin ZA, Siti-Azizah MN. 2015. Mitochondrial genetic variation and population structure of the striped snakehead, *Channa striata* in Malaysia and Sumatra, Indonesia. *Biochem Syst Ecol* 60: 99-105. DOI: 10.1016/j.bse.2015.04.006.
- Tan MP, Jamsari AFJ, Siti-Azizah MN. 2012. Phylogeographic pattern of the striped snakehead, *Channa striata* in Sundaland: Ancient river connectivity, geographical, and anthropogenic signatures. *PLoS One* 7 (12): e52089. DOI: 10.1371/journal.pone.0052089.
- Triantafyllidis A, Bobori D, Koliimitra C, Gbandi E, Mpanti M, Petriki O, Karaïskou N. 2011. DNA barcoding analysis of fish species diversity in four north Greek lakes. *Mitochondrial DNA* 22: 37-42. DOI: 10.3109/19401736.2010.542242.
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN. 2005. DNA barcoding Australia's fish species. *Philosophical Transactions Royal Society B Biol Sci* 360 (1462): 1847-1857. DOI: 10.1098/rstb.2005.1716.
- Wibowo A. 2012a. Struktur genetik populasi ikan belida (*Chitala lopis*, Bleeker 1851) di Waduk Kutopanjang. *Bawal* 4 (308): 53-58. [Indonesian]
- Wibowo A. 2012b. Genetic diversity of masheer *Tor tambroides* Bleeker 1854) in Manna River Bengkulu and Semangka River Lampung. *Bawal* 4 (2): 105-112. [Indonesian]
- Winarni ET, Kusbiyanto, Nuryanto A. 2021. Estimating Crustacean species utilize Segara Anakan Estuary Cilacap, Indonesia as nursery ground through DNA barcoding. *J Hunan Univ Nat Sci* 48 (10): 275-282.
- Xu X, Liu F, Chen J, Li D, Kuntner M. 2015. Integrative taxonomy of the primitively segmented spider genus *Ganthela* (Araneae: Mesothelae: Liphistiidae): DNA barcoding gap agrees with morphology. *Zool J Linnean Soc* 175 (2): 288-306. DOI: 10.1111/zoj.12280.
- Yudha IGNAP, Arya IW, Suryani SAMP. 2018. Studi identifikasi keragaman jenis, feeding habit dan food habit ikan demersal pada bagian hilir sungai yeh sungi, Tabanan, Bali. *Gema Agro* 23 (2): 114-123. [Indonesian]
- Zanella R, Peixoto JO, Cardoso FF, Cardoso LL, Biegelmeyer P, Cantao ME, Otaviano A, Freitas MS, Caetano AR, Ledur MC. 2016. Genetic diversity analysis of two commercial breeds of pigs using genomic and pedigree data. *Genet Selection Evol* 48: 24. DOI: 10.1186/s12711-016-0203-3.
- Zhu SR, Fu JJ, Wang Q, Li JL. 2013. Identification of *Channa* species using the partial cytochrome c oxidase subunit I (COI) gene as a DNA barcoding marker. *Biochem Syst Ecol* 51: 117-122. DOI: 10.1016/j.bse.2013.08.011.

# Genetically continuous populations of Striped Snakehead (Channa striata) in the Cingcingguling River fragmented by Sempor Reservoir, Central Java, Indonesia

## ORIGINALITY REPORT

12%	%	12%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

## PRIMARY SOURCES

1	"DNA Barcoding and Molecular Phylogeny", Springer Science and Business Media LLC, 2020 Publication	1%
2	Arielli Fabrício Machado, Maria Nazareth Ferreira da Silva, Izeni Pires Farias, Marina Anciães et al. "Recent past connections between Amazonian and Atlantic forests by comparative phylogeography and paleodistribution models for didelphid mammals", Research Square Platform LLC, 2023 Publication	1%
3	Abu Bakar Siti-Balkhis, Amirul Firdaus Jamaluddin Jamsari, Tan Shau Hwai, Zulfigar Yasin, Mohd Nor Siti-Azizah. "Evidence of geographical structuring in the Malaysian Snakehead, Channa striata based on partial segment of the CO1 gene", Genetics and Molecular Biology, 2011	<1%



4

A. Nuryanto, M. Kochzius. "Highly restricted gene flow and deep evolutionary lineages in the giant clam *Tridacna maxima*", Coral Reefs, 2009

Publication

---

5

Nuning Setyaningrum, Agus Nuryanto, W. Lestari, Krismono. " Spatial distribution and abundance of *striata* Bloch1793 in Sempor Reservoir, Kebumen Central Java ", E3S Web of Conferences, 2021

Publication

---

6

Rolando Robert, Noor H. Amit, Nor M. Sukarno, Richard J. Majapun, Subbiah V. Kumar. " Population genetic structure of Asian snakehead fish ( ) in North Borneo: Implications for conservation of local freshwater biodiversity ", Ecological Research, 2018

Publication

---

7

A Iswantari, W Cahyanti, F P Putri, D A Hedianto, V A Prakoso, A H Kristanto. "Reproductive and larval performance of artificially spawned javaen barb *Systemus orphoides* from two populations", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

<1 %

<1 %

<1 %

<1 %

8

Agus Nuryanto, Kusbiyanto Kusbiyanto, Dian Bhagawati. "Molecular barcoding of marine ornamental fish from the southern coast of West Java validates conventional identification", E3S Web of Conferences, 2021

Publication

<1 %

9

Vishwamitra Singh Baisvar, Mahender Singh, Ravindra Kumar. " Population structuring of from Indian waters using control region of mtDNA ", Mitochondrial DNA Part A, 2019

Publication

<1 %

10

Allen Benziger, Siby Philip, Rajeev Raghavan, Palakkaparambil Hamsa Anvar Ali et al. "Unraveling a 146 Years Old Taxonomic Puzzle: Validation of Malabar Snakehead, Species-Status and Its Relevance for Channid Systematics and Evolution", PLoS ONE, 2011

Publication

<1 %

11

A. Nuryanto. "Highly restricted gene flow and deep evolutionary lineages in the giant clam *Tridacna maxima*", Coral Reefs, 03/12/2009

Publication

<1 %

12

Mohamed Ahmed Mohamed Abozeid, Philobater Bahgat Adly Awad, Ahmed Shawki Sharouda, Basma Hussein Abdelaziz Hassan. "Reduction of the Recipient Spleen Size After Living Donor Liver Transplantation A

<1 %

13

Cecilia Conte-Grand, Ralf Britz, Neelesh Dahanukar, Rajeev Raghavan et al. "Barcoding snakeheads (Teleostei, Channidae) revisited: Discovering greater species diversity and resolving perpetuated taxonomic confusions", PLOS ONE, 2017

Publication

<1 %

14

Endang Hilmi, Amron, Dandhy Christianto. "The potential of high tidal flooding disaster in North Jakarta using mapping and mangrove relationship approach", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

15

YanJun Jin, Kun He, Peng Xiang, Xiaodong Wang, Lintian Tong, Zhen Wei, Xiuyue Zhang, Zhaobin Song. "Temporal genetic variation of the Chinese longsnout catfish (*Leiocassis longirostris*) in the upper Yangtze River with resource decline", Environmental Biology of Fishes, 2022

Publication

<1 %

16

Arni Sholihah, Erwan Delrieu-Trottin, Fabien L Condamine, Daisy Wowor et al. "Impact of Pleistocene Eustatic Fluctuations on Evolutionary Dynamics in Southeast Asian

<1 %



17

Sepideh Bazazi, Karin S. Pfennig, Nils Olav Handegard, Iain D. Couzin. "Vortex formation and foraging in polyphenic spadefoot toad tadpoles", Behavioral Ecology and Sociobiology, 2012

Publication

---

18

Tin-Yam Chan, Chien-Hui Yang, Kaori Wakabayashi. "Amended larval recruitment model for the Japanese spiny lobster *Panulirus japonicus* based on new larval records and population genetic data in Taiwan", Journal of Oceanography, 2018

Publication

---

19

Weitao Chen, Nicolas Hubert, Yuefei Li, Denggao Xiang, Xingwei Cai, Shuli Zhu, Jiping Yang, Chuanjiang Zhou, Xinhui Li, Jie Li. "Large - scale DNA barcoding of the subfamily Culterinae (Cypriniformes: Xenocyprididae) in East Asia unveils a geographical scale effect, taxonomic warnings and cryptic diversity", Molecular Ecology, 2022

Publication

---

20

Denis V. Prazdnikov. "Chromosome complements of *Channa lucius* and *C. striata* from Phu Quoc Island and karyotypic

<1 %

<1 %

<1 %

<1 %

21

Lucía Montoliu-Elena, Manuel Elías-Gutiérrez,  
Marcelo Silva-Briano. "Moina macrocopa  
(Straus, 1820): a species complex of a  
common Cladocera, highlighted by  
morphology and DNA barcodes", Limnetica,  
2019

Publication

---

22

Moustafa Sarhan, Ahmed M. Azab, Hassan  
M.M. Khalaf-Allah, Moharam A.M. Afifi. "DNA  
barcoding supports sexual dimorphism in two  
Labrid species; Cheilinus lunulatus and  
Halichoeres hortulanus (Family Labridae) in  
Red Sea, Egypt", The Egyptian Journal of  
Aquatic Research, 2019

Publication

---

23

Muhammad Abu Bakar Abdul-Latiff, Hanisah  
Baharuddin, Pazil Abdul-Patah, Badrul Munir  
Md-Zain. "Is Malaysia's banded langur,  
Presbytis femoralis femoralis, actually  
Presbytis neglectus neglectus? Taxonomic  
revision with new insights on the radiation  
history of the Presbytis species group in  
Southeast Asia", Primates, 2018

Publication

---

<1 %

<1 %

<1 %

24

Nuralif Fakhruallah Mohd Nur, Salwani Abdullah, Ying Giat Seah, Siti Azizah Mohd Nor et al. "DNA Barcoding of Commercially Important Trevallies, Carangoidesspp. (Carangiformes: Carangidae): A Baseline Report of Species from Malaysia", Thalassas: An International Journal of Marine Sciences, 2022

Publication

&lt;1 %

25

"Molecular Genetics and Genomics Tools in Biodiversity Conservation", Springer Science and Business Media LLC, 2022

Publication

&lt;1 %

26

Ingvild Riisberg, Bente Edvardsen. " Genetic variation in bloom-forming ichthyotoxic species (Dictyochophyceae, Heterokonta) using nuclear, mitochondrial and plastid DNA sequence data ", European Journal of Phycology, 2008

Publication

&lt;1 %

27

Mat Jaafar, Tun Nurul Aimi, Martin I. Taylor, Siti Azizah Mohd Nor, Mark de Bruyn, and Gary R. Carvalho. "DNA Barcoding Reveals Cryptic Diversity within Commercially Exploited Indo-Malay Carangidae (Teleostei: Perciformes)", PLoS ONE, 2012.

Publication

&lt;1 %



28

Teddy Triandiza, Agus Kusnadi, Novita Sari, Rosmi Nuslah Pesilette. "KERAGAMAN GENETIK KIMA KECIL (*Tridacna maxima*) DI PULAU KUR, PULAU BIAK, DAN MANADO SERTA IMPLIKASINYA UNTUK KONSERVASI", *Jurnal Penelitian Perikanan Indonesia*, 2020

Publication

&lt;1 %

29

Thuy-Yen Duong, Sophorn Uy, Phen Chheng, Nam So, Thanh-Hien Thi Tran, Ngoc-Tran Thi Nguyen, Robert Pomeroy, Hillary Egna. "Genetic diversity and structure of striped snakehead (*Channa striata*) in the Lower Mekong Basin: Implications for aquaculture and fisheries management", *Fisheries Research*, 2019

Publication

&lt;1 %

30

Yamamoto, Shoichiro, Hitoshi Kubota, Koh Hasegawa, and Tomoyuki Nakamura. "Census and effective population sizes of white-spotted charr (*Salvelinus leucomaenis*) in a fragmented landscape", *Ecology Of Freshwater Fish*, 2015.

Publication

&lt;1 %

31

Congrains, Carlos, Antônio F. Carvalho, Elder A. Miranda, Graeme S. Cumming, Dominic A. W. Henry, Shiiwua A. Manu, Jacinta Abalaka, Cristiano D. Rocha, Moussa S. Diop, Joãozinho Sá, Hamilton Monteiro, Lars H. Holbech,

&lt;1 %

Francis Gbogbo, and Silvia N. Del Lama.  
"Genetic and paleomodelling evidence of the  
population expansion of the cattle egret  
Bubulcus ibis in Africa during the climatic  
oscillations of the Late Pleistocene", Journal of  
Avian Biology, 2016.

Publication

---

32

Hafiz Muhammad Tahir, Alina Noor, Sana  
Mehmood, Sher Muhammad Sherawat,  
Muhammad Akram Qazi. "Evaluating the  
accuracy of morphological identification of  
insect pests of rice crops using DNA  
barcoding", Mitochondrial DNA Part B, 2018

Publication

---

33

L. M. Otwoma, V. Diemel, H. Reuter, M.  
Kochzius, A. Meyer. " Genetic population  
structure of the convict surgeonfish : a  
phylogeographic reassessment across its  
range ", Journal of Fish Biology, 2018

Publication

---

34

Xuhong Chai, Jerusha Bennett, Robert Poulin.  
"Decay of parasite community similarity with  
host phylogenetic and geographic distances  
among deep-sea fish (grenadiers)",  
Parasitology, 2022

Publication

---

35

"Species Diversity of Animals in Japan",  
Springer Science and Business Media LLC,

<1 %

<1 %

<1 %

<1 %

36

Aiguo Zhou, Shaolin Xie, Shulin Liu, Zhuolin Sun, Zhenlu Wang, Yue Zhang, Yanfeng Chen, Mingfei Luo, Jixing Zou. " Genetic diversity of Northern snakehead ( ) based on complete mitochondrial COI gene sequences ", Mitochondrial DNA Part B, 2019

Publication

---

37

M. Afzal Khan, Salman Khan, Kaish Miyan. "Stock identification of the *Channa striata* Inhabiting the Gangetic River System using Truss Morphometry", Russian Journal of Ecology, 2019

Publication

---

38

Nadjim Ahmed Mohamed, Qian Yu, Mohamed Ibrahim Chanfi, Yangping Li, Shi Wang, Xiaoting Huang, Zhenmin Bao. "Genetic diversity and population differentiation of small giant clam *Tridacna maxima* in Comoros islands assessed by microsatellite markers", SpringerPlus, 2016

Publication

---

39

Ricardo Zanella, Jane O. Peixoto, Fernando F. Cardoso, Leandro L. Cardoso et al. "Genetic diversity analysis of two commercial breeds of pigs using genomic and pedigree data", Genetics Selection Evolution, 2016

Publication

&lt;1 %

&lt;1 %

&lt;1 %

&lt;1 %

- 
- 40 Shoujia Jiang, Kai Zhang, Zhiqiang Ruan, Junmin Xu, Xinxin You, Qiong Shi. " The complete mitochondrial genome of Indonesian snakehead, (Channiformes, Channidae) ", Mitochondrial DNA Part B, 2016  
Publication <1 %
- 
- 41 "DNA Barcoding in Marine Perspectives", Springer Nature, 2016  
Publication <1 %
- 
- 42 Anindya Sundar Barman, Mamta Singh, Pramod Kumar Pandey. "DNA barcoding and genetic diversity analyses of fishes of Kaladan River of Indo-Myanmar biodiversity hotspot", Mitochondrial DNA Part A, 2017  
Publication <1 %
- 
- 43 Asadatun Abdullah, Hartmut Rehbein. "DNA barcoding for the species identification of commercially important fishery products in Indonesian markets", International Journal of Food Science & Technology, 2017  
Publication <1 %
- 
- 44 Hadi Alipour, Mohammad R. Bihamta, Valiollah Mohammadi, Seyed A. Peyghambari, Guihua Bai, Guorong Zhang. "Genotyping-by-Sequencing (GBS) Revealed Molecular Genetic Diversity of Iranian Wheat Landraces and Cultivars", Frontiers in Plant Science, 2017  
Publication <1 %
-



45

Shawn M. Lehman, John G. Fleagle. "Primate Biogeography", Springer Science and Business Media LLC, 2006

Publication

<1 %

46

Kaviarasu Munian, Subha Bhassu. "Genetic structure of locally threatened cyprinid, *Osteochilus melanopleurus*, in Peninsular Malaysia River systems inferred from mitochondrial DNA control region", Biochemical Systematics and Ecology, 2015

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On