

# 14.\_Article\_of\_effort\_to\_Reduce \_fat\_and\_cholsterol.pdf

*by*

---

**Submission date:** 23-Mar-2023 12:54AM (UTC-0700)

**Submission ID:** 2044289435

**File name:** 14.\_Article\_of\_effort\_to\_Reduce\_fat\_and\_cholsterol.pdf (186.04K)

**Word count:** 3160

**Character count:** 16081

# Effort to Reduce Fat and Cholesterol of Quail Eggs Using Citronella Oil in Feed

Elly Tugiyanti<sup>1,\*</sup>, and Soegeng Herijanto<sup>2</sup>

<sup>1</sup>Departemen <sup>11</sup>Animal Production, Faculty of Animal Science, University of Jenderal Soedirman

<sup>2</sup>Departemen of Animal Production, Faculty of Animal Science, University of Wijaya Kusuma

\*Corresponding author. Email: [elly.tugiyanti@unsoed.ac.id](mailto:elly.tugiyanti@unsoed.ac.id)

## 1 ABSTRACT

The purpose of this study was to determine the effect of citronella oil supplementation in feed on fat and cholesterol levels of quail eggs (*Coturnix coturnix japonica*). The materials used were 100 female quail *Coturnix-coturnix japonica*, quail commercial feed, commercial citronella oil. The design used was Completely Randomized Design (CRD) with four treatments and five replications. The treatments included citronella oil supplementation 0.3%, 0.6%, 0.9% per kg feed. The experimental research was carried out for 8 weeks. The data obtained were analyzed for variance (ANOVA) and Duncan's Multiple Range Test (DMRT). Analysis of variance showed that the treatment of citronella oil supplementation in the feed had no effect ( $P>0.05$ ) on egg protein but significantly ( $P<0.05$ ) on cholesterol and had a very significant effect ( $P<0.01$ ) on egg fat content. In conclusion, giving citronella oil as much as 0.6% per kg of feed can reduce fat and cholesterol in quail eggs.

**Keywords:** Egg quality, Female quail, Egg fat, Egg cholesterol.

## 1. INTRODUCTION

The current problem is that the productivity of quail eggs is not optimal, one of the causes is low feed efficiency [1]. This adversely affects the quail growth and results in low body weight gain. On the other hand Quail eggs are considered as healthy nutritious food. The growth and production of quail can be optimally if the metabolic process is good. Metabolic results are supported by the amount of feed consumed and the optimization of feed use.

The food consumed has better digestibility with the presence of essential oils in the feed, so that the nutrients in the feed are easily absorbed. Lemongrass essential oil contains antiseptic and antibacterial properties so that it can kill pathogenic bacteria in the digestive tract [2,3]. [4] explained that the content of citronellal, geraniol, and citronellol in Citronella oil can inhibit bacterial activity.

State that Citronella oil shows great potential as an antibacterial agent that can suppress the activity of bacteria [5,6]. Putriningtyas [7] in her study reported that citronella leaf essential oil from Tawangmangu was able to produce an inhibitory zone against *S. aureus* and *E. coli*. That essential oils can reduce the population of pathogenic bacteria that can damage the villi, and

increase the population of lactic acid bacteria which can increase the density of the villi, so that the number of villi increases, and nutrients can be absorbed completely [8].

Reported that supplements an herbal essential oil mixture on the diet can overcome coccidiosis broiler chickens [9]. The study results by [10] so that the essential oil found in lemongrass can increase body weight gain and function as an anti-bacterial, especially in the digestive tract, thereby increasing growth livestock.

Citronella oil extract is able to increase broiler body weight gain [11]. The highest weight gain was achieved in broilers who consumed rations with Citronella oil extract 0.5%, which was 1.160 g and tended to decrease in Citronella oil extract treatment, and the smallest body weight was obtained in broiler chickens that were given control rations which only reached 986 g. These results are contradictory with the finding of [12] that broilers fed diets treated with various levels of lemongrass oil (LGO) consumed significantly more amount of feed compared to the broilers fed control, but which was statistically non significant.

Supplementation of lemongrass leaf meal powder at the various levels in the broiler chick's diet substantially decreased serum cholesterol, triglyceride and increased the glucose, total protein level and improved the LDL/HDL ratio [13].

The results of these studies already knew that Citronella oil is able to inhibit the growth of bacteria, thus the livestock will easily digest their food and thus the growth and consumption of feed in livestock will increase periodically. Undoubtedly, based on the ability of Citronella oil as an antimicrobial and able to increase growth in poultry, it is necessary to have study the influence of Citronella to reduce egg cholesterol and egg fat of quail.

## 2. MATERIALS AND METHODS

### 2.1. Quails

As much as one hundred female quails (*Coturnix coturnix japonica*) (age 2 weeks) were used in the study. Quails were fed with four treatments in completely randomized design (CRD). Quail was housed in wire cages with dimensions of 35x25x25 cm (width, length, height). The experiment diets and water were given ad libitum for all groups. Quails were randomized to four diet treatments. Each treatment had five replicated of five female quails (*Coturnix coturnix japonica*).

### 2.2. Research Design

In four treatments, one was given 0% Citronella oil per kg of feed (T1) as the control diet, and the other diets were supplemented with the level of Citronella oil, namely, 0.3% Citronella oil per kg of feed (T2) per kg of feed, 0.6% Citronella oil per kg of feed (T3), and 0.9% Citronella oil per kg of feed (T4). The parameter measured were egg protein, cholesterol and egg fat content. The data obtained were analyzed for variance (ANOVA) and Duncan's Multiple Range Test (DMRT).

### 2.3. Collection of Egg

When the quails have been producing for four weeks, the eggs were then collected and analyzed for protein, fat and cholesterol content. Each cage unit was taken 5 eggs, so the number of eggs observed was 100 eggs. The method used to analyze egg cholesterol was the Liebermann Burchard method [14], while the method for protein analysis uses the Kjeldhal method (Rosaini et al., 2015) and the method for fat analysis uses the Soxhlet method [15].

### 2.4. Diet Experiment

The study used commercially diet available in packs produced by New Hope Indonesia. The ingredient composition and nutrient content of the diets for

experimental are presented in Table 1. The citronella oil extract used is available in the market in packaging produced by Agroatsiri.

**Table 1.** Nutrient content of the diets

Nutrients	Percentages (%)
Water content (max)	13.0
Protein	22 – 24
Fat (min)	5.0
Crude Fiber (max)	5.0
Ash (max)	7.0
Calcium (min)	0.9
Phosphor (min)	0.6
Metabolic energy (kcal/kg)	2900

References: PT. New Hope Indonesia

## 3. RESULTS AND DISCUSSION

The average mean of egg protein, egg cholesterol and egg fat content of female quail (*Coturnix coturnix japonica*) as influenced of dietary inclusion of citronella oil are presented in Table 2.

The statistical analysis on egg protein parameters had no significant difference ( $P > 0.05$ ) among the dietary treatment. Shown in Table 2 that the average the levels of egg protein in T1 as control ( $11.22 \pm 0.93$  mg/dl) was lowest as compared to the other treatments i.e. T2 ( $12.37 \pm 1.11$  mg/dl), T3 ( $11.34 \pm 1.17$  mg/dl) and T4 ( $11.71 \pm 1.07$  mg/dl), respectively. It is indicated that the optimum beneficial effect of citronella oil supplementation as an herbal feed additive up to 0.9% per kg of feed of can be increasing the level of egg protein.

The trend of average the levels of egg cholesterol of quail is given in Table 2. The egg cholesterol was lowest in T3 ( $362.19 \pm 11.19$  mg/dl), followed by T4 ( $363.68 \pm 5.21$  mg/dl), T2 ( $378.53 \pm 14.89$  mg/dl) and highest value in T1 as control group ( $382.15 \pm 9.87$  mg/dl).

Analysis of variance showed that the treatment of citronella oil supplementation in the feed had significantly effect ( $P < 0.05$ ) on egg cholesterol. Duncan's Multiple Range Test (DMRT) indicated that T3 and T4 significantly different ( $P < 0.05$ ) to T1, it is showed that the beneficial effect of citronella oil supplementation as an herbal feed additive as much 0.6% Citronella oil per kg of feed of can be reduce the level of egg cholesterol.

Similar observations by [16] who report that combination of probiotics with ginger were inoculated in laying quail diet can decrease total cholesterol levels in serum and yolk (107.05 mg/dl and 10.6 mg/g) respectively compared to control group (158 mg/dl and 14.1 mg/dl). Parade et al. (2019) that the cholesterol and high density lipoprotein (HDL) serum of the broiler

chickens was quantitatively lowest in experimental chickens receiving lemongrass leaf meal 1.5 per cent level as compared to control group without lemongrass leaf meal. Besides that low density lipoprotein (LDL) serum of the broiler was quantitatively highest in chickens receiving lemongrass leaf meal 1.5 per cent level as compared to control group without lemongrass leaf meal. The other study by [17] also reported that quail egg cholesterol decreased significantly in the 400 ppm niacin treatment, and continued to decrease the greater

0.5% and 1 % supplemented with ginger root powder (66.33 mg/dl and 58.33 mg/dl) compared to control group (94.33 mg/dl).

That study is contradictory with the finding of [18] who reported that the use of turmeric powder with a level of 54 mg/bird/day does not affect feed consumption and egg cholesterol levels quail produced at the age of 4 months, but the use of turmeric powder with a level of 54 mg/bird/day when became sexual maturity has an impact on increasing levels of quail egg cholesterol. As well the study by [19] report that serum cholesterol levels were not affected by the fenugreek seed powder as phytochemical feed additives in broiler diet.

Another study found that *Cymbopogon citratus* leaf meal has impact on poultry body weight. [13] that *Cymbopogon citratus* leaf meal in poultry diets resulted in significantly ( $P<0.05$ ) higher body weight compared to

**Table 2.** The egg protein, cholesterol and egg fat content of quail (Cortunix cortunix japonica)

Treatment	Egg protein (mg/dl) <sup>ns</sup>	Egg Cholesterol (mg/dl)	Egg fat (mg/dl)
T1	11.22±0.93	382.15± 9.87 <sup>a</sup>	23.95±0.41 <sup>c</sup>
T2	12.37±1.11	378.53±14.89	19.53±0.42 <sup>d</sup>
T3	11.34±1.17	362.19±11.19 <sup>b</sup>	17.42±0.81 <sup>e</sup>
T4	11.71±1.07	363.68± 5.21 <sup>b</sup>	17.40±0.44 <sup>e</sup>

<sup>ns</sup> non significant; <sup>ab</sup> significant different ( $P<0.05$ ); <sup>(cd)(ce)(de)</sup> very significant different ( $P<0.01$ )

control diet at the end of 6th week. In contradictory to the finding of [20] who report that the quail (unsex) diet supplemented with 3% Lemmon grass leaf meal showed a significantly lower final body weight compared to the control ( $p<0.05$ ). This was similar with [21] who reported that reduced body weight in broilers fed with 2% Lemmon grass leaf supplemented. The other study by [22] report that the lemongrass oil did not affect significantly on body weight gain during experiment period from initial to end of the experiment (36 days).

Our experiment showed that egg fat of quail treated with various levels of citronella oil quantitatively more lower compared to the control group as showed in Table 2. The egg fat was lowest in T4 (17.40±0.44 mg/dl), followed by T3 (17.42±0.81 mg/dl), T2 (19.53±0.42 mg/dl) and highest value in T1 as control group (23.95±0.41 mg/dl).

the niacin level. The further explanation is that the addition of niacin in the feed significantly reduce HDL and increase LDL. The function of LDL is to send cholesterol to the coronary vessel tissue which can cause calcium accumulation in the coronary blood vessels, while function HDL is absorbing cholesterol deposits in the tissue and then send it to the liver and then break down into bile acids. The study by [23] that serum cholesterol of unsexed broiler were significantly lower ( $P<0.05$ ) in

Analysis of variance showed that the treatment of citronella oil supplementation in the feed had very significant effect ( $P<0.01$ ) on egg fat. Duncan's Multiple Range Test (DMRT) indicated that T1 very significantly different ( $P<0.01$ ) to T2, T3 and T4. Other wise T2 significantly different ( $P<0.01$ ) to T3 and T4, but T3 had non different to T4. So this trend showed that the beneficial effect of citronella oil supplementation as an herbal feed additive as much 0.3% per kg of feed of can be reduce the level of egg fat.

These findings were in agreed with those of [16] found that the use of niacin as much as 1000 ppm in feed was able to reduce triglyceride serum levels from 155.52 become 100.19 (mg/dl) in female quails. These results contradictory with the finding by [12] who report that triglyceride serum of broiler with supplemented 1.5% level of lemongrass leaf meal (147.50 mg/dl) was

significantly different ( $P<0.05$ ) than other three groups i.e. supplemented 1.0% (140.25 mg/dl), supplemented 0.5% (136.75 mg/dl) and control group (130.50 mg/dl). Similar the study by Zomrawi et al. (2012) that serum triglyceride of unsexed broiler were significantly lower ( $P<0.05$ ) for control group (44 mg/dl) and 0.5% supplemented with ginger root powder (25.67 mg/dl) compared to other treatments i.e. 1.5% supplemented (99.33 mg/dl) and 1.0% supplemented (62.33 mg/dl)

#### 4. CONCLUSION

From this research it can be concluded that giving citronella oil as much as 0.6% per kg of feed can reduce fat and cholesterol in quail eggs.



## REFERENCES

- [1] Primacitra, Y.D., O. Sjojfan, dan M.H. Natsir. 2014. Pengaruh Penambahan Probiotik (*Lactobacillus* Sp.) dalam Pakan terhadap Energi Metabolis, Kecernaan Protein dan Aktivitas Enzim Burung Puyuh. *J. Ternak Tropika*, vol. 15, no. 1, pp. 74-79
- [2] Behura, S., dan V.K. Srivastav. 2004. Essential oil of leave of curcuma species. *Journal of Essential Oil Research*. 16(2): 109-110
- [3] Fawati, N.M., I.W. Suirta, dan S. Bahri. 2016. Isolasi, Identifikasi, Serta Uji Aktifitas Antibakteri Pada Minyak Atsiri Sereh Wangi (*Cymbopogon winterianus* Jowitt). *Jurnal Kimia*. 10 (2) : 219-227.
- [4] Luangnarumitchai S., Lamlerththong, S., and Tiyafoonchai, W. 2007. Antimicrobial activity of essential oils against five strains of *Propionibacterium acnes*. *Mahidol University Journal of Pharmaceutical Sciences*. 34: 60-64.
- [5] Ibrahim, Hayam M., Ferial M. Abu Salem. (2013). Effect of Adding Lemongrass and Lime Peel Extracts on Chicken Patties Quality. *Journal of Applied Sciences Research*. 9(8). 5036.
- [6] Paramartha D.A. 2015. Kajian Aktivitas Antibakteri Minyak Daun Sereh (*Cymbopogon citratus*) pada Adonan Sate Lilit Ikan Laut. *Jurnal Media Ilmiah Teknologi Pangan (Scientific Journal of Food Technology)*. 2 (1). 029-040.
- [7] Calsamiglia S., M. Busquet, P.W Cardozo, L. Catillejos, and A. Ferret. 2007. Invited Review: essential oils as modifiers of rumen microbial fermentation. *J Dairy Sci*. 90:2580- 2595
- [8] Sjojfan, O., D.N. Adli, M.H. Natsir, dan A. Kusumaningtyaswati. 2020. Pengaruh Kombinasi Tepung Kunyit (*Curcuma domestica* VAL.) dan Probiotik terhadap Penampilan Usus Ayam Pedaging. *Jurnal Nutrisi Ternak Tropis dan Ilmu Pakan*. 2(2) : 19-24.
- [9] Bozkurt M., N. Aysul, K. Küçükyılmaz, S. Aypak, G. Ege, A. U. Çatli, H. Akşit, F. Çöven, K. Seyrek, and M. Çınar. 2014. Efficacy of in-feed preparations of an anticoccidial, multienzyme, prebiotic, probiotic, and herbal essential oil mixture in healthy and *Eimeria* spp.-infected broilers. *Poultry Science* 93 :389–399
- [10] Lee K. W., Everts, H. and Beyne, A. C. 2004. Essential oils in broiler nutrition. *International Journal of Poultry Science*. 3 (12): 738 – 752
- [11] Rumokoy L., Charles Kaunang, and W. Toar. 2017. Efek ekstrak minyak citronela *Cymbopogon nardus* L. terhadap proporsi frekuensi kontak fisik *Musca domestica* L. (Diptera: Muscidae) pada ransum dan performa broiler. *Indonesian Journal of Entomology*. 14 (2): 89–96.
- [12] Tiwari M. R., Jha P. K., Sah B., Kunwar G. and Jha A. K. 2018. Performance of Lemongrass (*Cymbopogon citratus*) Oil as Growth Promoter in Broiler. *Bang. J. Anim. Sci.*, 47(2): 85-91.
- [13] Parade A. K., B. M. Thombre, R. A. Patil, P. V. Padghan, S. S. Shinde and P. B. Meshram. 2019. Inclusion of Lemongrass (*Cymbopogon citratus*) Leaf Meal as a Natural Feed Additive for Evaluation Serum Lipid Profile of Broilers. *Int. J. Curr. Microbiol. App. Sci*. 8(10): 1835-1841.
- [14] Adu, J.K., Cedric D. K. A, Naomi K., Emmanuel O., Stella A.G.P., Bernice K.O. 2019. Validation of a Simple and Robust Liebermann–Burchard Colorimetric Method for the Assay of Cholesterol in Selected Milk Products in Ghana. *Int J. Food Science*, 2019, Article ID 9045938, 7 pages. <https://doi.org/10.1155/2019/9045938>
- [15] Aminullah, Mardiah, Muhammad Reza Riandi, Arum Puspito Argani, Gustini Syahbirin, and Tetty Kemala. 2018. The Effect of Types of Fat Extraction Methods on Total Lipids of Chicken and Pork Fat. *Journal of Halal Agroindustry* 4 (1): 094– 100
- [16] Khalifa M I and Noseer E A. 2019 Cholesterol quality of edible eggs produced by quail fed diets containing probiotic and/or ginger (*Zingiber officinale*). *Livestock Research for Rural Development*. Volume 31, Article #165. Retrieved August 1, 2021, from <http://www.lrrd.org/lrrd31/10/mrwa31165.html>
- [17] Aviati, V., S.M. Mardiaty and T.R. Saraswati. 2014. Kadar Kolesterol Telur Puyuh Setelah Pemberian Tepung Kunyit Dalam Pakan. *Buletin Anatomi dan Fisiologi*. 22(1): 58-64
- [18] Fenita Y and T Suteky. 2006. Pengaruh Pemberian Niacin terhadap Kadar Kolesterol Telur dan Perlemakan Serum Darah Puyuh (*Cortunix cortunix japonica*). *Jurnal Sain Peternakan Indonesia*. 1(2): 45-48.
- [19] Weerasakha A. S. and N.S.B.M. Atapattu. 2013. Effect of Fenugreek (*Trigonella foenumgraecum* L.) Seed powder on Growth Performance,

- Visceral Organ Weight, Serum Cholesterol Levels and the Nitrogen Retention of Broiler Chicken. *Tropical Agricultural Reserch*. 24(3):289-295.
- [20] Sariözkan S, Y Konca, GüçlüBk, E 10 tuğ, M Kaliber, Ş Özkan and ST Ay (2016). *The effect of dietary supplementation of lemongrass (Cymbopogoncitratu)s on performance, carcass*
- [21] Thayalini, K., Shanmugavelu, S., Saminathan, P.M., Siti Masidayu, M.S., Nor Idayusni, Y., Zainuddin, H., Nurul Akmal, C.A. and Wong, H.K. 2016. Effects of Cymbopogon citratus leaf and Zingiber officinale rhizome supplementation on growth performance, ileal morphology and lactic acid concentration in broilers. *Mal. J. Anim. Sci.* 14:43-49
- [22] Tiwari M. R., Jha P. K., Sah B., Kunwar G. and Jha A. K. 2018. Performance of Lemongrass (Cymbopogon citrates) Oil as Growth Promoter in Broiler. *Bang. J. Anim. Sci.*, 47(2): 85-91.
- [23] Zomrawi W. B., KH. A. Abdel Atti, B. M. Dousa and A. G. Mahala. 2012. The Effect of Ginger Root Powder (Zingiber Officinale)Supplementation on Broiler ChicksPerformance, Blood and Serum Constituents. *Online Journal Anim. and Feed Reserch*. Vol. 1(6): 457-

# 14.\_Article\_of\_effort\_to\_Reduce\_fat\_and\_cholsterol.pdf

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

3%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	repository.ub.ac.id Internet Source	2%
2	www.ijcmas.com Internet Source	2%
3	download.atlantis-press.com Internet Source	1%
4	altifani.org Internet Source	1%
5	Submitted to University of the Philippines Diliman Student Paper	1%
6	uad.portalgaruda.org Internet Source	1%
7	www.chemijournal.com Internet Source	1%
8	conference.khuisf.ac.ir Internet Source	1%
9	Luana Roland Ferreira Contini, Thais de Souza Zerlotini, Isadora Fernandes Brazolin, Jackson	1%

Wesley Silva dos Santos et al. "Antioxidant chitosan film containing lemongrass essential oil as active packaging for chicken patties", Journal of Food Processing and Preservation, 2021

Publication

10

G. Zanaty, Eman Hussein. "EFFECT OF ADDING DIFFERENT SOURCES OF PROBIOTICS TO DIETS CONTAINING OXIDIZED PALM OIL ON SOME PRODUCTIVE AND REPRODUCTIVE TRAITS IN GIMMIZAH LAYING HENS", Egyptian Journal of Nutrition and Feeds, 2021

Publication

<1 %

11

M A Yusuf, A L Toleng, M F Syafar. "Distribution of Cows by Days in Milk (DIM) at First AI and Calving to Conception Interval in Dairy Cows", Media Peternakan, 2012

Publication

<1 %

12

Yosi Fenita. "Pengaruh Pemberian Niasin Terhadap Kadar Kolesterol Telur dan Perlemakan Serum Darah Puyuh (Coturnix coturnix japonica)", Jurnal Sain Peternakan Indonesia, 2006

Publication

<1 %

13

[ejurnal.bunghatta.ac.id](http://ejurnal.bunghatta.ac.id)

Internet Source

<1 %

14

[pub.epsilon.slu.se](http://pub.epsilon.slu.se)

Internet Source

<1 %



15

gjournals.org

Internet Source

<1 %

16

sipora.polije.ac.id

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off