

jurnal Eggs effectiveness

by Turnitin Checker

Submission date: 01-May-2023 01:56AM (UTC+1000)

Submission ID: 2079837361

File name: jurnal_Eggs_effectiveness.pdf (279.12K)

Word count: 4563

Character count: 24321

EGGS EFFECTIVENESS LEAVES EXTRACT *PIPER BETLE* TO ATTACK PREVENTION FUNGUS *SAPROLIGNIA SP* IN *OSTEOCHILUS HASSELTII*

^{1*}Kasprijo, ²Nuni Aisah Dan and ³Rudi Wijaya

^{1,3}Teaching staff, Faculty of Fisheries and Marine Sciences, Jenderal Soedirman University,
Indonesia.

²Students, Faculty of Fisheries and Marine Sciences, Jenderal Soedirman University,
Indonesia.

Article Received on
21 June 2021,
Revised on 12 July 2021,
Accepted on 02 August 2021
DOI: 10.20959/wjpr202110-21072

*Corresponding Author

Kasprijo

Teaching staff, Faculty of
Fisheries and Marine
Sciences, Jenderal
Soedirman University.
Indonesia.

ABSTRACT

Betle leaf (*Piper betle*) is one of the traditional native plant to Indonesia that known to contain antifungal substances. The beetle leaves contained essential oils, tannins, phenoids, saponins that able to inhibit the growth of fungi. This study aims to determine the optimal extract concentration of beetle leaf on egg hatchability and survival of nilem larvae. This study used an experimental method with a completely randomized design consisting of 5 preparations and 3 replications. The treatment given is soaking nilem fish eggs for 20 minutes into the media with beetle leaf extract contents namely P1: 0 ppm (control), P2: 20 ppm, P3: 40 ppm, P4: 60 ppm, and P5: 80 ppm. Test parameters that observed including egg hatchability, larval

survival, prevalence of fungal infection and water quality. The results of egg immersion in beetle leaf extract at a concentration of 80 ppm (P4) showed the best results with 95 ± 3.36% hatching rate off eggs. The prevalence of fungal infection on P5 was 3.67 ± 2.46%, and the larval spreading on P4 was 87.34 ± 3.17%. Water quality in the study showed a pH value of 7, temperatures between 21 - 23°C, and dissolved oxygen of 6.06 - 7 mg / L.

KEYWORDS: *Osteochilus hasseltii* egg, Beetle Leaf Extract, prevalence fungus Hatching Rate, Intensity of Fungal Infection.

INTRODUCTION

Nilem fish is an endemic fish of Indonesia that normally lives in fresh water with clear water quality and is one of the freshwater aquaculture commodities concentrated on the island of Java. Nilem fish (*O. hasselti*) is a freshwater fish that is not only popular with meat but the eggs are also consumed by the people. Nilem fish eggs have a delicious flavor and can be explored in certain countries. High public interest in Nilem makes Nilem fish production continue to increase (Firdaus *et al.*, 2017). This causes the Nilem fish must be cultivated to meet the demand so that fish stocks are maintained.

One obstacle in the cultivation of Nilem is the availability of seeds that are not continuous. The availability of seeds is determined by the number of eggs that hatch.

The hatching of eggs is the process of developing an embryo in the egg until it hatches (Sa'diah *et al.*, 2015). At the egg hatching stage, Nilem fish often experience problems, one of which is the egg does not hatch. Eggs that do not hatch are usually caused by fungus. Eggs that are attacked by fungi can cause eggs to not hatch or rot so that the hatching rate is low (Dian *et al.*, 2015). Overcoming this, prevention is needed.

One of the efforts made in preventing eggs attacked by fungi is usually by adding chemicals whose purpose is to inhibit the growth of fungi in eggs. Chemicals that are frequently used include Methylene blue, formalin, melachine green, NaCl, KMnO_4 , (Rivanto *et al.*, 2014). The use of synthetic chemicals including antibiotics can have a resistant effect on pathogenic microbes and cause environmental pollution if their use is inappropriate (Sabrina *et al.*, 2014).

Overcoming this problem, it is necessary to have an alternative prevention by using materials that are environmentally friendly and do not cause effects resistant to fungi and certainly can prevent fungal diseases (Rosidah *et al.*, 2017). One alternative is to use traditional plants that are anti-fungal. In addition to being antifungal, these plants are also easily obtained and are easy to use for preventing and treating fish or egg disease (Zuraidah and Silkhairi, 2016). One of the traditional plants that can potentially prevent fungal diseases is the betel leaf plant (*Piper betle* L) (Rivanto *et al.*, 2014).

5

MATERIALS AND METHODS

3.1.3. Material

The materials used for making the extract were fresh betel leaf, distilled water and filter paper. Other materials used for broodstock injection of nilem fish include ovaprim and aquadest.

3.1. Research methods

The research method used in this study is an experimental method using a completely randomized design (CRD) with 5 treatments and 3 replications so that it becomes 15 experimental units. The treatment given was soaking nilem fish eggs into media with different content of betel leaf extract, namely:

P1: Betel leaf extract concentration 0 ppm (control)

P2: Betel leaf extract concentration 20 ppm

P3: Betel leaf extract concentration 40 ppm

P4: Betel leaf extract concentration 60 ppm

P5: Concentration of betel leaf extract 80 ppm

The research variables observed were egg hatchability, fungal attack rate, and larval survival. The main data calculated were the number of eggs that hatched and the survival of larvae that were reared until the yolks were exhausted. Supporting data in the form of the number of eggs attacked by fungi and water quality include temperature, pH, and dissolved oxygen.

3.2. RESEARCH PROCEDURE

3.3.1. Production of betel leaf extract (*Piper betle* L)

Making betel leaf extract is done by selecting fresh betel leaves, then washed and drained without the help of sunlight. Then the betel leaves were dried using an oven at 60°C for 1 day. Then the dried betel leaves are mashed using a blender which is then sieved with a sieve to obtain powder. Then the powder was dissolved using 1 liter of distilled water with the amount of simplicia according to the treatment (20 ppm, 40 ppm, 60 ppm, and 80 ppm). The powder was brewed with distilled water at 90°C for 30 minutes. Betel leaf extract can be used after cooling (Dwiyanti, 1996 in Zuraidah and Silkhairi, 2016).

3.3.2. Container Preparation

Container preparation includes soaking and maintenance containers. The container is a basin measuring 6 liters of water as many as 15 pieces. Previously, the basin was washed and dried,

then filled with water and given a solution of Potassium Permanganate for 24 hours. After that, it was rinsed and dried, then filled with 2 liters of water again with a water level of 4.8 cm and aerated for 24 hours.

3.3.3. Nilem Fish Spawning

Nilem fish spawning is done artificially. There are 2 brooders (1 female and 1 male) and the gonads are ripe and ready to be spawned. Before spawning, the female and male Nile broodstock were weighed to determine the dose of ovaprim. The female Nile broodstock were injected with 0.5 ml/kg ovaprim and the male parent was injected with 0.3 ml/kg ovaprim. After being injected, the broodstock is put into a fiber bath for the spawning process. After spawning, Nile fish eggs are ready to use.

3.3.4. Giving Betel Leaf Extract (Piper betle L)

The betel leaf extract was given using the soaking method for 20 minutes. Soaking was carried out in a 6 l basin with a volume of 1 l of betel leaf extract. 100 eggs were taken using a sieve, then soaked in betel leaf extract according to treatment for 20 minutes. After soaking, the eggs are transferred to a pre-prepared rearing basin. The immersion method refers to the research of Ghofur et al., (2014).

3.3.5. Maintenance of Nilem Fish Eggs

Eggs that have been transferred to the rearing basin are maintained until the eggs hatch. During the hatching process, each rearing container is aerated. After the eggs hatch, the eggs that hatch are counted and observed the eggs that are attacked by fungus by looking at the number of dead fish eggs with fungus and dead eggs without fungus. Eggs that are attacked by fungus are characterized by the presence of threads that cover the eggs like cotton. Then after hatching, the larvae are reared for as long as the egg yolk runs out. Then calculated to determine the survival value of Nile fish larvae.

RESULTS AND DISCUSSION

Research on the effect of betel leaf extract (Piper betle L) on egg hatchability and larval survival of Nile fish (*Osteochilus hasselti*) with 5 different treatments. The treatments were P1 (control), P2 (20 ppm betel leaf extract), P3 (40 ppm betel leaf extract), P4 (60 ppm betel leaf extract), P5 (80 ppm betel leaf extract).

4.1. Egg Hatchability

Observation of egg hatchability was carried out after $\pm 24 - 48$ hours by counting the number of eggs that hatched into larvae (Marthin *et al.*, 2018). Egg hatchability was obtained by observing fertilized eggs and hatched into larvae. The results of observations of the percentage of egg hatchability by immersion treatment on betel leaf extract with different concentrations can be seen in Figure 1.




Figure 1: Hatchability of Nilem Fish Eggs.

Note: The same superscript letter indicates that the betel leaf extract with different concentrations has a significantly different effect ($P < 0.05$).

The results showed an increase in egg hatchability at P1 (88.67 ± 2.26), P2 (93.34 ± 1.29), P3 (94.67 ± 0.72), P4 (95 ± 3.36), and P5 (94 ± 1.21). The results of statistical analysis (ANOVA) showed that the administration of betel leaf extract with different concentrations in each treatment showed a significant difference ($P < 0.05$). The statistical test was continued with the Tukey Test. Tukey's test results showed that P1 was not different from P2 but significantly different from P3, P4, and P5. However, at P2, P3, P4, and P5 were not significantly different. The hatchability of eggs at P4 was the best to prevent the growth of fungus on the eggs. P4 with a value of $95 \pm 3.36\%$ showed the best results compared to P2, P3, and P5, presumably because the betel leaf extract contains various anti-bacterial or fungal compounds and the concentrations used are effective so that they are not harmful to egg development. Munurut Huzni *et al.*, (2015), active compounds found in betel leaf such as alkaloids, flavonoids, and saponins are able to protect the chorion from fungal attacks.

In P3 the hatchability value of eggs was lower than in P4, this was presumably because the concentration of the extract given did not have an effect on resisting fungal attacks so that the eggs would also rely on chorion to resist fungal attacks. According to Sumantadinata (2012), if the embryonic development process is attacked by fungi, the ability of the eggs to hatch will be reduced and even cause the death of the eggs so that the success of the eggs is low.

At P5 with a concentration of 80 ppm it was lower than P4 with a value of $94 \pm 1.21\%$, this was caused by active compounds such as saponins which were able to poison organisms at certain concentrations. Inaya *et al.*, (2015) confirmed that saponin compounds can inhibit egg development by damaging cells so that there is a change in the structure of the egg which causes fluid in the cell to come out and cell dehydration occurs. Cell dehydration that occurs will cause the eggs to fail to hatch, because in their development the eggs require fluids that contain nutrients.

4.2. The prevalence of fungal infection rate

Fish eggs are known to be relatively susceptible to fungal attack. The fungus will attack infertile (dead) eggs and attack healthy eggs. How to calculate the number of eggs infected with fungus according to Helmiati (2005) is the number of eggs infected with fungi divided by the number of initial eggs. Percentage of fungal attack rate obtained during the study:




Figure 2: The prevalence of fungal infection rate.

Note: The same superscript letter indicates that the betel leaf extract with different concentrations has a significantly different effect ($P < 0.05$).

Based on the results of the research that has been done, the more concentration given, the less the level of fungal attack. The results of the ANOVA test showed a significant difference ($P < 0.05$) to the treatment. Tukey's test results show that P1 is not different from P2 and P3, but different from P4 and P5. At P1, P2, and P3 are not different, but different from P4 and P5. The results of the Tukey test showed that the lowest level of fungal attack was found in P5 with a value of $3.67 \pm 2.46\%$, it is suspected that the concentration given to P5 was more capable of inhibiting fungal growth. Rivanto *et al.*, (2014), added that the higher the concentration of betel leaf extract to a certain extent will be able to kill the fungus but also the high concentration of the extract will be absorbed by the eggs so that it is toxic to the eggs and causes the eggs not to hatch.

Eggs that were not soaked with betel leaf extract in P1 (control) gave the fungus *Saprolegnia* sp. the highest is $9 \pm 2.01\%$. The high fungal attack is because the eggs are not soaked by betel leaf extract so they are not protected by antifungal substances contained in the extract, so the fungus will easily stick to the eggs, enter and infect the eggs. This causes uncontrolled fungal growth so that it continues to attack healthy eggs and results in healthy eggs that die and do not hatch. According to Effendie (2001), fungal attacks can weaken the chorion, then it becomes wrinkled because the fungus attaches and penetrates the chorion to take the food substances in it.

According to Fitri (2007), eggs are a good growing medium for microorganisms, because eggs contain chemical compounds that serve as a source of nutrition for these microorganisms. Without antifungal compounds, the resistance of eggs to attack by the fungus *Saprolegnia* sp. weakened and only rely on the strength of the chorion alone. Eggs attacked by *Saprolegnia* sp. Visible hyphae in the form of fine threads attached to the surface of the egg. The eggs that do not hatch due to the attack of the fungus *Saprolegnia* sp. appears to show signs around the egg there are fine threads like white cotton. This is in accordance with the opinion of Kordi (2004), who said that eggs that are attacked by fungi will usually appear covered with thread-like formations known as white fungal hyphae.

Unhatched eggs are thought to be the influence of low water quality Eggs that died were not attacked by fungi did not show the presence of hyphae of *Saprolegnia* sp. on the surface of the fish eggs. According to Rosidah *et al.*, (2017), the greater the concentration given, the higher the average percentage of dead eggs.

4.4. Water quality

The results of water quality measurements during the study took place the temperature values in the morning ranged from 22 - 23 oC and in the afternoon ranged from 22 - 24 oC. From the results of temperature measurements during the study, it can be said that it is still in the normal range. Where according to Sunarma (2004), the normal temperature for hatching eggs is 22-32 C. While *Saprolegnia* sp. can grow at a temperature of 0-35 C with the best growth temperature in the range of 15-30 C and pH 4-6 (Irianto, 2005). According to Rozaldi, et al., (1990), stating that the speed of mushroom growth is closely related to environmental temperature, in general *Saprolegnia* mushrooms can grow at a minimum temperature of 0-5 C and an optimum of 15-30 C. So eggs that do not hatch are not caused by poor water quality but by *Saprolegnia* sp. or treatment.

The results of measuring the pH of the water during the study were 7. Based on the PP No. 82 of 2001 (class II), a good pH for aquaculture activities is in the range of 6-9. According to Amalia and Subanditono, (2013) that acidity (pH) that is not optimal can cause fish stress, disease susceptibility, productivity, and low growth. A low pH can cause a decrease in growth rate, and a high pH will increase ammonia which is indirectly harmful (Paulinus and Revol, 2015). So from the results of the pH measurements above, it shows that the pH during the study was within the feasibility range. Dissolved oxygen levels during the hatching process averaged 6.44 mg/L, while at the time of rearing Nile fish larvae, dissolved oxygen averaged 6.73 mg/L. Thus, the dissolved oxygen level during the study was still within reasonable limits for hatching Nile fish eggs. Sumarna (2004) confirmed that a good range of dissolved oxygen is > 3 mg/L. From the results of this study, the dissolved oxygen content is still within the tolerance limit. The provision of dissolved oxygen in the hatching media water is needed as an oxygen supply through the addition of aeration. The oxygen enters the egg by diffusion through the surface layer of the egg shell. If oxygen levels are low, it will affect biological functions and can even lead to death (Sutisna and Sutarmanto, 1995).

CONCLUSIONS

Based on the results of the study, the following conclusions can be drawn:

1. Soaking eggs with different concentrations of betel leaf extract has an effect on egg hatchability and survival of Nile fish larvae.
2. The best concentration was at P4 with egg hatchability value of $95 \pm 3.36\%$.

REFERENCES

1. Adilfiet. 1994. *Buku Ajar Mikrobiologi Kedokteran*. Binarupa Aksara. Jakarta.
2. Andika, H., E.Dewantoro, E.I. Raharjo. Perendaman Telur Ikan Gurame (*Osphronemus gouramy*) Dengan Ekstrak Meniran (*Phyllanthus niruri* L) Sebagai Anti Jamur. *Jurnal Ruaya*, 2014; 1(1): 71-76.
3. Aqil, I. D. Pemanfaatan Plankton Sebagai Sumber Makanan Ikan Bandeng (*Chanoschanos*) di Waduk Ir. H. Juanda, Jawa Barat. *Skripsi*. Universitas Islam Negeri Syarif Hidayatullah. Jakarta, 2010.
4. Arfah, H., L. Muftucha dan O. Carman. Pemijahan Secara Buatan Pada Ikan Gurame (*Osphronemus gouramy* Lac.) Dengan Penyuntikan Ovaprim Induced Spawning Of Giant Gouramy *Osphronemus Gouramy* Lac. By Ovaprim. *Jurnal Akuakultur Indonesia*, 2006; 5(2): 103-112.
5. Atmawati, S.N. *Perbedaan Keanekaragaman Zooplankton di Daerah Sekitar Keramba dan Sekitar Warung Apung Rawa Jombor Hubungannya dengan Kualitas Perairan*. Universitas Negeri Yogyakarta: Yogyakarta, 2012.
6. Augusta, T.S. Dinamika Perubahan Kualitas Air Terhadap Pertumbuhan Pertumbuhan Ikan Lele Dumbo (*Clarias gariepinus*) yang Dipelihara di Kolam Tanah. *Jurnal Ilmu Hewan Tropika*, 2016; 5(1): 41-44.
7. Ayer, Y., J. Mudeng, H. Sinjal. Daya Tetas Telur dan Sintasan Larva Dari Penambahan Madu Pada Bahan Pengencer Sperma Ikan Nila (*Oreochromis niloticus*). *Jurnal Budidaya Perairan*, 2015; 3(4): 149-153.
8. Bachtiar, Y. *Mencemerlangkan Warna Koi*. Agromedia Pustaka. Jakarta, 2012.
9. Dian, D. A., Hastiadi, H., dan Eka, I. R. Pengaruh Ekstrak Daun Kemangi (*Ocimum basilicum* L) Terhadap Daya Tetas Telur Ikan Lele Dumbo (*Clarias gariepinus*) yang Diinfeksi Jamur *Saprolegnia* sp. *Skripsi*. Fakultas Perikanan Ilmu Kelautan Universitas Muhammadiyah Pontianak. Pontianak, 2015.
10. Diansyah, S., dan F. Diana. Pengaruh Ekstrak *Tagetes erecta* L Terhadap Pertumbuhan Jamur *Saprolegnia* sp. Secara Invito. *Jurnal Perikanan Tropis*, 2017; 4(2): 199-208.
11. Effendie, M. I. *Biologi Perikanan*. Yayasan Nusatama. Bogor, 2001.
12. Fazriel, M., C. Yulvizar., I. Hasri. Pengaruh Suplemen Dan Probiotik Pada Pakan Terhadap Pertumbuhan dan Kelangsungan Hidup Larva Ikan Peres (*Osteochilus vittatus*). *Jurnal Ilmiah Mahasiswa Kelautan dan Perikanan Unsyiah*, 2017; 2(1): 158-168.
13. Firdaus, N., Iskandar, H. Hamdani. Pengaruh *Lemna* sp. Sebagai Pakan Dalam Budidaya Ikan Nilem Organik. *Jurnal Perikanan dan Kelautan*, 2017; 8(1): 9-13.

14. Firmantin, I. T., A. Sudaryono, R. A. Nugroho. Pengaruh Kombinasi Omega-3 dan Klorofil Dalam Pakan Terhadap Fekunditas, Derajat Penetasan dan Kelulushidupan Benih Ikan Mas (*Cyprinus carpio* L). *Journal of Aquaculture Management and Technology*, 2015; 4(1): 19-25.
15. Fitri, A. Pengaruh Penambahan Daun Salam (*Eugenia polyantha* W) Terhadap Kualitas Mikrobiologis, Kualitas Organoleptis dan Daya Simpan Telur Asin Pada Suhu Kamar. *Jurnal Pangan dan Agroindustri*, 2007; 2(5): 6-28.
16. Fitria, A.S. Analisis Kelulushidupan dan Pertumbuhan Benih Ikan Nila Larasati (*Oreochromis niloticus*) F5 D30-D70 Pada Berbagai Salinitas. *Skripsi*. Fakultas Perikanan dan Ilmu Kelautan, Universitas Diponegoro. Semarang, 2012.
17. Ghofur, M., M. Sugihartono, R. Thomas. Efektifitas Pemberian Ekstrak Daun Sirih (*Piper betle* L) Terhadap Penetasan Telur Ikan Gurame (*Osphronemus gouramy* Lac). *Jurnal Ilmiah Universitas Batanghari Jambi*, 2014; 14(1): 37-44.
18. Hasan, H., Raharjo, E.I., Ariyani, D.D. Pengaruh Ekstrak Daun Kemangi (*Ocimum basilicum* L) Terhadap Daya Tetas Telur Ikan Lele Dumbo (*Clarias gariepinus*) yang Diinfeksi Jamur *Saprolegnia* sp. *Jurnal Ruaya*, 2016; 4(1): 18-23.
19. Herawati, Ratna. Ekstrak Daun Sirih (*Piper betle* L) Sebagai Insektisida Nabati Untuk Membasmi Larva Nyamuk *Aedes aegypti* L. *Skripsi*. Program Studi Biologi. Fakultas Teknobiologi. Universitas Atma Jaya Yogyakarta. Yogyakarta, 2010.
20. Hijriyati, K. H. 2012. Kualitas Telur dan Perkembangan Awal Larva Ikan Kerapu Bebek [*Cromileptes altevis*, Valenciennes (1998)] di Desa Air Saga, Tanjung Pandan, Belitung. *Tesis*. Program Studi Magister Ilmu Kelautan. Fakultas Matematika dan Ilmu Pengetahuan Alam. Universitas Indonesia. Jakarta.
21. Husni, M., G. Saptiani, Agustina. Pemberian Ekstrak Lengkuas (*Alpinia galanga*) Terhadap Daya Tetas Telur Ikan Lele Sangkuriang (*Clarias gariepinus*). *Jurnal Ilmu Perikanan Tropis*, 2016; 2(2): 80-84.
22. Huzni, M., B. T. Rahardho, H. Tarno. Uji Laboratorium Ekstrak Kiriuh (*Chomoloena adorata*: King dan Robinson) Sebagai Nematocida Nabati Terhadap *Meloidogyne* sp. (Chitwood). *Jurnal HPT*, 2015; 3(1): 93-100.
23. Inaya, A., F. N. Kismiyati., Subekti. Pengaruh Perasan Biji Pepaya (*Carica papaya*) Terhadap Kerusakan Telur *Argulus japonicus*. *Jurnal Ilmiah Perikanan dan Kelautan*, 2015; 7(2): 159-164.
24. Jangkaru, Zulkifli. 2001. *Pembesaran Ikan Air Tawar di Berbagai Lingkungan Pemeliharaan*. Penebar Swadaya. Jakarta.

25. Kamsuri, A. I., N. P. L. Pangemanan., A.T. Reiny. Kelayakan Lokasi Budidaya Ikan Di Danau Tondano Ditinjau dari Parameter Fisika Kimia Air. *Jurnal Budidaya Perairan*, 2013; 1(3): 31-42.
26. Kordi, K.M., Ghufuran. *Budidaya Perairan*. PT Citra Aditya Bakti. Bandung. Marthin, Lutfi., Juliaan Ch. Watung., O. J. Kalesaren., E. L. Ginting., H. J. Sinjai., I. R., 2004.
27. N. Salindeho. Penambahan Madu Dalam Pengenceran Sperma Terhadap Mortalitas Soermatozoa, Fertilisasi dan Daya Tetas Telur Ikan Patin Siam *Pangasius hypophthalmus*. *Budidaya Perairan*, 2018; 6(2): 45-52.
28. Mentari, N.L., Safrida, Khairil. Potensi Pemberian Ekstrak Daun Sirih (*Piper betle* L) Sebagai Pengawet Alami Ikan Selar (*Selaroides leptolepis*). *Jurnal Ilmiah Mahasiswa Pendidikan Biologi*, 2016; 1(1): 1-9.
29. Mufrod, Suwaldi, S. Wahyuono. Patch Ekstrak Daun Sirih (*Piper betle* L): Pengaruh Penambahan *Release Enhancer Substances* Terhadap Sifat Fisikokimia Dan Aktivitas Antibakteri. *Majalah Farmaseutik*, 2016; 12(2): 431-442.
30. Nugroho, A., e. Arini., T. Elfitasari. Pengaruh Kepadatan Yang Berbeda Terhadap Kelulushidupan dan Pertumbuhan Ikan Nila (*Oreochromis niloticus*) Pada Sistem Resirkulasi dengan Filter Arang. *Journal of Aquaculture and Technology*, 2013; 2(3): 94-100.
31. Putri, M. R. Anwar, Y. Sugianti, Krismono. Aspek Biologi Ikan Nilem (*Osteocholus vittatus*) di Danau Talaga. Sulawesi Tengah. *BAWAL*, 2015; 7(2): 111- 120.
32. Putri, S.K. *Budidaya Ikan Nilem (Osteochilus hasselti)*. Laporan praktikum Universitas Negeri Diponegoro. Semarang, 2010.
33. Raharjo, E.I., H. Hastiadi., Dermawan. Pergantian Pakan Terhadap Pertumbuhan dan Kelangsungan Hidup Larva Ikan Gurame (*Osphronemus gouramy*). *Jurnal Ruaya*, 2016; 4(1): 13-17.
34. Rivanto, I. Sidabalok, H. Hasan. Ekstrak Daun Sirih (*Piper betle* L) Untuk Pencegahan Infeksi Jamur *Saprolegnia* sp. Pada Telur Ikan Patin Siam (*Pangasius pangasius*). *Jurnal Ruaya*, 2014; 1(1): 17-22.
35. Rosidah, Y. Andriani, W. Lili, I. Herdiawan. Efektivitas Lama Perendaman Telur Ikan Lele Sangkuriang dalam Ekstrak Bunga Kecombrang untuk Mencegah Serangan Jamur *Saprolegnia* sp. *Jurnal Perikanan dan Kelautan*, 2017; 7(2): 199-209.
36. Sa'diah, I. N., Garnida, D., Mushawir, A. Mortalitas Embrio dan Daya Tetas Itik Lokal (*Anas* sp.) Berdasarkan Pola Pengaturan Temperatur Mesin Tetas. *Student e-Joernals*, 2015; 4(3): 1-12.

37. Saanin, H. 1984. *Taksonomi dan Kunci Identifikasi Ikan*. Bina Cipta. Jakarta.
38. Sabrina T. I., Sudarno, Suprpto. H. Uji Aktiivitas Antifungi Perasan Daun Kemangi (*Ocimum bacilicum*) Terhadap *Asperigillus terreus* Secara In Vitro. *Jurnal Ilmiah Perikanan dan Ilmu Kelautan*, 2014; 6(2): 7-12.
39. Samsundari, S., dan G.A. Wirawan. Analisis Penerapan Biofilter Dalam Sistem Resirkulasi Terhadap Mutu Kualitas Air Budidaya Ikan Sidat (*Anguila bicolor*). *JURNAL GAMMA*, 2013; 8(2): 86-97.
40. Soeminto. 2008. *Buku Dan Petunjuk Praktikum Struktur dan Perkembangan Hewan II*. Universitas Jenderal Soedirman. Purwokerto.
41. Srihati. 1997. *Pengaruh Suhu Terhadap Penetasan Telur, Pertumbuhan Dan Daya Tahan Hidup Larva Ikan Bandeng (Dicentrarchus labrax L.)*. Seminar biologiXV. Bandar Lampung. Lampung.
42. Sugianti, Budi. 2005. Pemanfaatan Tumbuhan Obat Tradisional Dalam Pengendalian Penyakit Ikan. *Makalah Pribadi Falsafah Sains (PPS-702)*. Sekolah Pasca Sarjana. Institut Pertanian Bogor. Bogor.
43. Sugihartono, Muhammad. Respon Tingkat Kepadatan Telur Ikan Gurami (*Osphronemus gouramy* Lac) Yang Berbeda Terhadap Daya Tetas Telur. *Jurnal Ilmiah Universitas Batanghari Jambi*, 2013; 13(4): 127-131.
44. Sumantadinata, K. 2010. *Pengembangan Ikan-ikan Peliharaan di Indonesia*. Sastra Hudaya. Surabaya.
45. Sunarma, A. 2004. *Peningkatan Produktifitas Usaha Lele Sangkuriang (Clarias sp.)*. Departemen Kelautan dan Perikanan. Direktorat Jenderal Perikanan Budidaya. Balai Budidaya Air Tawar Sukabumi. Sukabumi.
46. Syamsu, H.S.S., dan Hutapea, J.R. 1997. *Invetaris Tanaman Obat Obat Indonesia (1)*. Departemen Kesehatan Republik Indonesia. Badan Penelitian dan Pengembangan Kesehatan Jakarta. Jakarta.
47. Syamsuri, A.I., M. W. Alfian, V.P. Muharta, A.T. Mukti, Kismiyati, W.H. Satyaantini. Teknik Pembesaran Ikan Nilem (*Osteochilus hasselti*) di Balai Pengembangan Dan Pemacuan Stok Ikan Gurame dan Nilem (BPPSIGN) Tasikmalaya, Jawa Barat. *Journal of Aquaculture and Fish Health*, 2017; 7(2): 57-62.
48. Tang, U.M. 1999. *Aspek Biologi dan Kebutuhan Lingkungan Larva Ikan Baung (Mystus nemurus C.V)*. Institut Pertanian Bogor. Bogor.
49. Widya, D.R., Suryanto, D., Desrita. Aktivitas Antomikroba Biji Teratai (*Nymphaea purbescens* L.) Terhadap Bakteri *Aeromonas hydrophila*, *Streptococcus agalactine* dan

- Jamur *Saprolegnia* sp. *Jurnal Mikrobiologi*, 2014; 2(1): 7-17.
50. Wijayanti, K. *Pengaruh Pemberian Pakan Alami Yang Berbeda Terhadap Sintasan dan Pertumbuhan Benih Ikan Palmas (Polyterus senegalus senegalus Curvier, 1829)*. Universitas Indonesia. Jakarta, 2010.
51. Zuraidah, S., dan Silkhairi. Penggunaan Larutan Daun Sirih (*Piper betle* L) Dengan Dosis Yang Berbeda Untuk Mencegah Pertumbuhan Jamur (*Saprolegnia* sp.) Pada Telur Ikan Tawes (*Puntius javanicus*). *Jurnal Perikanan Tropis*, 2016; 3(2): 119-130.

Journal Eggs effectiveness

ORIGINALITY REPORT

14%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Texas A&M University, College Station Student Paper	4%
2	garuda.ristekbrin.go.id Internet Source	1%
3	Submitted to Angeles University Foundation Student Paper	1%
4	journal.unram.ac.id Internet Source	1%
5	I Hafizah, Y Aisyah, D Hasni. "Effect of betel type (Piper sp) and concentration of betel leaf extract on quality and antibacterial activities of glycerine bar soap", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1%
6	jppipa.unram.ac.id Internet Source	1%
7	Shara Sasmita, Indah Anggraini Yusanti, Sofian Sofian. "The effectiveness of basil	1%

extract on the mildew saprolegnia within in vitro", JP BIO (Jurnal Pendidikan Biologi), 2021

Publication

8

Meinita, Maria Dyah Nur, Bintang Marhaeni, Tjahjo Winanto, Dwi Setyaningsih, and Yong-Ki Hong. "Catalytic efficiency of sulfuric and hydrochloric acids for the hydrolysis of Gelidium latifolium (Gelidiales, Rhodophyta) in bioethanol production", Journal of Industrial and Engineering Chemistry, 2015.

Publication

1 %

9

repository.unmuhpnk.ac.id

Internet Source

1 %

10

E Setyaningsih, Kismiyati, S Subekti. " The effect of noni fruits () with different ripeness stages against the total erythrocytes and leukocytes of comet goldfish () infested by ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

1 %

11

www.wjpr.net

Internet Source

1 %

12

Bukola Gbemisola Oguntuase, O. T Adebayo. "Sperm Quality and reproductive performance of male Clarias Gariepinus induced with synthetic hormones (Ovatide and Ovaprim)", International Journal of Fisheries and Aquaculture, 2014

1 %

13	jurnalfkip.unram.ac.id Internet Source	1 %
14	www.ijres.org Internet Source	1 %
15	Halimatuddahlia Nasution, Elisa Julianti, Nisaul Fadilah Dalimunthe, Gita Wulandari. "Tensile properties and antimicrobial activity of bioplastic based on sago starch utilized piper betel leaf", AIP Publishing, 2022 Publication	1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%