

# Quality evaluation of polypropylene packaged corn yogurt during storage

*by Nur Aini*

---

**Submission date:** 11-Feb-2022 03:16PM (UTC+0700)

**Submission ID:** 1759941565

**File name:** ISFA.pdf (302.25K)

**Word count:** 4449

**Character count:** 21767

## Quality evaluation of polypropylene packaged corn yogurt during storage

Nur Aini<sup>1,2</sup>, V Prihananto<sup>1</sup>, B Sustriawan<sup>1</sup>, Y Astuti<sup>1</sup> and M R Maulina<sup>1</sup>

Department of Food Science and Technology, Jenderal Soedirman University,  
Purwokerto, Indonesia

Corresponding author : nur.aini@unsoed.ac.id

**Abstract.** Packaging is an important factor to control the process of quality decrease of any food product, including to determine the shelf life. The objective of this study was to determine changes quality of corn yogurt packaged using polypropylene. The method were using was package yogurt polypropylene, then it was stored in a refrigerator at 5, 10, or 15°C during 21 days. The yogurt was analysed every 7 days over a 21-day period. The results indicate that protein content decreased during storage, while the lactic acid bacteria, total acid, pH, viscosity, and total solids were increased. At the end of storage, the amount of lactic acid bacteria still fulfil the minimum requirements of a probiotic food, with a count of 6.407 log CFU/g. Overall scoring by panelist (scores ranged from 0 to 5) have a 4.78 at the beginning of storage. By the 21<sup>st</sup> day of storage, yogurt was packaging using transparent polypropylene having a score of 3.85, and that stored in opaque white packaging having a value of 3.95.

### 1. Introduction

Yoghurt is one of dairy-based probiotic products which is much sold on the market. People who are allergic to milk protein require an alternative ingredient of yoghurt. For example, Yoghurt with vegetable as the raw material such as peanuts and soybeans [1], [2]. [3] developed probiotic corn extract which has similar characteristic to yogurt in which it has 8.74 log CFU/g of total lactic acid bacteria (BAL). This shows that sweet corn extract is qualified as a probiotic food which requires lactic acid bacteria of 6 log CFU/g [4].

Changes during the storage will affect the food quality. The stability of food products is related to the speed of physical damage due to the changes of chemical substance, physic and microbiology. Temperature is a factor which affects the changes in food product quality. When food is stored, the storage room temperature is constant. For fermentation products' storage, the temperature is very much affect the storage time. For fermentation products whose bacteria is alive, its storage time is short and it should be stored at 4° C. According to [5], the storage of yogurt at 4°C will decrease the viability of lactic acid bacteria.

Pasteurized or sterilized fermentation products have a longer storage time, but its condition also depends on the type of packaging [6]. The packaging commonly used for yoghurt is a polypropylene cup due to its small surface area so it can reduce the contact of the ingredients with oxygen. Generally,



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

the polypropylene packaging used is transparent and opaque. The opaque polypropylene packaging is not transparent so it can provide different qualities during the storage.

The aim of this research is to know the quality change of corn yoghurt which is packed with transparent and opaque polypropylene at 5, 10 and 15°C.

## 2. Experimental details

### 2.1. The making of corn yogurt

The method used to produce corn yogurt was that reported by [3]. A corn extract was prepared and combined with 15% sweet potato, 10% mung bean extract, 15% sugar, and 10% skim milk powder. The mixture was pasteurized at 70°C for 15 minutes, then cooled to 40-43°C. A mixed culture of *Lactobacillus bulgaricus* and *Streptococcus thermophilus* were then added. The final mixture was incubated for 8 hours at 37°C.

### 2.2. Package and storage of corn yogurt

Corn yogurt was then packed with transparent and white cups. Storage was done on refrigerator at 5°C, 10°C and 15°C for 21 days. The data were analyzed every 7 days for 21 days.

### 2.3. Analysis of samples

The research variables observed was total lactic acid bacteria using plate count method and M.R.S Agar media [7]; the pH measurement was carried out by potentiometric method, by using pH meters in all experimental units, total soluble solids (using refractometer), lactic acid levels determined by titration using alkaline solution (Mann's Acid Test), total protein level (micro Kjeldahl method). Sensory analysis including color, acid flavor, aroma and preference was done by using scoring test. Panelists used were trained panelists consisting of 20 people. The scoring is on a scale of 1 to 5.

### 2.4. Analysis of data


Data analysis was done by using simple linear regression method and descriptive analysis.

## 3. Result and discussion

### 3.1. Viability of lactic acid bacteria

According to [8], *Lactobacillus* is a potential lactic acid bacteria (BAL) to make food to be functional. At first, the amount of BAL was 7,573 log CFU/ml, which fulfilled the minimum requirement of probiotic food. According to [9], the total minimum of BAL in probiotic food is 6 log CFU/g at expiration time.

During the storage, BAL decreased slightly from 0.67 to 0.86 log cycle. Until the 21<sup>st</sup> day, the amount of BAL still met the minimum requirement of probiotic food, in which the lowest was 6.407 log CFU/ml (Figure 1). The decrease of BAL during this storage was better than [5] result in which BAL of yogurt decreased as much as 2.34 log cycles during the 28 days storage time. The decrease viability of lactic acid bacteria is caused by the excessive production of acids resulting in the lactic acid bacteria death. Meanwhile, according to [10] the more total BAL exists in food, the tighter the competition among BAL will be. It means that the more total BAL in food will decrease the availability of nutrients and the BAL survival. So, the longer the storage time, the number of BAL will decrease.




**Figure 1.** The number of lactic acid bacteria in corn yogurt using transparent and opaque packaging

The amount of lactic acid bacteria in yogurt using transparent and opaque packaging tended to be the same during the storage. According to [4], the ability of bacteria to grow and develop is influenced by the nutrition and environmental conditions. Nutrition that can be used by BAL to survive is proteins, fats and minerals (calcium, phosphorus and magnesium), of which have the same amount.

### 3.2. Protein content

During the storage, the protein content of corn yoghurt tended to decrease until the 7<sup>th</sup> day, after that, it tended to increase until the 21<sup>st</sup> day. The protein levels decreased until the 7<sup>th</sup> day because *Lactobacillus* was proteolytic, breaking proteins into simple peptides which was decreasing the protein levels.

Protein levels which tended to increase after the 7<sup>th</sup> day were likely because the protease enzymes degraded the proteins in microbes. Microbes are composed by various components, including 60-70% of protein. According to [11], microbial cells lysis during the storage. The protease and peptidase enzymes will break the protein down in the BAL, so the more cells lyse, the higher the protein level in yoghurt. This protein changes pattern is almost the same for transparent and opaque polypropylene packaging yogurt during the storage, but for yoghurt which was stored at the highest temperature (5°C), the protein level tends to be higher.



**Figure 2.** The protein content of corn yogurt was packaged by transparent and opaque polypropylene during the storage

### 3.3. Total acid and pH

The most produced organic acids in yogurt and as the main product is lactic acid [12]. During the storage, total acids in each package [15] and temperature tended to increase, initially from 0.765 to 0.819-1.026 (Table 1). This shows that the production of lactic acid in corn yoghurt had increased during the storage. This is similar to [5] result in which during the 21 days storage, lactic acid levels increased to 1.11 until 1.42%. [13] also almost the same, in which during the 21 days storage, the total acidity of yogurt is 1.

According to [11], lactic acid bacteria has the optimum temperature to produce acids at 10-40°C. At 10°C and 15°C, the production of lactic acid approaches the optimal temperature of lactic acid bacteria activity so that the production of lactic acid increases. This corn yogurt also fulfill the standard of yogurt SNI no 01-2981-1992 which has total acid of 0.5-2.

**Table 1.** The pH and total acid of corn yogurt was packaged by polypropylene during storage

Packaging	Storage temperature	pH on the day				Total acid on the day (%)			
		0	7	14	21	0	7	14	21
Transparent	5	4.4	4.5	4.5	4.7	0.765	0.774	0.882	0.882
	10	4.4	4.4	4.4	4.6	0.765	0.819	1.053	0.999
	15	4.4	4.5	4.5	4.6	0.765	0.81	0.882	0.918
Opaque white	5	4.4	4.5	4.5	4.6	0.765	0.783	0.864	0.819
	10	4.4	4.4	4.4	4.6	0.765	0.837	0.927	1.026
	15	4.4	4.5	4.5	4.6	0.765	0.747	0.963	0.945

Lactic acid produced during the fermentation can increase the acidity or decrease the pH. In this research, there was no decrease in pH, instead the pH increased (Table 1). On the 21<sup>st</sup> day, the pH which was originally 4.4 became 4.6-4.7. This was due to the delayed growth of lactic acid bacteria and the growth of proteolytic and other lipolytic organisms which have the ability to perform lactic

acid metabolism and therefore produced base compounds. The increase of pH of yoghurt during the storage is also consistent with [14] which stated that the pH of yogurt stored in cold temperatures ( $4 \pm 1^\circ\text{C}$ ) tends to increase during the storage. The initial value of pH which was 3.2 rose to 3.5 on the 10<sup>th</sup> day and 3.8 on the 35<sup>th</sup> day.

[15] reported that increasing the total acid in yogurt does not always lead to a decrease in pH. [6] also reported that a low storage temperature can inhibit the action of lactase, leading to a reduction in lactic acid production and no decrease in pH.

The production of lactic acid was increased during storage. This is consistent with [5], who found that the lactic acid content of corn yogurt increased from 1.11 to 1.42% during 21 days of storage. [13] also reported that on the 21<sup>st</sup> day, the total acidity of corn yogurt was 1%.

### 3.4. Total solid and viscosity

During the storage, total solids of corn yoghurt decreased slightly as shown in Table 2. The total solids of corn yoghurt was initially 23°Brix, and on the 21<sup>st</sup> day it became 22-22.7°Brix. This total decrease of solids is similar to [13] that during the storage, the total solids of yoghurt decrease. Despite the decreases during the storage, this total soluble solids still fulfil the SNI of yogurt standard of at least 8.2°Brix.

**Table 2.** Total solid and viscosity of corn yogurt was packaged by polypropylene during storage

Packaging	Storage temperature	Total solid at day (°Brix)				Viscosity at day (cP)			
		0	7	14	21	0	7	14	21
Transparent	5	23	23	23	23	160	190	170	160
	10	23	23	23	22,7	160	160	180	140
	15	23	23	23	22,7	160	170	180	140
Opaque white	5	23	23	23	22,5	160	180	180	200
	10	23	23	23	22	160	150	180	190
	15	23	23	23	22,6	160	150	180	160

The desired viscosity of yogurt is viscous liquids and homogeneous consistency. According to [16], the viscosity of yogurt is also influenced also by milk protein, which is casein. At pH close to 4.6, the solubility of casein is lost therefore the hydrophobic interaction occurs between the casein forming the structure and the main consistency of yogurt, and the process may affect the physic of yogurt, such as texture, viscosity, water holding capacity, and syneresis.

Casein is coagulated as the degree of acidity decreases due to the production of lactic acid. Lactic acid can agglomerate milk protein, sweet corn protein or mungbean protein. [16] explained that lactic acid leads to the change of casein/phosphate complexes to be soluble fraction of calcium and phosphate so that the casein destabilizes.

During the storage, there is increased yoghurt viscosity due to the increase of lactic acid. This can be seen from yogurt which was stored in opaque packaging until the 21<sup>st</sup> day and transparent packaging until the 14<sup>th</sup> day. It is caused by syneresis or separation of whey causing unstable curd. Syneresis can occur due to high storage temperatures, low total solids in milk, and no vibration during transport or during storage. Syneresis occurred because the transparent packaging can be penetrated by light that disrupts the stability of yogurt.



At the end of storage, the opaque packaging corn yoghurt has a higher viscosity than the transparent packaging. Storage was done in the refrigerator which has light bulb which directly highlights the product. During the storage, there is increased yoghurt viscosity due to the increase of lactic acid. This can be seen from yogurt which was stored in opaque packaging until the 21<sup>st</sup> day and transparent packaging until the 14<sup>th</sup> day. It is caused by syneresis or separation of whey causing unstable curd. Syneresis can occur due to high storage temperatures, low total solids in milk, and no vibration during transport or during storage. Syneresis occurred because the transparent packaging can be penetrated by light that disrupts the stability of yogurt.

Yogurt is required to have the form of a viscous fluid, with a homogenous consistency. According to [17], the viscosity of yogurt is also influenced by milk protein, i.e., casein. At a pH of approximately 4.6, casein becomes insoluble, resulting in hydrophobic interactions between casein micelles that form the main structure and consistency of yogurt. This process can influence the physical properties of yogurt, including its viscosity, water-holding capacity, and syneresis.

Casein is coagulated when acidity decreases due to lactic acid production. Lactic acid can agglomerate protein in milk, corn, or mungbean. [17] explain that the production of lactic acid resulted in a complex change whereby casein-phosphate was converted into soluble calcium and phosphate, with the casein being destabilized.

### 3.5. Sensory properties

The yoghurt quality is determined by texture or viscosity, the degree of acidity and flavor which is highly influential on sensory quality and consumer acceptance. Sensory analysis done included color, flavor, aroma and overall acceptance.

Corn yoghurt is yellow due to the presence of carotenoids. Color changes also occur because of oxidation reactions which can cause changes in flavor, color, and microbial growth in large quantities which also can damage the color or appearance of food.

Color is an important food attribute and becomes the first characteristic seen by consumers. The color of yoghurt according to the panelist assessment is 4.97 (yellow). Yellow color in yoghurt is caused by the presence of carotenoids in corn and sweet potatoes. The longer the storage, the more decrease the carotenoid color pigment will be. This is due to the degradation of pigments carried out by lactic acid bacteria. According to [16] the color changes of the fermentation medium which is getting faded after the fermentation can be caused by anthocyanin pigment degradation due to the decrease of acidity (pH) during the fermentation process. Yogurt which packaged by white opaque polypropylene was more yellow (4.2-4.35) than yogurt which packaged by transparent polypropylene (4.15) as shown in Table 3.

The formation of lactic acid in yogurt resulted in sour taste. The panelist rate on the yoghurt sour taste at the beginning of storage was 3.98 (acid), and at the end of the storage range at 3.95 (sour) to 4.4 (very sour). According to [18] the sour taste characteristic of yogurt is due to their lactic acid. The increased acid taste during storage in accordance with [13] that sour taste during the storage increased because the amount of lactic acid increased.

The panelist assessment of yogurt aroma indicated that yogurt aroma decreased during the storage. At first, the aroma value was 4.85 and at the end of the storage (the 21<sup>st</sup> day) it decreased to 3.5 - 4.35. This is caused by alcohol formed as the result of further degradation of lactic acid during the storage. According to [5], this is considered to be due to a contamination of fermentative yeast that will form alcohol during the storage.

According to [18], flavor yogurt compounds consists of 1) non-volatile acids (lactic acid, pyruvate or succinate); 2) volatile acids (formic acid, acetate, propionate or butyrate); 3) carbonyl compounds (acetaldehyde, acetone, or diacetyl), and 4) other compounds (amino acids and compounds formed by thermal degradation of proteins, fats or lactose). Among these compounds, acetaldehyde and diacetyl is the most predominant in determining the scent of yoghurt. According to [5],

acetaldehyde and other carbonyl compounds will decrease during the storage of yoghurt products. In addition, during the storage, there may be oxidation so that the volatile compounds evaporate. This results in the decrease of yoghurt aroma during the storage.

**Table 3.** Sensory properties of corn yogurt was packaged by polypropylene during storage

Sensory attributes	Packaging		Day			
			0	7	14	21
Color	Transparent	5	4.97	4.4	4.35	4.15
		10	4.97	4.4	4.1	4.1
		15	4.97	4.35	4.1	4.1
	Opaque white	5	4.97	4.45	4.35	4.3
		10	4.97	4.4	4.4	4.35
		15	4.97	4.25	4.25	4.2
Flavour	Transparent	5	4.85	4.4	4.5	4.35
		10	4.85	3.85	3.75	3.75
		15	4.85	4.05	4	3.95
	Opaque white	5	4.85	4.3	4.25	4.15
		10	4.85	4.45	3.55	3.5
		15	4.85	4.35	3.85	3.7
Taste	Transparent	5	3.98	3.85	4.15	4.2
		10	3.98	3.95	4	4.15
		15	3.98	3.8	3.9	3.95
	Opaque white	5	3.98	3.8	4.05	3.95
		10	3.98	3.7	4.05	4.25
		15	3.98	3.75	4.2	4.4
Overall acceptance	Transparent	5	4.78	4.5	4.1	3.85
		10	4.78	4.4	3.3	3.15
		15	4.78	4.35	3.75	2.7
	Opaque white	5	4.78	4.55	4.15	3.85
		10	4.78	4.65	4	3.3
		15	4.78	4.5	3.2	2.85

Panelists' preference on yogurt is affected by several factors, including flavor, aroma, and consistency. According to [18] the preference on yoghurt is much related to the fermentation by lactic



acid bacteria which resulted in a distinctive flavor of yoghurt. The stability of yoghurt also affects the preference, which can be seen from no occurrence of yoghurt damage in the form of wheying off or syneresis. According to [5] moderate viscosity and subtle texture are preferred by consumers.

The longer the storage time, the more decrease the panelists' preference judgment on yogurt is. At first, panelist's preference on yogurt was 4.78 and after storage for 21 days, panelists' preference on yogurt was 2.7-3.85 (Table 3). In metabolism reactions, the acid will turn into alcohol so that other aromas which arise are the result of the process. This is also reason of the decrease preference of panelists' preference on yoghurt.

#### 4. Conclusion

During corn yogurt storage, the protein content decreased, while the total acid, pH, viscosity, and total solids remained consistent for both types of packaging at all three temperatures tested. At the end of storage, the amount of lactic acid bacteria still fulfil the minimum requirements of a probiotic food, with a count of 6.407 log CFU/g. Panel ratings of sensory variables tended to decrease as the length of storage increased. Corn yogurt was rated by a sensory assessment panel, and scores ranged from 0 to 5, with a mean rating of 4.78 at the beginning of storage. By the 21<sup>st</sup> day of storage, these ratings had decreased, with yogurt stored in transparent packaging having a rating of 3.85, and that stored in opaque white packaging having a value of 3.95.

#### Acknowledgement

This study was supported by the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia under the National Strategic Research program in 2016 grant number 2045/UN23.14/PN/2016.

#### Reference

- [1] J. Isanga and G. Zhang, "Production and evaluation of some physicochemical parameters of peanut milk yoghurt," *LWT-Food Sci. Technol.*, vol. 42, no. 6, pp. 1132–1138, 2009.
- [2] M. Sveučilište u Zagrebu. Prehrambeno-biotehnološki fakultet., L. Hrvatsko društvo za biotehnologiju., Saveza hrvatskih društava za unapređenje prehrane., and Slovenskog mikrobiološkog društva., "Physicochemical, Textural and Sensory Characteristics of Probiotic Soy Yogurt Prepared from Germinated Soybean," *Food Technol. Biotechnol.*, vol. 48, no. 4, pp. 490–496, Dec. 2010.
- [3] N. Aini, V. Prihananto, G. Wijonarko, A. Arimah, and M. Syaifudin, "Effect of Culture Concentration and Sweet Potato Prebiotic to the Properties of Sweet Corn Juice Probiotic," *Agritech*, vol. 37, no. 2, pp. 165–172, 2017.
- [4] D. A. . Saccaro, A. Y. Tamime, S. Pilleggi, Ana Lu Cia O.P, and M. N. Oliveira, "The viability of three probiotic organisms grown with yoghurt starter cultures during storage for 21 days at 4 C," *Int. J. Dairy Technol.*, vol. 62, no. 3, pp. 397–404, 2009.
- [5] M. R. Damin, E. Minowa, M. R. Alcantara, Oliveira, and Marice Nogueira, "Effect of Cold Storage on Culture Viability and Some Rheological Properties of Fermented Milk Prepared with Yogurt and Probiotic Bacteria," *J. Texture Stud.*, vol. 39, no. 1, pp. 40–55, Feb. 2008.
- [6] D. . Johnson, S. Duncan, L. . Bianchi, and H. . Chang, "Packaging modifications for protecting flavor of extended-shelf-life milk from light," *J. Dairy Sci.*, vol. 98, pp. 1–10, 2015.
- [7] P. Romano, C. Fiore, M. Paraggio, M. Caruso, and A. Capece, "Function of yeast species and strains in wine flavour," vol. 86, pp. 169–180, 2003.
- [8] W. Gustaw, M. Kordowska-Wiater, and J. Kozio, "The influence of selected prebiotics on the growth of lactic acid bacteria for bio-yoghurt production," *Acta Sci. Polym. Technol.*, vol. 10,

- no. 4, pp. 455–466, 2011.
- [9] A. Y. Tamime, M. Saarela, A. K. Sondergaard, V. V. Mistry, and Shah. N.P, “Production and maintenance of viability of probiotic microorganisms in dairy product,” *Probiotic Dairy Prod.*, vol. 3, pp. 39–63, 2005.
- [10] M. Gurses, B. Cetin, and M. Sengul, “Change in Probiotic Microflora and Physico-chemical Characteristics of Rose Hip Marmalade Bio-yoghurts During Refrigerated Storage,” *Asian J. Chem.*, vol. 21, no. 5, pp. 4097–4103, 2009.
- [11] M. Granito and G. Álvarez, “Lactic acid fermentation of black beans (*Phaseolus vulgaris*): microbiological and chemical characterization,” *J. Sci. Food Agric.*, vol. 86, no. 8, pp. 1164–1171, Jun. 2006.
- [12] E. Mani-López, E. Palou, and A. López-Malo, “Probiotic viability and storage stability of yogurts and fermented milks prepared with several mixtures of lactic acid bacteria,” *J. Dairy Sci.*, vol. 97, no. 5, pp. 2578–2590, May 2014.
- [13] T. E. Obi, F. . Henshaw, and O. . Atanda, “Quality Evaluation of Plain-Stirred Probiotic Yoghurt Produced From Skim and Whole Milk Powder during Refrigerated Storage,” *Electron. J. Environ. Agric. Food Chem.*, vol. 9, no. 7, pp. 1203–1213, 2010.
- [14] A. G. Evrendilek, “Survival of *Escherichia coli* O157:H7 in yogurt drink, plain yogurt and salted (tuzlu) yogurt: Effects of storage time, temperature, background flora and product characteristics,” *Int. J. Dairy Technol.*, vol. 60, no. 2, pp. 118–122, May 2007.
- [15] A. B. Dantas *et al.*, “Manufacture of probiotic Minas Frescal cheese with *Lactobacillus casei* Zhang,” *J. Dairy Sci.*, vol. 99, no. 1, pp. 18–30, Jan. 2016.
- [16] A. P. Marafon, A. Sumi, M. R. Alcântara, A. Y. Tamime, and M. Nogueira de Oliveira, “Optimization of the rheological properties of probiotic yoghurts supplemented with milk proteins,” *LWT - Food Sci. Technol.*, vol. 44, no. 2, pp. 511–519, 2011.
- [17] A. P. Marafon, A. Sumi, M. R. Alcântara, A. Y. Tamime, and M. Nogueira de Oliveira, “Optimization of the rheological properties of probiotic yoghurts supplemented with milk proteins,” *LWT - Food Sci. Technol.*, vol. 44, no. 2, pp. 511–519, 2011.
- [18] L. Allgeyer, M. Miller, and S. Lee, “Sensory and microbiological quality of yogurt drinks with prebiotics and probiotics,” *J. Dairy Sci.*, 2010.

# Quality evaluation of polypropylene packaged corn yogurt during storage

## ORIGINALITY REPORT

13%	12%	8%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://doc-pak.undip.ac.id">doc-pak.undip.ac.id</a> Internet Source	4%
2	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	3%
3	<a href="http://www.insightsociety.org">www.insightsociety.org</a> Internet Source	1%
4	<a href="http://psasir.upm.edu.my">psasir.upm.edu.my</a> Internet Source	1%
5	Béal, Catherine, and Gisèle Chammas. "Production of Laban", Handbook of Animal- Based Fermented Food and Beverage Technology Second Edition, 2012. Publication	<1%
6	<a href="http://ebin.pub">ebin.pub</a> Internet Source	<1%
7	<a href="http://www.foodandnutritionjournal.org">www.foodandnutritionjournal.org</a> Internet Source	<1%
8	<a href="http://pubs.sciepub.com">pubs.sciepub.com</a> Internet Source	

<1 %

9

Olga S. Papadopoulou, Anthoula A. Argyri, Evangelos E. Varzakis, Chrysoula C. Tassou, Nikos G. Chorianopoulos. "Greek functional Feta cheese: Enhancing quality and safety using a Lactobacillus plantarum strain with probiotic potential", Food Microbiology, 2018

Publication

<1 %

10

[vuir.vu.edu.au](http://vuir.vu.edu.au)

Internet Source

<1 %

11

MARK A. BUONO. "CARBOHYDRATE UTILIZATION and GROWTH KINETICS IN the PRODUCTION of YOGURT FROM SOYMILK. PART II: EXPERIMENTAL and PARAMETER ESTIMATION RESULTS", Journal of Food Processing and Preservation, 6/1990

Publication

<1 %

12

[itjfs.com](http://itjfs.com)

Internet Source

<1 %

13

[journals.tubitak.gov.tr](http://journals.tubitak.gov.tr)

Internet Source

<1 %

14

R D Andriani, P P Rahayu, M W Apriliyani, M E Sawitri, A Manab, A R Azkariahman. "Anti-Obesity Effect of Yoghurt Synbiotic in High Fat Diet Induced Wistar Rats", IOP Conference Series: Earth and Environmental Science, 2020

<1 %

- |    |   |      |
|----|---|------|
| 15 | <a href="http://www.hindawi.com">www.hindawi.com</a><br>Internet Source   | <1 % |
| 16 | <a href="http://www.mdpi.com">www.mdpi.com</a><br>Internet Source   | <1 % |
| 17 | "Lactic Acid Bacteria", Springer Nature, 2014<br>Publication  | <1 % |
| 18 | N Aini, B Sustriawan, V Prihananto, T Heryanti.<br>"Characteristics of cheese analogue from corn<br>extract added by papain and pineapple<br>extract", IOP Conference Series: Earth and<br>Environmental Science, 2019<br>Publication         | <1 % |
| 19 | <a href="http://iopscience.iop.org">iopscience.iop.org</a><br>Internet Source   | <1 % |
| 20 | Gueimonde, Miguel, Clara de Los Reyes-<br>Gavilán, and Borja Sánchez. "Stability of Lactic<br>Acid Bacteria in Foods and Supplements",<br>Lactic Acid Bacteria Microbiological and<br>Functional Aspects Fourth Edition, 2011.<br>Publication | <1 % |

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Quality evaluation of polypropylene packaged corn yogurt during storage

## GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9



## CLAIM

Take an arguable position on the scientific topic and develop the essay around that stance.

---

ADVANCED	The essay introduces a precise, knowledgable, qualitative and/or quantitative claim based on the scientific topic or text(s), regarding the relationship between dependent and independent variables, and establishes the significance of the claim. The essay develops the claim and counterclaim fairly and thoroughly, distinguishing the claim from alternate or opposing claims.
PROFICIENT	The essay introduces a precise, qualitative and/or quantitative claim based on the scientific topic or text(s), regarding the relationship between dependent and independent variables. The essay develops the claim and counterclaim fairly, distinguishing the claim from alternate or opposing claims.
DEVELOPING	The essay introduces a qualitative and/or quantitative claim based on the scientific topic or text(s), regarding the relationship between dependent and independent variables, but it may be somewhat unclear. The essay may not present the claim and counterclaim evenly or objectively, failing to adequately distinguish the claim from alternate or opposing claims.
EMERGING	The essay does not clearly make a qualitative and/or quantitative claim based on the scientific topic or text(s), or the claim is overly simplistic or vague. The essay does not address counterclaims.

## EVIDENCE

Include relevant facts, definitions, and examples to back up the claim.

---

ADVANCED	The essay supplies the most relevant and appropriate qualitative and/or quantitative data and evidence related to the scientific topic or text(s) to support its claim and counterclaim.
PROFICIENT	The essay supplies sufficient qualitative and/or quantitative data and evidence related to the scientific topic or text(s) to support its claim and counterclaim.
DEVELOPING	The essay supplies some qualitative and/or quantitative data and evidence, but it may not be closely related to the scientific topic or text(s), thereby not effectively supporting the essay's claim and counterclaim.
EMERGING	The essay supplies very little or no qualitative and/or quantitative data and evidence to support its claim and counterclaim.

## REASONING

Explain how or why each piece of evidence supports the claim.

---

ADVANCED	The essay applies scientific ideas and principles in order to thoroughly explain how or why the cited evidence supports the claim. The essay demonstrates consistently logical reasoning and full understanding of the scientific topic and/or text(s), comprehensively stating the strengths and limitations of both the claim
----------	---

and counterclaim. The essay's explanations effectively anticipate the audience's knowledge level, concerns, values, and potential biases about this scientific topic.

PROFICIENT	The essay applies scientific ideas and principles in order to explain how or why the cited evidence supports the claim. The essay demonstrates some logical reasoning and understanding of the scientific topic and/or text(s) and states the strengths and limitations of both the claim and counterclaim. The essay's explanations anticipate the audience's knowledge level and concerns about this scientific topic.
DEVELOPING	The essay includes some reasoning and understanding of the scientific topic and/or text(s), but it does not effectively apply scientific ideas and principles to explain how or why the evidence supports the claim. The strengths and limitations of the claim and counterclaim may be unclear.
EMERGING	The essay does not demonstrate clear or relevant reasoning to support the claim or to demonstrate an understanding of the scientific topic and/or text(s).

## FOCUS

Focus your writing on the purpose and task.

---

ADVANCED	The essay maintains strong focus on the purpose and task, using the whole essay to support and develop the claim and counterclaims fairly while thoroughly addressing the demands of the prompt.
PROFICIENT	The essay maintains focus on the purpose and task, using most of the essay to support and develop the claim and counterclaims while thoroughly addressing the demands of the prompt.
DEVELOPING	The essay may not fully address the demands of the prompt or stay focused on the purpose and task. The writing may stray off topic at times.
EMERGING	The essay does not maintain focus on purpose or task.

## ORGANIZATION

Organize your writing in a logical sequence.

---

ADVANCED	The essay incorporates an effective organizational structure throughout that logically sequences the claim(s), counterclaims, reasons, and evidence. It also includes an effective introduction and a conclusion which follows from and supports the arguments presented. Transitional words and phrases are included to enhance the relationships between and among ideas (i.e. claim and reasons, reasons and evidence, claim and counterclaim) and to create cohesion and clarity throughout.
PROFICIENT	The essay incorporates an adequate organizational structure and transitional phrases throughout that establish clear relationships between and among the claim(s), counterclaims, reasons, and evidence, including an introduction and concluding statement or section.

DEVELOPING	The essay uses a basic organizational structure but relationships between and among ideas are not consistently clear. The essay moves from beginning to end; however, an introduction and/or conclusion may be overly formulaic, repetitious, or missing.
EMERGING	The essay does not have a clear organizational structure. An introduction and conclusion are not evident.

## LANGUAGE

Pay close attention to your tone, style, word choice, and sentence structure when writing.

---

ADVANCED	The essay effectively establishes and maintains a formal style and objective tone and incorporates language that anticipates the reader's knowledge level, concerns, values, and potential biases. The essay demonstrates a clear command of conventions, while also consistently employing discipline-specific word choices and varied sentence structure.
PROFICIENT	The essay generally establishes and maintains a formal style and somewhat objective tone with few possible exceptions and incorporates language that anticipates the reader's knowledge level and concerns. The essay demonstrates a general command of conventions, while also employing discipline-specific word choices and some variety in sentence structure.
DEVELOPING	The essay attempts a formal style and objective tone, and language inconsistently shows an awareness of the reader's knowledge or concerns. The essay may contain errors in conventions that interfere with meaning. Some attempts at discipline-specific word choices are made, and sentence structure may not vary.
EMERGING	The essay is not formal in style or objective in tone and employs language that does not anticipate the reader's knowledge or concerns. The essay may contain pervasive errors in conventions that interfere with meaning. Word choice is not discipline-specific, and sentence structures are simplistic and unvaried.