

Transpiration on the rebound in lowland Sumatra

by Afik Hardanto

Submission date: 25-Mar-2023 09:52AM (UTC+0700)

Submission ID: 2045955743

File name: 10.1016_j.agrformet.2019.04.017.pdf (1.22M)

Word count: 14401

Character count: 77549



Transpiration on the rebound in lowland Sumatra

A. Röhl^{a,*}, F. Niu^{a,b}, A. Meijide^{c,d,e}, J. Ahongshangbam^a, M. Ehbrecht^f, T. Guillaume^{g,h,i},
D. Gunawan^j, A. Hardanto^{a,k}, Hendrayanto^l, D. Hertel^m, M.M. Kotowska^{m,n}, H. Kreft^{o,p},
Y. Kuzyakov^{g,p}, C. Leuschner^{m,p}, M. Nomura^o, A. Polle^{p,q}, K. Rembold^{o,r}, J. Sahner^q, D. Seidel^f,
D.C. Zemp^{a,o}, A. Knohl^{c,p}, D. Hölscher^{a,p}

^a University of Goettingen, Tropical Silviculture and Forest Ecology, Germany

^b University of Arizona, School of Natural Resources and the Environment, USA

^c University of Goettingen, Bioclimatology, Germany

^d Universidad de Granada, Departamento de Ecología, Spain

^e University of Goettingen, Department of Crop Sciences, Division Agronomy, Germany

^f University of Goettingen, Silviculture and Forest Ecology of the Temperate Zones, Germany

^g University of Goettingen, Soil Science of Temperate Ecosystems and Agricultural Soil Science, Germany

^h Ecole Polytechnique Fédérale de Lausanne, Ecological Systems Laboratory, Switzerland

ⁱ Swiss Federal Institute for Forest, Snow and Landscape Research, Site Lausanne, Switzerland

^j Agency for Meteorology, Climatology and Geophysics, Center for Climate Change Information, Jakarta, Indonesia

^k Jenderal Soedirman University, Faculty of Agriculture, Indonesia

^l Bogor Agricultural University, Forest Management, Indonesia

^m University of Goettingen, Plant Ecology and Ecosystems Research, Germany

ⁿ Macquarie University, Biological Sciences, Australia

^o University of Goettingen, Biodiversity, Macroecology and Biogeography, Germany

^p University of Goettingen, Centre of Biodiversity and Sustainable Land Use, Germany

^q University of Goettingen, Forest Botany and Tree Physiology, Germany

^r University of Bern, Botanical Garden Bern, Switzerland

ARTICLE INFO

Keywords:

Forest
Jungle rubber
Land-cover change
Land-use intensification
Oil palm
Tropics
Rubber plantation
Sap flux
Water use

ABSTRACT

Following large-scale conversion of rainforest, rubber and oil palm plantations dominate lowland Sumatra (Indonesia) and other parts of South East Asia today, with potentially far-reaching ecohydrological consequences. We assessed how such land-use change affects plant transpiration by sap flux measurements at 42 sites in selectively logged rainforests, agroforests and rubber and oil palm monoculture plantations in the lowlands of Sumatra. Site-to-site variability in stand-scale transpiration and tree-level water use were explained by stand structure, productivity, soil properties and plantation age. Along a land-use change trajectory forest-rubber-oil palm, time-averaged transpiration decreases by $43 \pm 11\%$ from forest to rubber monoculture plantations, but rebounds with conversion to smallholder oil palm plantations. We uncovered that particularly commercial, intensive oil palm cultivation leads to high transpiration ($827 \pm 77 \text{ mm yr}^{-1}$), substantially surpassing rates at our forest sites ($589 \pm 52 \text{ mm yr}^{-1}$). Compared to smallholder oil palm, land-use intensification leads to 1.7-times higher transpiration in commercial plantations. Combined with severe soil degradation, the high transpiration may cause periodical water scarcity for humans in oil palm-dominated landscapes. As oil palm is projected to further expand, severe shifts in water cycling after land-cover change and water scarcity due to land-use intensification may become more widespread.

1. Introduction

Climate is governed by changes in atmospheric water vapour content and associated feedback mechanisms (Inamdar and Ramanathan, 1998; Held and Soden, 2000; Ellison et al., 2012). Terrestrial water

fluxes from land surface to the atmosphere are dominated by plant transpiration (Jasechko et al., 2013; Good et al., 2015), which in turn is strongly affected by human land use. Land-cover change by forest conversion to agricultural systems usually substantially decreases (evapo)transpiration (Sampaio et al., 2007; Ellison et al., 2012; Silvério

* Corresponding author at: University of Goettingen, Tropical Silviculture and Forest Ecology, Buesgenweg 1, 37077, Goettingen, Germany.

E-mail address: aroell@gwdg.de (A. Röhl).

<https://doi.org/10.1016/j.agrformet.2019.04.017>

Received 1 February 2019; Received in revised form 13 April 2019; Accepted 22 April 2019

Available online 13 May 2019

0168-1923/© 2019 Elsevier B.V. All rights reserved.

et al., 2015), which is associated with increases in land surface temperature (Alkama and Cescaati, 2016; Ellison et al., 2017; Sabajo et al., 2017) and decreases in regional precipitation (Bagley et al., 2014; Spracklen and Garcia-Carreras, 2015; Zemp et al., 2017a, 2017b). Potential effects of land-use intensification have received less attention so far. Recent research proposes that increases in cropland extent, productivity and intensification could influence climate through increasing (evapo)transpiration (Betts et al., 2013; Mueller et al., 2016, 2017).

Today, large-scale forest conversion mainly occurs in tropical regions, with currently highest rates in Southeast Asia and particularly in Indonesia (Hansen et al., 2013; Margono et al., 2014). In many Indonesian post-conversion landscapes, rubber and oil palm monocultures are dominant land-use types, while remaining rainforests have largely been selectively logged or degraded otherwise (FAO, 2014; Abood et al., 2015; Clough et al., 2016). Rubber has been cultivated in Indonesia since the early 1910s, formerly often in so-called 'jungle rubber' agroforests, a mixture of planted and spontaneously established trees of various ages (Gouyon et al., 1993; Joshi et al., 2003). Today, monocultural rubber plantations are much more widespread, and over the past 25 years oil palm monocultures have strongly increased in area (FAO, 2014; Abood et al., 2015; Clough et al., 2016).

Forest conversion to rubber and oil palm monocultures decreases biodiversity (Clough et al., 2016), ecological functioning (Barnes et al., 2014), soil fertility (Guillaume et al., 2015, 2016) and carbon storage (Guillaume et al., 2018) and leads to changes in microclimate (Meijide et al., 2018). In Amazonia, large-scale forest conversion to pasture and soy bean regionally decreased evapotranspiration, subsequently reducing downwind precipitation and resulting in more frequent and severe droughts (Bagley et al., 2014; Spracklen and Garcia-Carreras, 2015; Zemp et al., 2017a, 2017b). In contrast, woody perennial rubber and oil palm plantations in Southeast Asia could be more similar to forests regarding transpiration than Amazonian pastures or soy bean, and ecohydrological consequences thus potentially less severe. Previous studies found indications of quite high evapotranspiration from oil palm plantations (e.g. Radersma and de Ridder, 1996). Recent studies report very high annual transpiration and evapotranspiration from mature commercial, intensively managed oil palm plantations (Meijide et al., 2017; Manoli et al., 2018). It was indicated that smallholder oil palm plantations had lower transpiration than commercial plantations, but the study was restricted to short periods and did not provide annual estimates (Röhl et al., 2015). Smallholder rubber plantations in lowland Sumatra had lower transpiration compared to adjacent smallholder oil palm plantations and also compared to commercial rubber plantations on the Asian mainland (Niu et al., 2017). To our knowledge, there are no previous studies in Southeast Asia that directly compare transpiration of different land-use types including rubber and oil palm with adjacent reference forest sites to allow a more comprehensive ecohydrological assessment. As oil palm is one of the most rapidly expanding crops worldwide (FAO, 2014), there is an urgent need to better understand the ecological consequences of its large-scale cultivation.

Establishing representative means of (evapo)transpiration for certain land-use types is one important aspect of ecohydrological studies; it can be accomplished by a high number of spatial replicates. Of further importance for analysing effects of land-cover and land-use change at the landscape level is to assess and understand variability, both within and across land-use types. Hourly, daily and seasonal fluctuations in transpiration at a certain site can commonly be well explained by micrometeorological variables (for rubber: Hardanto et al., 2017a; Niu et al., 2017; for oil palm: Röhl et al., 2015; Meijide et al., 2017), sometimes in combination with soil moisture fluctuations (e.g., Tromp-van Meerveld and McDonnell, 2006; Hardanto et al., 2017a). Over larger time-scales, changes in transpiration can also be driven by functional changes in communities. Spatial, site-to-site variability in transpiration is often much harder to explain. Studies encompassing multiple sites are rare, and often there is a lack of data on potential explanatory variables describing e.g. differences in stand structure,

productivity and soil. Available previous studies have related differences in transpiration among stands to diameter-dependent variables (e.g., basal area) as well as to specific leaf area (e.g., Cermák, 1989; Hatton et al., 1995; Roberts et al., 2001). In monoculture landscapes, plantation age was further found to substantially influence transpiration (Röhl et al., 2015; Niu et al., 2017).

From 1990–2011, the lowlands of Sumatra underwent substantial land-use changes, losing about 45% of their forest area. In the same period, rubber plantations increased by almost 30%, and both smallholder and commercial oil palm plantations more than doubled in area (Clough et al., 2016; BPS, 2013; Margono et al., 2012; Villamor et al., 2014). Land-use change in lowland Sumatra is largely driven by profitability, and today oil palm is often the preferred land-use option (Clough et al., 2016; Kubitzka et al., 2018). One main trajectory for the use of a certain unit of land leads from forest degradation over forest conversion to jungle rubber agroforest to monoculture rubber plantations to smallholder or commercial oil palm plantations (Clough et al., 2016; Drescher et al., 2016). It thus encompasses both land-cover change and land-use intensification as potential drivers of shifts in transpiration. Regarding the integrity of the hydrological cycle, this raises the question of how transpiration as a key ecohydrological process is affected along this linear main land-use change trajectory. At larger scales, current land-use dynamics in lowland Sumatra are mosaic-like and highly complex, further raising the question of how regional transpiration could be affected.

At 42 sites across the dominant land-use types of lowland Sumatra, Indonesia, we estimated transpiration using calibrated sap flux sampling schemes adapted to each land-use type. The land use types included selectively logged rainforests, jungle rubber agroforests, monoculture smallholder rubber plantations and smallholder and commercial oil palm plantations (Fig. 1). We (1) assessed annual stand-level transpiration and (2) analysed potential covariates to explore relationships with and discuss explanations for site-to-site variability in transpiration across land-use types. We (3) present a data-based concept on how land-use change along a trajectory forest-rubber-oil palm affects transpiration, and discuss what ecohydrological consequences this might have.

2. Materials and methods

2.1. Study region

The study region was Jambi Province (Sumatra, Indonesia, Fig. 1). From 1991–2011, annual temperature in Jambi was $26.7 \pm 0.2^\circ\text{C}$ (mean $\pm$ SD, data from Sultan Thaha airport, Jambi). Annual precipitation in Jambi was $2235 \pm 385\text{ mm}$. A relatively dry season (below 120 mm monthly precipitation) usually occurs between June and September (Drescher et al., 2016). Temperature and precipitation were similar in the vicinity of our study sites (at approx. 100 km distance, Figure S-1).

Soil types at the study sites were well-drained loamy Acrisols dominated by kaolinites. Soils in the Harapan landscape (Fig. 1) are chromic Acrisols (loamic) and in the Bukit Duabelas landscape chromic Acrisols (clayic). In the Bukit landscape, soils contained more clay (31–60%) and were more saturated with base cations (20–38%) than in the Harapan landscape (26–37% clay and 10–28% base saturation) (Allen et al., 2015; Guillaume et al., 2015). Our study sites contain no pedosequences, as the soil parent materials and relief were very similar among sites and landscapes.

Our study encompassed 42 sites distributed over dominant land-use types of lowland Sumatra (Fig. 1): selectively logged rainforest ($n = 8$ sites), which is equivalent to 'primary degraded forest' (Margono et al., 2014), 'jungle rubber' (Gouyon et al., 1993) agroforests ($n = 8$), smallholder rubber plantations ($n = 10$), smallholder oil palm plantations ($n = 14$) and commercial oil palm plantations ($n = 2$). They were located at similar altitude and landscape positions (non-valley,




Fig. 1. Study sites in the lowlands of Jambi Province, Sumatra, Indonesia. 32 core sites in forests, jungle rubber agroforests and medium-aged smallholder monoculture rubber and oil palm plantations. Eight additional age-class sites in smallholder rubber and oil palm plantations and two additional sites in commercial oil palm plantations. Eight micrometeorological stations in the study region and one long-term station in the greater region (Province map).

moderate slopes, 70 m asl). The 42 sites comprised 32 'core sites', for which a large data set of stand structural, productivity, microclimate and soil related variables (Table S-1) were available within the framework of the EForTS project (www.uni-goettingen.de/efforts; Drescher et al., 2016), as well as ten additional sites for which basic site and stand data (location, elevation, stand density, diameters, heights, age) had been recorded (Table S-2).

The eight core sites each in rubber and oil palm monocultures were 'productive' (i.e. fruit- or latex-yielding), medium-aged (8–16 years old) smallholder plantations (Drescher et al., 2016). Management among oil palm smallholder plantations was variable, but characterized by little fertilizer input (24–96 kg N ha⁻¹ yr⁻¹, Fan et al., 2015) compared to intensively managed, commercial oil palm plantations in the study region (240–456 kg N ha⁻¹ yr⁻¹, Fan et al., 2015; Teuscher et al., 2016). The ten additional sites encompassed two smallholder rubber and eight smallholder oil palm plantations along a gradient of plantation age as well as two commercial, intensively managed oil palm plantations (PTPN VI, PT Humusindo) (Fig. 1).

2.2. Sap flux derived daily transpiration

Sap flux measurements at 41 of 42 sites were performed between March 2013 and April 2014 (Table S-2, Table S-3). Measurements at the remaining site (PT Humusindo) were performed in October 2016, under similar environmental conditions and with identical methodology. At all 42 sites, we measured sap flux density (J_s , g cm⁻² h⁻¹) with thermal dissipation probes (TDP, Granier, 1985) in oil palm leaf petioles and in the outer xylem of tree trunks to derive daily stand transpiration (E_t , mm d⁻¹).

For oil palm, we used a species-specific set of parameters for the Granier sap flux equation (Granier, 1985) to estimate J_s ; it had been derived in previous laboratory calibration experiments on petioles with

simultaneous gravimetric measurements as a reference (Niu et al., 2015). We followed an oil-palm specific measurement scheme with simultaneous measurements in the petioles of 16 leaves on four different palms (Niu et al., 2015). At a mature oil palm site (PTPN VI), this approach yielded reasonable, i.e. 31–33% lower estimates for sap flux derived stand transpiration than for evapotranspiration derived from simultaneous eddy covariance measurements. The divergences are likely accounted for by canopy and soil evaporation, but could also partly root in estimation uncertainties associated with the sap flux measurements on a limited number of oil palms (Niu et al., 2015; Meijde et al., 2017) as well as in uncertainties associated with the eddy covariance technique at the PTPN VI site (e.g., regarding energy balance closure, Meijde et al., 2017) and in general (e.g., Teuling, 2018).

For rubber, we also applied a species-specific measurement scheme. It uses the original Granier equation parameters (Granier, 1985) for estimating J_s (confirmed by calibration experiments), with simultaneous sap flux measurements on six trees (with two sensors each) per site (Niu et al., 2017).

For the structurally more heterogeneous forest and jungle rubber stands, we confirmed the applicability of the original Granier equation parameters in exploratory laboratory experiments in analogy to previous work (Niu et al., 2017). On average, we found close agreement between TDP-derived and reference gravimetric measurements when using the original Granier equation parameters ($J_s = 0.94 \cdot J_G$; $R^2 = 0.92$; $P < 0.001$, where J_G is the gravimetrically-derived sap flux density in g cm⁻² h⁻¹). This is in line with results from previous calibration studies on diffuse-porous dicot trees (Lu et al., 2004). We adjusted the above-described measurement scheme for rubber trees to the more heterogeneous stand structure of forest and jungle rubber by measuring on eight instead of six trees per site (also with two sensors per tree).

Four out of 32 core sites were 'long-term' J_s monitoring sites (up to

1 year), one in each land-use type (Table S-4). At the remaining 28 core sites and at the ten additional sites, J_s was successively measured for reduced periods (on average 27 days, Table S-5). As sample trees, we chose predominant, dominant and co-dominant trees ('Kraft classes' 1–3, Kraft, 1984, see English description in Grala-Michalak and Kazmierczak, 2011). In all land-use types, the minimum diameter at breast height (DBH, cm) for sample trees was 10 cm. At jungle rubber sites, sensors were evenly distributed among rubber and non-rubber trees. In accordance with a well-accepted observation that tree water use scales universally with tree size across species (Meinzer et al., 2005), we did not take into account tree species in our sap flux design in the very species rich land-use types forest and jungle rubber. Instead, we evenly chose individuals of relatively larger, medium and smaller DBH to assure well-balanced samples with respect to tree dimension. Nonetheless, the 259 sap flux sample trees across the 42 study sites commonly belonged to the most dominant species within each stand (assessed by species rank of relative contribution to total stand basal area), even in the very species-rich land-use types forest and jungle rubber (Table S-6). In jungle rubber (24–56 tree species per site), more than 80% of the sap flux sample trees belonged to the 10 most dominant species at the according sites. In forest (51–97 tree species per site), about 70% of the sample trees belonged to the 20 most dominant species (Figure S-2).

Sensor installation and protection were carried out in analogy to previous work on trees and palms in the study region (Niu et al., 2015, 2017). The sensors were connected to AM16/32 multiplexers connected to CR1000 data loggers (Campbell Scientific Inc., Logan, USA); signals were recorded every 30 s and averaged and stored every 10 min. To scale-up from TDP point measurements of J_s to stand-scale E_t , stand water conductive areas (the summed-up tree cross sectional areas that are constituted by active xylem) were established for each site. For oil palm and rubber, methodological approaches were available (Niu et al., 2015; Niu et al., 2017). For forest and jungle rubber, in analogy to the approach for rubber (Niu et al., 2017), we measured radial patterns of J_s with increasing depth into the xylem (0–8 cm, 1 cm resolution) with heat field deformation sensors (HFD, ICT International, Armidale, Australia; Nadezhdina et al., 2012) on ten sample trees per land-use type, in parallel to TDP measurements in the outer xylem. By averaging daily J_s for each of the eight HFD measurement depths (from at least two weeks per tree), radial patterns of J_s were established. They were then normalized by the value at the standard TDP measurement depth and used to calculate the adjusted water conductive area (WCA_{adj} , cm²) for each sample tree, i.e. the diameter-dependent basal area corrected for radial differences in J_s in relation to the standard TDP measurement depth (see Niu et al., 2017 for methodological details). This approach yielded significant relationships between DBH and WCA_{adj} for both rubber and non-rubber trees at jungle rubber sites as well as for non-rubber trees at forest sites (Methods S-1). Using stand density and DBH distribution data of the study sites subsequently allowed calculating stand water conductive areas (WCA_{stand}).

To calculate daily E_t estimates for each site, J_s day-sums of all sensors running simultaneously were averaged and multiplied by the respective WCA_{stand} . Commonly, only days with more than 13 (oil palm), ten (rubber), and 14 (jungle rubber and forest) sensors running simultaneously were included due to high associated estimation errors at smaller sample sizes. Using these sampling schemes, E_t estimates in the heterogeneous land-use types forest and jungle rubber are associated with sample size-induced estimation errors of approx. 30% (Granier et al., 1996), while potential errors for rubber and oil palm are reported to be substantially smaller (< 15%, Niu et al., 2015, 2017). Days with severe data abnormalities or data gaps larger than two hours were excluded from the respective datasets; to fill data gaps up to two hours, linear interpolation of hourly values was performed before deriving daily sums.

2.3. Microclimate, soil moisture and reference potential evapotranspiration

We monitored key micrometeorological variables at eight open-area reference meteorological stations in the vicinity of the study sites (Fig. 1), as well as below-canopy microclimate and soil moisture and temperature with 32 micrometeorological stations at the center of each core site.

At the 32 below-canopy stations, below-canopy air temperature and relative humidity were measured. Simultaneously, soil moisture was monitored at 0.3 m depth below the soil surface (Meijide et al., 2018). To derive a complete data series of soil moisture for the study region (Figure S-3), we averaged data from all stations running simultaneously.

The eight meteorological reference stations near our study sites were located at similar altitude (60 m ± 15 m a.s.l.) in open terrain. Air temperature and relative humidity were measured at 2 m height with a thermohygrometer (type 1.1025.55.000, Thies Clima, Göttingen, Germany). Wind speed was measured with an anemometer (Thies Clima) at 4 m height. A net radiation sensor (NR Lite2, 200–100,000 nm, Kipp & Zonen, Delft, The Netherlands) and short-wave radiation sensor (CMP3 Pyranometer, Kipp & Zonen) were installed at 3 m. Measurements were taken every 15 s and averaged and stored every 10 min on a DL16 logger (Thies Clima). Recorded micrometeorological variables were the basis for deriving daily values of potential evapotranspiration (E_o , mm d^{−1}) with the Priestley-Taylor equation (Priestley and Taylor, 1972) using the R package Evapotranspiration (Guo and Westra, 2016). To obtain a complete (Jan 2013–Jan 2015) E_o series as a reference for sap flux derived E_t and as a basis for modelling annual values, we used an almost complete data series from one of our climate stations ('Reki station') and gap-filled missing values with the average of all other stations running simultaneously (distance 30–90 km, < 25% of daily data gap-filled). This seemed justified because daily E_o from all stations were highly and significantly correlated to E_o from Reki when running simultaneously (average divergences < 10%, Figure S-4). Annual E_o was 1408 mm in 2013 and 1310 mm in 2014.

More basic, but longer-term measurements of temperature and precipitation were further available from a station at Jambi Airport (1991–2014, approx. 100 km east of our sites) (Fig. 1). The station is operated by the Indonesian Weather Service BMKG.

2.4. Annual transpiration

Temporal variability of E_t is generally largely explained by day-to-day fluctuations in micrometeorological driving variables, given that soil moisture remains non-limiting (i.e. not too low and not too high) (Oren et al., 1999; O'Brien et al., 2004; Kume et al., 2007; Hardanto et al., 2017a). Our study period (Jan 2013–Jan 2015) at 41 of the 42 study sites ended before the 2015 ENSO event, while measurements at the remaining site (PT Humusindo) were performed well after the ENSO event, in October 2016. Environmental conditions during the respective study periods were representative of long-term average climatic conditions regarding temperature and precipitation (Figure S-1). Soil moisture remained relatively stable at sufficiently high levels over our study period and at all sites to be considered non-limiting for tree water use (Figure S-3, Table S-7). The sites were not flooded during the sap flux measurements.

We consequently followed a linear regression approach between daily E_t (sap flux derived) and potential evapotranspiration E_o derived from simultaneous micrometeorological measurements to estimate annual E_t . It is similar to previous cropping coefficient approaches for calculating crop evapotranspiration (Allen et al., 1998) and was previously applied for estimating annual E_t of two of the oil palm (Meijide et al., 2017) and one of the rubber plantations (Niu et al., 2017) in our study. In those two oil palm plantations, the approach yielded estimates for the contribution of E_t to total annual evapotranspiration (derived

from simultaneous eddy covariance measurements) of 7% (very young plantation) and 68% (mature commercial plantation) (Meijide et al., 2017). These results were interpreted as ecologically reasonable considering additional evaporation from soil and canopy (including leaf axils of mature palms, Tarigan et al., 2018), and, in the case of the very young plantation, the unassessed transpiration of a dense grass layer surrounding the planted, still very small oil palms (Meijide et al., 2017). Exploratory analyses with data from the four long-term monitoring plots showed that annual E_t derived from our simple linear regression approach differed less than 5% from several tested non-linear approaches (e.g. logarithmic, Weibull, Mitscherlich), with the advantage of higher robustness in case of study sites with relatively few available daily E_t values for model fitting.

At each site, we used the mentioned E_o time series for the study region (Figure S-3) and performed linear regressions (forced through origin) of daily TDP-derived E_t vs. E_o from the corresponding days. Resulting E_t/E_o ratios (i.e. the slopes of the regressions) were subsequently multiplied by annual E_o to obtain estimates of annual E_t (mm yr^{-1}). This approach of using linear E_t/E_o ratios to extrapolate to annual E_t rates accounts for potential differences in climatic driving variables (in our study represented by E_o) among the non-simultaneous sap flux measurement periods at the 42 sites. Exploratory Monte-Carlo analyses (with 10,000 iterations) using the data from the long-term monitoring plots (plot codes BF3, BJ5, BR3, BO3) and systematically varying the sampling size from 1 up to 365 consecutive days showed that our linear regression approach of deriving annual E_t estimates for the remaining 28 study sites based on reduced measurement periods (13–66 days of usable data, Table S-5) is associated with potential divergences of 15–20% compared to using fully measured data.

Our linear regression approach yielded strong linear relationships ($R^2 > 0.92$, $P < 0.001$) between daily E_t and E_o for all 42 study sites, based on 45 ± 11 days (mean $\pm$ SE) of simultaneous measurements per site (Table S-5). Comparing day-to-day measured E_t with E_o -based modeled E_t series shows good agreement across sites and land-use types (Figures S-5 and S-6). Divergences between measured and modelled mean daily E_t were consistently below 5% and RMSEs mostly below 20% (max. 28%). Parametric bootstrapping suggests low uncertainties (expressed by SEs) of annual E_t estimates, less than 5.2% at all sites (Table S-5).

A study from the same region reported about 20% reductions in monoculture rubber E_t due to gradual leaf coverage reductions of 30–60%, with the partial leaf shedding period lasting approx. three months (Niu et al., 2017). Because of this, we firstly corrected all annual monoculture rubber E_t estimates that base on measurements under fully-leaved conditions (five core sites and two additional sites) by a shedding period of 86 days, with E_t reductions of 20% (in analogy to observations at the rubber long-term monitoring site BR3, also see Figure S-3). Secondly, for two monoculture rubber sites among the 32 core sites (plot codes HR2 and HR3, Table S-3) the E_t/E_o ratios were linearly corrected before extrapolation to annual E_t , because partial leaf shedding occurred during the entire respective sap flux measurement periods. Based on the study by Niu et al. (2017), we multiplied the two respective E_t/E_o ratios by 1.25 (i.e. accounting for on average 20% lower E_t during partial leaf shedding, Table S-5). Assuming about 20–40% shedding at our sites (visual assessment), this magnitude of correction is in line with research on several tropical tree species showing a similar (and linear) relationship between degree of shedding and E_t (Kunert et al., 2010).

2.5. Covariates to explore site-to-site variability in transpiration

We hypothesized that site-to-site variability in stand transpiration E_t (or mean tree water use WU) is explained by stand structural, productivity and soil variables. For our 32 core sites, a large set of covariates were available to test this hypothesis. General site and stand characteristics had previously been published. They include mean stand

diameter and height, above- and belowground biomass (Kotowska et al., 2015) and estimates of canopy gap fraction (Drescher et al., 2016) (Table S-3). Further available covariates were estimates of above- and belowground net primary productivity (Kotowska et al., 2015) as well as a variety of classic pedological parameters (e.g., pH, bulk density, nitrogen- and carbon-related variables; Allen et al., 2015; Guillaume et al., 2015) and annual means of soil moisture and temperature (Clough et al., 2016). Furthermore, basic microclimatic variables (e.g., average annual below-canopy temperature) were available from each of the 32 sites (Clough et al., 2016; Meijide et al., 2018).

As additional, previously unpublished stand structural variables, estimates of canopy cover and different effective leaf area indices (LAI , $\text{m}^2 \text{m}^{-2}$) were calculated from hemispherical photographs (Drescher et al., 2016) with CAN-EYE (Version 6.4.6, INRA Centre de recherche PACA, Avignon, France). Simultaneously to taking hemispherical photographs, below-canopy photosynthetically active radiation (PAR , $\mu\text{mol s}^{-1} \text{m}^{-2}$) and PAR at an open reference site were measured with LI-190SA quantum sensors connected to LI-Cor 250 A light meters and LI-1400 data loggers (Li-Cor Biosciences, Lincoln, NE, USA) to estimate canopy light transmission (%).

Based on nine systematically distributed single terrestrial laser scans made at 29 of the 32 core sites (notably: in 2016; three sites missing), a stand structural complexity index ($SSCI$, Ehbrecht et al., 2017) and the effective number of canopy layers (ENL , Ehbrecht et al., 2016) were calculated. The computations were based on 3D points clouds obtained using a FARO Focus 3D terrestrial laser scanner (Faro Technologies Inc., Lake Mary, USA).

Soil samples taken from different horizons were analysed regarding their carbon (C) and nitrogen (N) content, their C/N ratio and the abundance of stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) (Guillaume et al., 2015).

2.6. Shifts in local and regional transpiration

One main land-use trajectory for a certain unit of land in lowland Sumatra leads from forest degradation over forest conversion to jungle rubber to monoculture rubber plantations to smallholder or commercial oil palm plantations (Clough et al., 2016; Drescher et al., 2016). To examine shifts in transpiration along this linear trajectory, we applied the annual E_t estimates from the 42 study sites. For forest and jungle rubber agroforests (non-rotational systems), we assumed steady-state E_t over time. For the rubber and oil palm monoculture plantations (rotation systems), we provide time-averaged E_t values over rotation cycles of 20 (rubber) and 25 years (oil palm); they base on age- E_t relationships as derived from the analysis of 14 smallholder oil palm and 10 smallholder rubber plantations along a gradient of plantation age (including the eight core sites each, see *Statistical analyses*). We further expanded our measurement-based E_t estimates by literature values from previous studies in undisturbed tropical lowland forests (average of 12 sites in Kunert et al., 2017; Kume et al., 2011; Bruijnzeel, 1990; Lion et al., 2017).

We combined stand-level E_t in dominant land-use types (as presented in the linear trajectory) with area fraction estimates for 1990 and 2011 (Clough et al., 2016) to estimate regional shifts in transpiration for lowland Sumatra. We further sub-categorized forests into 'undisturbed' and 'selectively logged' forests using data from Margono et al. (2012). The category oil palm was further subdivided into 'smallholder' and 'commercial' plantations, using data from the Statistical Office of Indonesia (BPS, 2013). We applied the value of jungle rubber agroforests for all 'intermediate' land-use types (e.g., fallows, regrowth). Due to a lack of any further information, for our calculations all thus far not classified 'other areas' were assigned the transpiration value of the average of all other land-use types.

2.7. Statistical analyses

To ensure the stability of our linear regression approach between

daily transpiration E_t (sap flux derived) and potential evapotranspiration E_o (Priestly-Taylor-derived) for estimating annual transpiration at our study sites, and to provide uncertainty estimates for annual E_t , we performed parametric bootstrapping (with the R package 'boot', Canty and Ripley, 2017) with 50,000 iterations and derived means and standard errors. In analogy to previous work (Meijide et al., 2017), we bootstrapped the linear relationship between E_t and E_o (forced through the origin) (Table S-5).

To test for differences in mean annual E_t (and WU) among land-use types and in the forest-rubber-oil palm land-use change trajectory, we applied linear regressions with land-use type as the categorical variable. We tested for overall significance with ANOVA ($P < 0.05$) and for significant differences among groups with a post-hoc Tukey-LSD test. The data sets were log-transformed for the ANOVA to minimize patterns in residuals.

To identify and analyse potential covariates for explaining site-to-site variability in transpiration, we first applied univariate linear regression models and tested all available single variables. We further applied multiple linear regression models ($E_t = a + b_1 \cdot x_1 + b_2 \cdot x_2 + \dots + b_n \cdot x_n$, where a is the intercept, x are the different available variables and b are the parameters to be fitted) and tested different blocks of variables from the categories stand structure, productivity and soil (with and without statistical interaction) in a 'brute-force' approach using the R package leaps (Lumley and Miller, 2009). In our study, only significant ($P < 0.05$) models of substantial explanatory power ($R^2 > 0.60$) are presented. They are ordered from lower to higher Akaike information criterion (AIC) values and provided with F-statistics and degrees of freedom (DF). Models were further examined regarding patterns in residuals, leverage and robustness.

To examine the influence of plantation age on E_t , we separately pooled the data from all available smallholder rubber ($n = 10$ sites) and smallholder oil palm plantations ($n = 14$), respectively. We normalized E_t values by the respective means of the eight core sites each and plotted them against plantation age. We then fitted exponential functions ($y = y_0 + A \cdot \exp(R_0 \cdot x)$, where y_0 , A and R_0 are the parameters to be estimated) to each of the two datasets.

Statistical analyses were performed with R 3.1.1 (R Core Team, 2014). For graphing, Origin 8.5 (Origin Lab, Northampton, MA, USA) was used.

3. Results

3.1. Transpiration among land-use types

Annual transpiration (E_t , mm yr^{-1}) for the 32 core sites decreased in the sequence selectively logged forest ($589 \pm 52 \text{ mm yr}^{-1}$, mean $\pm$ SE), medium-aged smallholder oil palm plantations ($543 \pm 24 \text{ mm yr}^{-1}$), jungle rubber agroforests ($521 \pm 80 \text{ mm yr}^{-1}$) and medium-aged smallholder monoculture rubber plantations ($391 \pm 38 \text{ mm yr}^{-1}$) (Fig. 2). Mean E_t differed significantly among land-use types (ANOVA, $P < 0.05$). E_t in rubber monoculture was lower than in forest and oil palm (post-hoc LSD test, $P < 0.05$), while jungle rubber E_t was intermediate. Jungle rubber was characterized by at least 37% higher site-to-site variability in E_t than the other land-use types and encompassed the sites with the highest ($995 \pm 43 \text{ mm yr}^{-1}$, mean $\pm$ SE from bootstrapping) and lowest E_t ($264 \pm 14 \text{ mm yr}^{-1}$) (Fig. 2, Table S-3).

3.2. Site-to-site variability in transpiration

Only small parts of the observed variability in E_t among the 32 core sites (Fig. 2) were explained using univariate regressions and the large set of potential covariates (Table S-8). A multiple linear regression model allowing for interactions among stand structural, productivity and soil variables explained 73% of variability in E_t . Stand density, mean tree diameter and topsoil stable $\delta^{15}\text{N}$ isotopic signature were




Fig. 2. Stand-level transpiration across land-use types. Annual transpiration (E_t , mm yr^{-1}) and the ratio of E_t to potential evapotranspiration (E_t / E_o) of eight core sites each in forest (F), jungle rubber agroforests (J) and medium-aged smallholder rubber (R) and oil palm plantations (O). Overall means (and standard errors) are presented as large dots (with bars). Significant differences between the groups as based on an ANOVA with post-hoc LSD test are indicated by different letters.

significant covariates ($P < 0.001$, Table 1).

We further analysed potential covariates of mean tree water use WU across the 32 sites (Table S-9). Crown projection area per tree explained 91% of the variability in WU ($P < 0.001$). Several further stand structural variables (e.g. mean tree diameter, stand density, fine root biomass and gap fraction) also explained much of the observed variability in WU ($R^2 > 0.6$, $P < 0.001$). WU was further positively related to productivity-related variables, particularly per-tree total and aboveground net primary productivity (NPP) ($R^2 > 0.85$, $P < 0.001$) (Table 1).

3.3. Influences of plantation age and management

For both rubber and oil palm smallholder plantations (combined data from core sites and additional sites, Fig. 1), we observed strong increases in annual E_t up to an age of about seven years. E_t subsequently maintained a steady state regardless of plantation age (Fig. 3). Of site-to-site variability in E_t , 51% (rubber, $n = 10$) and 80% (oil palm, $n = 14$) were explained by plantation age ($P < 0.001$). In intensively managed, commercial oil palm plantations (PTPN VI and PT Humusindo), E_t was on average 73% higher than in smallholder plantations of similar age (Fig. 3).

3.4. Shifts in transpiration at the local and regional scale

The previously introduced linear main land-use trajectory for a certain unit of land in lowland Sumatra (forest-rubber-oil palm) goes along first with a decrease and then a rebound of E_t (Fig. 4). Conversion of selectively logged forests to monoculture rubber plantations (land-cover change) decreases E_t by $43 \pm 11\%$. Eventual land-cover change to oil palm leads to a rebound in transpiration: conversion to smallholder oil palm almost raises E_t back to the level of forests, while land-use intensification to commercial oil palm increases E_t by a factor of 1.7 compared to smallholder oil palm, thus even exceeding E_t in logged forests (Fig. 4).

At the regional scale, land-use dynamics are more complex than depicted in the 'local' linear main trajectory. Combining stand-level E_t of dominant land-use types with area fraction estimates for lowland Sumatra suggests 9% decreases in regional E_t from 1990 to 2011 (Table 2). The predicted continuing conversion to oil palm and particularly land-use intensification to commercial oil palm will likely

Table 1

Covariates to explore site-to-site variability in stand-level transpiration (E_t) and tree-level water use (WU). Results from linear regression analyses among the 32 core sites based on variables related to stand structure, productivity and soil characteristics.

	Stand structure	Covariates	Soil	AIC	R ²	Statistics	F-stat	DF
		Productivity				P		
Stand transpiration E_t	stand density * diameter _{mean} ³	–	* ¹⁵ N signature ³	–55.5	0.70	0.0002	7.1	7, 21
Mean tree water use WU	per-tree crown area	–	–	251.3	0.91	2.20E-16	321.4	1, 30
	diameter _{mean} ¹	–	–	256.7	0.90	2.20E-16	266.7	1, 30
	–	per-tree NPP _{total} ¹	–	261.1	0.88	1.43E-15	228.5	1, 30
	–	per-tree NPP _{aboveground} ¹	–	263.2	0.87	3.81E-15	212.2	1, 30
	per-tree necromass _{fineroot} ¹	–	–	283.1	0.76	4.46E-11	100.2	1, 30
	mean tree height ²	–	–	292.0	0.69	3.00E-09	68.7	1, 30
	canopy cover ²	–	–	292.1	0.68	3.13E-09	68.4	1, 30
	stand density ²	–	–	293.2	0.67	5.34E-09	65.0	1, 30
	per-tree biomass _{fineroot} ¹	–	–	296.9	0.63	3.19E-08	54.5	1, 30

– = not significant.

* = multivariate regression with interaction.

¹ positive influence.

² negative influence.

³ interaction.

rebound regional E_t . In comparison to times when the forests of Sumatra were still more extensive and undisturbed as a baseline (e.g. 1932), regional reductions in E_t until 2011 are estimated to be much larger (Table S-10).

4. Discussion

4.1. Transpiration among land-use types

Annual transpiration at our 32 core sites decreased in the sequence selectively logged forest, medium-aged smallholder oil palm plantations, jungle rubber agroforests and medium-aged smallholder monoculture rubber plantations (Fig. 2).

Mean E_t of our forest sites ($589 \pm 52 \text{ mm yr}^{-1}$) was considerably lower than in undisturbed, old growth tropical lowland rainforests in Amazonia and South East Asia (1023 mm yr^{-1} , mean of 12 sites, Bruijnzeel, 1990; Kume et al., 2011; Kunert et al., 2017; Lion et al., 2017). Among these, Pasoh Forest Reserve in Malaysia is located closest to our study region; there, E_t was estimated at 801 mm yr^{-1} (Lion et al., 2017). In Amazonian undisturbed forest, it was found that large emergent canopy trees account for up to 71% of E_t (Kunert et al., 2017). Such trees were largely absent in the selectively logged forests (Clough et al., 2016; Drescher et al., 2016) where our study sites were located. It thus seems reasonable that we found lower E_t than the mean of previous studies in undisturbed tropical forests.

Jungle rubber was characterized by high site-to-site variability in E_t and encompassed both the sites with the highest and lowest E_t . Mean

jungle rubber E_t ($521 \pm 80 \text{ mm yr}^{-1}$, or on average 1.4 mm day^{-1}) appears centred within a variety of tropical agroforestry and reforestation system estimates from previous studies ($0.5\text{--}2.5 \text{ mm day}^{-1}$, Cienciala et al., 2000; Dierick and Hölscher, 2009; Dierick et al., 2010; Köhler et al., 2014; Hardanto et al., 2017b).

Medium-aged smallholder rubber monoculture plantations had the lowest mean E_t ($391 \pm 38 \text{ mm yr}^{-1}$), significantly lower than forests and smallholder oil palm (Fig. 2). The lower transpiration in rubber plantations is in line with 33–53% lower NPP (Kotowska et al., 2015), and with 20–80% lower basal area, biomass and leaf area compared to forests (Table S-3). Periodically occurring partial leaf shedding during dry periods with reductions of E_t of 30–70% contributes to low annual rubber E_t (Niu et al., 2017). At first sight, our low rubber E_t estimate contrasts reports of high rubber evapotranspiration from mainland Asia (Tan et al., 2011). However, available estimates of annual E_t from mainland Asia are only 10–20% higher than our mean rubber estimate (Kobayashi et al., 2014; Giambelluca et al., 2016). Potential reasons for divergences are 11–14% higher stand densities and higher management intensity at the mainland sites as well as higher evaporative demand in mainland Asia compared to equatorial Indonesia (Niu et al., 2017).

Mean E_t in medium-aged smallholder oil palm plantations ($543 \pm 24 \text{ mm yr}^{-1}$) was only 8% lower than in forest (Fig. 2), despite 85% lower total biomass (Kotowska et al., 2015). Annual evapotranspiration of oil palm plantations derived from eddy covariance measurements was also found to be high, with rates similar to tropical forests (Meijide et al., 2017; Manoli et al., 2018). Mean per-tree water use WU of oil palms was substantial ($108 \pm 6 \text{ kg d}^{-1}$, mean $\pm$ SE




Fig. 3. Transpiration as influenced by plantation age in rubber and oil palm plantations. Annual transpiration (E_t) normalized by previously presented means of eight core sites each in rubber and oil palm plantations (Fig. 2), plotted against plantation age. Exponential functions were fitted to each dataset (excluding the commercial oil palm plantations PTPN VI and PT Humusindo).




Fig. 4. Data-based concept of annual transpiration (E_t) changes along a forest-rubber-oil palm trajectory. Equivalent changes in the ratio of E_t to potential evapotranspiration (E_t / E_o). Values for rubber and oil palm monoculture systems are age-adjusted averages over rotation cycles, while 'steady state' transpiration over time is assumed for forests and agroforests. We used mean E_t of 12 undisturbed tropical lowland rainforests across the tropics (Table 2) as the starting point of the trajectory. Bars indicate bootstrapping-derived standard errors. Significant differences in E_t among land-use types (ANOVA with post-hoc LSD-test) are indicated by different letters.

among sites) in our study, four- to five-times higher than in the other land-use types (Table S-3). High E_t and WU in oil palm plantations go along with very high NPP (47% higher than at forest sites, Guillaume et al., 2018). Oil palm NPP is to 58% allocated to fruit yield (Guillaume et al., 2018), making it the globally highest-yielding oil crop.

4.2. Site-to-site variability in transpiration

Site-to-site transpiration varied considerably across the 32 core sites (Fig. 2). Using univariate regressions, only small parts of the observed variability in E_t were explained (Table S-8). Basic stand structural variables, more complex indices derived from terrestrial laser scans (SSCI, ENL) as well as soil moisture alone did not explain site-to-site variability in E_t . While it is undisputed that spatial variations in soil moisture in general have significant control over E_t (Berry et al., 2005; Tromp-van Meerveld and McDonnell, 2006; Hardanto et al., 2017a), soil moisture did not significantly influence E_t in our study because of high precipitation (Merten et al., 2016) and loam and clay Acrisol soil properties (Allen et al., 2015) at the study sites. Soil moisture fluctuated little among days and seasons and remained relatively high on average and at all measurement stations in the study region throughout the study period (Table S-7) due to the absence of prolonged periods without precipitation (Figure S-3). Exploratory analyses showed no significant influence of soil moisture fluctuations on E_t within or across land-use types at any time scale (e.g. hourly, daily, monthly), which is

in line with previous results from rubber and oil palm long-term monitoring sites in the study region (Niu et al., 2015, 2017).

A multiple linear regression model allowing for interactions among stand structural, productivity and soil variables explained 73% of variability in E_t , with the variables stand density, mean tree diameter and topsoil stable $\delta^{15}N$ isotopic signature as significant covariates (Table 1). Stand density and mean tree diameter (or basal area) are usual suspects for explaining variability in E_t (e.g. Cermák, 1989; Hatton et al., 1995; Cienciala et al., 2000); as such, with site-specific upper limits, E_t tends to be higher in stands with a higher density and a larger mean diameter. Soil $\delta^{15}N$ has, to our knowledge, not been connected to E_t in previous studies. Its ecological interpretation is thus more difficult; in our case, it likely functions as an indicator of disturbance. As such, higher soil $\delta^{15}N$ have been related to lower canopy cover (Evans and Ehleringer, 1993) and a higher degree of soil disturbance (Matson et al., 1987) and erosion (Martinelli et al., 1999, by bringing up deeper soil layers with a higher $\delta^{15}N$ signature). It can potentially also be related to fertilization. However, further in-depth studies will be needed to comprehensively address the role of soil $\delta^{15}N$ as an indicator for ecohydrological processes.

We further expected a productivity-related variable to be part of significant (multiple) linear E_t models. However, due to the exceptionally high water use efficiency of oil palm (NPP divided by E_t), these models do not hold across land-use types. For smallholder oil palm plantations, mean water use efficiency was 44% higher (3.1 g C

Table 2

Land-use types, area fractions and shifts in regional transpiration between 1990 and 2011. Ratios of transpiration to potential evapotranspiration (E_t / E_o) and transpiration (E_t , mm yr⁻¹) as presented in our study for selectively logged forests, jungle rubber agroforests and monoculture rubber and oil palm plantations in lowland Sumatra. Time-averaged E_t over the respective common rotation cycles are presented for rubber and oil palm monoculture systems.

Annual precipitation (P) 2199 mm Annual potential evapotranspiration (E_o): 1408 mm P/ E_o ratio: 1.56	E_t / E_o Ratio Time-averaged means	E_t (mm yr ⁻¹) Measurements in our study (mean $\pm$ SE)	1990 area fraction (%) Adjusted from Clough et al., 2016	2011 area fraction (%) Adjusted from Clough et al., 2016
Undisturbed forest	0.73 ^{*1}	1023 ^{*1}	17.9 ^{*3}	9.5 ^{*3}
Logged forest	0.42	589 $\pm$ 52	48.6 ^{*3}	27.1 ^{*3}
Intermediate land-use, jungle rubber	0.37	521 $\pm$ 80	3.4	16.4
Small-holder rubber	0.24	336 $\pm$ 33	20.4	26.2
Small-holder oil palm	0.34	477 $\pm$ 22	4.7 ^{*4}	11.6 ^{*4}
Commercial oil palm	0.59	827 $\pm$ 77	2.8 ^{*4}	7.2 ^{*4}
Other	0.45 ^{*2}	629 ^{*2}	2.2	2.0
Relative E_t differences between 1990 and 2011	–	–	–	–9 %

^{*1} Average E_t values reported for 12 undisturbed tropical lowland rainforests across the tropics in Kunert et al., 2017; Kume et al., 2011; Lion et al., 2017 and Bruijnzeel, 1990.

^{*2} Due to the lack of any further information on area type, 'other areas' were assigned the E_t and E_r values of the average of all other land-use types.

^{*3} Data on extent of undisturbed vs. logged forests from Margono et al., 2012.

^{*4} Data on extent of commercial vs. small-holder oil palm cultivation from BPS, 2013 (same ratio assumed for 1990 and 2011).

kg⁻¹ H₂O) than for forest, jungle rubber and rubber systems.

In our study, tree-level measurements were used to estimate stand-level E_t . We thus extended our analyses to covariates of mean tree water use WU (Table S-9). The best predictor for WU was the stand structural variable crown projection area per tree ($R^2 > 0.9$), but several further stand structural and productivity-related variables also explained much of the observed variability in WU ($R^2 > 0.6$, Table 1). In accordance with the results of our study, close relationships between water use and crown or leaf area related variables, often in contrast to classic stem size related variables such as diameter, have been described in a variety of previous studies (Cermák, 1989; Hatton et al., 1995; Medhurst and Beadle, 2002; Radersma et al., 2006; Alcorn et al., 2013). However, previous studies had to our knowledge never confirmed this across such a large number of study sites and across a land-use trajectory from forest to perennial agricultural monocultures.

So far having discussed the variability across land-use types ($n = 32$) finally raises the question whether the observed within land-use type variability of E_t (Fig. 2) can be explained. Among the eight forest sites, we found a strong trend of decreasing E_t with increasing $\delta^{15}\text{N}$ in the topsoil ($R^2 = 0.86$, $P < 0.001$), which can be interpreted as reduced transpiration with increasing disturbance regime. In jungle rubber stands, E_t rose with increasing water conductive area of the stand (WCA_{stand} , $R^2 = 0.52$, $P < 0.05$), which in turn increases with increasing stand density and mean tree diameter. The available datasets further allow for a speculation on the reasons behind very high or low E_t values at single sites. For example, the by far highest E_t at one single jungle rubber site ('BJ6') was particularly due to very high sap flux density (J_s). The stand is characterized by fewer but taller and thicker trees and higher structural complexity (laser-scanning based $SSCI$) and leaf area (PAI) than at any of the other jungle rubber sites (Table S-3). Tree species composition (Table S-6) could be a further potential reason for the high observed J_s ; as such, leaf stomatal conductance and sap flux velocities were found to be much higher in tropical early- than in late-successional tree species (Jührbandt et al., 2004; Bretfeld et al., 2018). For the monoculture rubber and oil palm plantations, we further found strong effects of age and land-use intensity, which we address in the following section.

The observed strong divergences in covariate performance between the tree- and the stand-level could potentially be due to several reasons. (1) The experimental design of the study was not optimized towards explaining site-to-site variability, but rather towards providing sound mean values for each land-use type. A gradient study design rather than our replicated design likely would have been more successful in identifying underlying ecological drivers of stand-scale E_t (Kreyling et al., 2018). (2) Both our E_t estimate and all potential covariates are associated with estimation uncertainties that may mask underlying relationships at the stand-scale. For assessing E_t at a single site, uncertainties that arise from limited sap flux sampling size may be as high as 30% in the more heterogeneous land-use types forest and jungle rubber (Granier et al., 1996), but likely lower than 15% in the rubber and oil palm monocultures (Niu et al., 2015, 2017). As our study encompassed several replicates per land-use type, uncertainties associated with the overall means are likely substantially lower. However, future studies could benefit from more within-site replicates in heterogeneous land-use types. (3) There might be scaling problems from trees to ecosystem. For scaling our tree level sap flux measurements to stand-level E_t , our measurement scheme relies on the key structural variable stand water conductive area. It mainly depends on stand density and mean tree diameter, both of which then form part of the significant multiple linear model. Independent ecosystem-scale measurements of E_t , for example via remote sensing, could help to circumvent this problem in future studies. (4) There may be further, unassessed variables such as topographic landscape position or neighborhood effects potentially influencing E_t . Still, we found that a high degree of variability in tree-level WU was explained by tree structure, and stand-level E_t was largely explained by stand structure and topsoil $\delta^{15}\text{N}$. This offers

opportunities for a better understanding of the controls of E_t in rapidly changing tropical landscapes.

4.3. Influences of plantation age and management

Site-to-site variability in E_t was to 51% (rubber) and 80% (oil palm) explained by plantation age. Over the commonly observed rotation cycles (rubber: 20 years, oil palm: 25 years), E_t was characterized by a rapid increase up to about seven years plantation age (approx. corresponding to canopy closure) and a subsequent steady state for both rubber and oil palm (Fig. 3). Time-averaged E_t over the mentioned rotation cycles was 16% (rubber) and 12% (oil palm) lower (Table 2) than the means at the respective medium-aged core sites (Fig. 2). We found no indications of declining E_t at advanced plantation age for both rubber and oil palm, likely because of the short rotation cycles in these agricultural systems. This stands in contrast to previous studies in even-aged forest stands, which cover much larger time scales per rotation cycle and commonly show, after a relatively early peak, lower E_t with increasing stand age (Jayasuriya et al., 1993; Roberts et al., 2001; Vertessy et al., 2001; Delzon and Loustau, 2005).

In lowland Sumatra, rubber is predominantly grown by smallholders, with no or very little fertilizer input. In contrast, oil palm is grown by smallholders and in commercial, more intensively managed plantations. In two commercial oil palm plantations, E_t was on average 73% higher than in smallholder plantations and 40% higher than at our forest sites (Fig. 3, Table 2). Commercial oil palm E_t was 9% lower than mean E_t in previously studied undisturbed tropical rainforests (Bruijnzeel, 1990; Kume et al., 2011; Kunert et al., 2017; Lion et al., 2017) (Table 2), which is substantial considering e.g. the much lower biomass or leaf area. The substantial sap flux-derived E_t at the commercial oil palm sites is in line with equally substantial evapotranspiration estimates derived from simultaneous eddy covariance measurements at one of the sites (PTPN VI) (Meijide et al., 2017). Reasons for much higher E_t in commercial than in smallholder oil palm plantations are likely related to management intensity. Among our study sites, commercial oil palm plantations received five- to ten-times higher nitrogen fertilization than smallholder plantations (Fan et al., 2015; Teuscher et al., 2016). Although single oil palm smallholder plantations in Indonesia and Jambi, respectively, have been reported to receive much larger amounts of N-fertilizer (up to 170 kg N ha⁻¹ yr⁻¹, Woittiez et al., 2018) than our smallholder study sites (24–96 kg N ha⁻¹ yr⁻¹, Fan et al., 2015), averages (141 kg N ha⁻¹ yr⁻¹, Van Noordwijk et al., 2017) remain substantially below the inputs at our two commercial sites (240–456 kg N ha⁻¹ yr⁻¹, Fan et al., 2015; Teuscher et al., 2016). Commercial oil palm cultivation also involves more intensive herbicide and pesticide application (Teuscher et al., 2016), as well as optimized harvesting practises and more quality control in the selection of hybrid seedlings used as planting material (Woittiez et al., 2018). Such advanced and more intensive plantation management is associated with substantial increases in oil palm fruit yield (Fan et al., 2015), which is also reflected in higher E_t : Among oil palm sites for which approximate data were available (core and commercial sites), we found a linear trend of higher E_t at higher fruit yield ($n = 10$, $R^2 = 0.34$, $P < 0.05$).

Even though we only had two study sites in commercial oil palm plantations (as opposed to a total of 14 smallholder sites) and the results will thus have to be strengthened further by systematic experimental trials, we found sound indications that agricultural land-use intensification from smallholder to commercial oil palm is accompanied by a substantial increase in E_t . To our knowledge, no studies examining in-situ E_t from perennial tropical agricultural crops under different management intensity regimes are available thus far for comparison. However, our results are in line with recent research proposing that increases in productivity of annual crops, or management intensification, increase (evapo)transpiration (Mueller et al., 2016, 2017).

4.4. Shifts in transpiration at the local and regional scale

In lowland Sumatra, land-use change is driven by profitability (Clough et al., 2016). In recent years, rubber commodity prices have declined. Together with other socio-economic drivers, this has made oil palm a more attractive land-use option compared to rubber (Kubitza et al., 2018). One main land-use trajectory for a certain unit of land leads from forest degradation over forest conversion to jungle rubber to monoculture rubber plantations to smallholder or commercial oil palm plantations (Clough et al., 2016; Drescher et al., 2016). This linear trajectory results in a U-shaped shift in transpiration, with relatively high E_t in oil palm, particularly in intensively managed commercial plantations (Fig. 4).

Land cover and land use are quite heterogeneous in lowland Sumatra and have changed considerably over time (Clough et al., 2016; Drescher et al., 2016). Oil palm is often more commonly cultivated near larger populated areas with established infrastructure, while rubber often remains the preferred land-use option in remote areas (Euler et al., 2016). In between, some protected forest remnants and other land-use types can be found (Clough et al., 2016; Drescher et al., 2016). At the regional scale, land-use dynamics are thus more complex than depicted in the linear main trajectory for a certain unit of land. Combining stand-level E_t estimates of dominant land-use types from our study with adjusted area fraction estimates for lowland Sumatra (Margono et al., 2012; BPS, 2013; Clough et al., 2016) suggests 9% decreases in regional E_t from 1990 to 2011 (Table 2). Decreases of E_t until 2011 are much more substantial (43%) when taking a time when the forest cover of Sumatra was still largely intact as a baseline for comparison; on the other hand, the predicted continuing conversion to oil palm and particularly land-use intensification to commercial oil palm will likely result in a rebound of regional E_t (Table S-10). The latter is in line with recent studies suggesting regionally increasing (evapo)transpiration as a result of cropland expansion and intensification (Betts et al., 2013; Mueller et al., 2016, 2017).

Land-cover change to oil palm and particularly land-use intensification to commercial oil palm plantations can enhance periodical local water scarcity for humans during dry periods (consecutive weeks without precipitation), as has been indicated in interviews with rural villagers (Merten et al., 2016). Soil structure degradation and associated erosion after forest conversion strongly reduce soil water infiltration and increase surface run-off, more so in oil palm plantations than in rubber plantations (Guillaume et al., 2015, 2016; Tarigan et al., 2018). Further, very high canopy interception of precipitation has been reported for oil palm plantations (higher than in forests, Tarigan et al., 2018). In oil palm dominated landscapes, high E_t as reported in our study can lead to additional reductions of available water. In contrast, rubber cultivation seems less concerning in terms of water scarcity for humans due to 30–60% lower E_t (our study), 39% lower canopy interception (Tarigan et al., 2018) and less severe soil degradation (Guillaume et al., 2016).

5. Conclusions

We conclude that land-cover change and land-use intensification substantially alter transpiration in lowland Sumatra. Site-to-site variability in E_t is largely explained by stand structural and soil variables and plantation age. Transpiration along a land-use trajectory forest-rubber-oil palm first substantially decreases and then rebounds. We uncovered that particularly commercial, intensive oil palm cultivation leads to very high transpiration. As commercial oil palm cultivation is on the rise in many tropical regions, increases in transpiration due to land-cover change and to intensification in oil palm plantations will become more widespread in the future.

Declarations of interest

None.

Acknowledgements

This study was supported by a grant from the German Research Foundation (DFG, CRC 990 'EForTS' project: sub-projects A02, A03, A04, B04, B06, B07, B11, Z02, ABS). Afik Hardanto received a scholarship from the Indonesian-German Scholarship Programme (IGSP) and Furong Niu from the China Scholarship Council (CSC). The sap flux measurements as the core of the here presented work were carried out under the research permit number 15/EXT/SIP/FRP/SM/II/2013, as issued by the Indonesian Institute of Sciences LIPI. We thank Heri Junaedi, Agusta Herdhata and Andrea Hanf for field support and co-operation, our technical staff Edgar Tunsch, Malte Puhon and our field assistants Ali, Bayu, Ferry, and Basri for field support. We thank Christian Stiegler for providing micrometeorological data for October 2016. Thanks to all 'EForTS' colleagues and friends in Indonesia, Germany, and around the world. Terimah kasih!

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.agrformet.2019.04.017>.

References

- Abood, S.A., Lee, J.S.H., Burivalova, Z., Garcia-Ulloa, J., Koh, L.P., 2015. Relative contributions of the logging, Fiber, oil Palm, and mining industries to forest loss in Indonesia. *Conserv. Lett.* 8 (1), 58–67. <https://doi.org/10.1111/conl.12103>.
- Alcorn, P.J., Forrester, D.I., Thomas, D.S., James, R., Smith, R.G.B., Nicotra, A.B., Bauhus, J., 2013. Changes in whole-tree water use following live-crown pruning in young plantation-grown *Eucalyptus pilularis* and *Eucalyptus cloeziana*. *Forests* 4 (1), 106–121. <https://doi.org/10.3390/f4010106>.
- Alkama, R., Cescatti, A., 2016. Biophysical climate impacts of recent changes in global forest cover. *Science* 351 (6273), 600–604. <https://doi.org/10.1126/science.123325>.
- Allen, R.G., Pereira, L.S., Raes, D., Smith, M., W, a B., 1998. Crop evapotranspiration - Guidelines for computing crop water requirements - FAO Irrigation and drainage paper 56. *Irrig. Drain.* 1–15. <https://doi.org/10.1016/j.eja.2010.12.001>.
- Allen, K., Corre, M.D., Tjoa, A., Veldkamp, E., 2015. Soil nitrogen-cycling responses to conversion of lowland forests to oil palm and rubber plantations in Sumatra, Indonesia. *PLoS One* 10 (7). <https://doi.org/10.1371/journal.pone.0133325>.
- Bagley, J.E., Desai, A.R., Harding, K.J., Snyder, P.K., Foley, J.A., 2014. Drought and deforestation: Has land cover change influenced recent precipitation extremes in the Amazon? *J. Clim.* 27 (1), 345–361. <https://doi.org/10.1175/JCLI-D-12-00369.1>.
- Barnes, A.D., Jochum, M., Mumme, S., Haneda, N.F., Farajallah, A., Widarto, T.H., Brose, U., 2014. Consequences of tropical land use for multitrophic biodiversity and ecosystem functioning. *Nat. Commun.* 5. <https://doi.org/10.1038/ncomms6351>.
- Berry, S.L., Farquhar, G.D., Roderick, M.L., 2005. 12: co-evolution of climate, soil and vegetation. *Encyclop. Hydrol. Sci.* 177–192. <https://doi.org/10.1002/0470848944>.
- Betts, A.K., Desjardins, R., Worth, D., Cerkowniak, D., 2013. Impact of land use change on the diurnal cycle climate of the Canadian Prairies. *J. Geophys. Res. Atmos.* 118 (21), 11–996. <https://doi.org/10.1002/2013JD020717>.
- BPS, 2013. Badan Pusat Statistik, Statistical Office of Indonesia. Last accessed on April 28, 2018. <http://www.bps.go.id/eng/>.
- Bretfeld, M., Ewers, B.E., Hall, J.S., 2018. Plant water use responses along secondary forest succession during the 2015–2016 El Niño drought in Panama. *New Phytol.* 219, 885–899. <https://doi.org/10.1111/nph.15071>.
- Bruijnzeel, L.A., 1990. Hydrology of moist tropical forests and effects of conversion: a state of knowledge review. *Hydrology of Moist Tropical Forests and Effects of Conversion: a State of Knowledge Review*. Faculty of Earth Sciences, Free University of Amsterdam.
- Canty, A., Ripley, B., 2017. Boot: Bootstrap R (S-Plus) Functions. R Package Version 1.3.
- Cermák, J., 1989. Solar equivalent leaf area: an efficient biometrical parameter of individual leaves, trees and stands. *Tree Physiol.* 5 (3), 269–289. Retrieved from. <http://www.ncbi.nlm.nih.gov/pubmed/14972974>.
- Cienciala, E., Kučera, J., Malmer, A., 2000. Tree sap flow and stand transpiration of two *Acacia mangium* plantations in Sabah, Borneo. *J. Hydrol.* 236 (1–2), 109–120. [https://doi.org/10.1016/S0022-1694\(00\)00291-2](https://doi.org/10.1016/S0022-1694(00)00291-2).
- Clough, Y., Krishna, V.V., Corre, M.D., Darras, K., Denmead, L.H., Meijide, A., Moser, S., Musshoff, O., Steinebach, S., Veldkamp, E., Allen, K., Barnes, A., Breidenbach, N., Brose, U., Buchori, D., Daniel, R., Finkeldey, R., Harahap, I., Hertel, D., Holtkamp, A.M., Hörandl, E., Irawan, B., Jaya, I.N.S., Jochum, M., Klarner, B., Knoch, A., Kotowska, M., Krashevskaya, V., Kreft, H., Kurniawan, S., Leuschner, C., Maraun, M.,

- Melati, D.N., Opfermann, N., Perez-Cruzado, C., Prabowo, W.E., Rembold, K., Rizali, A., Rubiana, R., Schneider, D., Tjitrosedirdjo, S.S., Tjoa, A., Tschamtké, T., Scheu, S., 2016. Land-use choices follow profitability at the expense of ecological functions in Indonesian smallholder landscapes. *Nat. Commun.* 7. <https://doi.org/10.1038/ncomms13137>.
- Delzon, S., Loustau, D., 2005. Age-related decline in stand water use: sap flow and transpiration in a pine forest chronosequence. *Agric. For. Meteorol.* 129 (3–4), 105–119. <https://doi.org/10.1016/j.agrformet.2005.01.002>.
- Dierick, D., Hölscher, D., 2009. Species-specific tree water use characteristics in reforestation stands in the Philippines. *Agric. For. Meteorol.* 149 (8), 1317–1326. <https://doi.org/10.1016/j.agrformet.2009.03.003>.
- Dierick, D., Kunert, N., Köhler, M., Schwendenmann, L., Hölscher, D., 2010. Comparison of tree water use characteristics in reforestation and agroforestry stands across the tropics. In: Tschamtké, T., Leuschner, C., Veldkamp, E., Faust, H., Guhardja, E., Bidin, A. (Eds.), *Tropical Rainforests and Agroforests Under Global Change*. Environmental Science and Engineering, Springer, Berlin Heidelberg, Berlin, Heidelberg, pp. 293–308.
- Drescher, J., Rembold, K., Allen, K., Beckschäfer, P., Buchori, D., Clough, Y., Faust, H., Fauzi, A.M., Gunawan, D., Hertel, D., Jaya, N.S., Klamer, B., Klein, C., Knohl, A., Kotowska, M., Krashevskaya, V., Krishna, V., Leuschner, C., Lorenz, W., Meijide, A., Melati, D., Nomura, M., Perez-Cruzado, C., Qaim, M., Siregar, I.Z., Steinebach, S., Tjoa, A., Tschamtké, T., Wick, B., Wiegand, K., Kreft, H., Scheu, S., 2016. Ecological and socio-economic functions across tropical land use systems after rainforest conversion. *Philos. Trans. Biol. Sci.* 371 (1694), 20150275. <https://doi.org/10.1098/rstb.2015.0275>.
- Ehbrecht, M., Schall, P., Juchheim, J., Ammer, C., Seidel, D., 2016. Effective number of layers: a new measure for quantifying three-dimensional stand structure based on sampling with terrestrial LIDAR. *For. Ecol. Manage.* 380, 212–223.
- Ehbrecht, M., Schall, P., Ammer, C., Seidel, D., 2017. Quantifying stand structural complexity and its relationship with forest management, tree species diversity and microclimate. *Agric. For. Meteorol.* 242, 1–9. <https://doi.org/10.1016/j.agrformet.2017.04.012>.
- Ellison, D., Futter, M.N., Bishop, K., 2012. On the forest cover-water yield debate: from demand- to supply-side thinking. *Glob. Chang. Biol.* <https://doi.org/10.1111/j.1365-2486.2011.02589.x>.
- Ellison, D., Morris, C.E., Locatelli, B., Sheil, D., Cohen, J., Mudiaryso, D., Gutierrez, V., van Noordwijk, M., Creed, I.F., Pokorny, J., Gaveau, D., Spracklen, D.V., Barges, Tobella, A., Ilstedt, U., Teuling, A.J., Gebrehiwot, S.G., Sands, D.S., Muys, B., Sullivan, C.A., 2017. Trees, forests and water: cool insights for a hot world. *Glob. Environ. Change Part A* 43, 51–61. <https://doi.org/10.1016/j.gloenvcha.2017.01.002>.
- Euler, M., Schwarze, S., Siregar, H., Qaim, M., 2016. Oil palm expansion among smallholder farmers in Sumatra, Indonesia. *J. Agric. Econ.* 67 (3), 658–676. <https://doi.org/10.1111/1477-9552.12163>.
- Evans, R.D., Ehleringer, J.R., 1993. A break in the nitrogen-cycle in Aridlands - evidence from Delta-N-15 of soils. *Oecologia* 94 (3), 314–317. <https://doi.org/10.1007/BF00317104>.
- Fan, Y., Rouspard, O., Bernoux, M., Le Maire, G., Panferov, O., Kotowska, M.M., Knohl, A., 2015. A sub-canopy structure for simulating oil palm in the Community Land Model (CLM-Palm): phenology, allocation and yield. *Geosci. Model. Dev.* 8 (11), 3785–3800. <https://doi.org/10.5194/gmd-8-3785-2015>.
- FAO, 2014. FAOSTAT Online Statistical Service. Retrieved January 3, 2018, from <http://www.fao.org/faostat/en/#data/QC>.
- Giambelluca, T.W., Mudd, R.G., Liu, W., Ziegler, A.D., Kobayashi, N., Kumagai, T., Miyazawa, Y., Lim, T.K., Huang, M., Fox, J., Yin, S., Mak, S.V., Kasemsap, P., 2016. Evapotranspiration of rubber (*Hevea brasiliensis*) cultivated at two plantation sites in Southeast Asia. *Water Resour. Res.* 52 (2), 660–679. <https://doi.org/10.1002/2015WR017755>.
- Good, S.P., Noone, D., Bowen, G., 2015. Hydrologic connectivity constrains partitioning of global terrestrial water fluxes. *Science* 349 (6244), 175–177. <https://doi.org/10.1126/science.1259311>.
- Gouyon, A., de Foresta, H., Levang, P., 1993. Does “jungle rubber” deserve its name? An analysis of rubber agroforestry systems in southeast Sumatra. *Agrofor. Syst.* 22 (3), 181–206. <https://doi.org/10.1007/BF00705233>.
- Grala-Michalak, J., Kaźmierczak, K., 2011. Discriminant analysis for Kraft's classes of trees. *Biom. Lett.* 48 (1), 67–81.
- Granier, A., 1985. Une nouvelle méthode pour la mesure du flux de seve brute dans le tronc des arbres. *Ann. Des Sci. For.* 42 (2), 193–200. <https://doi.org/10.1051/forest:19850204>.
- Granier, A., Biron, P., Bréda, N., Pontailier, J.-Y., Saugier, B., 1996. Transpiration of trees and forest stands: short and long-term monitoring using sapflow methods. *Glob. Change Biol.* 2 (3), 265–274. <https://doi.org/10.1111/j.1365-2486.1996.tb00078.x>.
- Guillaume, T., Damris, M., Kuzyakov, Y., 2015. Losses of soil carbon by converting tropical forest to plantations: erosion and decomposition estimated by $\delta^{13}\text{C}$. *Glob. Change Biol.* 21 (9), 3548–3560. <https://doi.org/10.1111/gcb.12907>.
- Guillaume, T., Holtkamp, A.M., Damris, M., Brümmer, B., Kuzyakov, Y., 2016. Soil degradation in oil palm and rubber plantations under land resource scarcity. *Agrofor. Ecosyst. Environ.* 232, 110–118.
- Guillaume, T., Kotowska, M.M., Hertel, D., Knohl, A., Krashevskaya, V., Murtlaksono, K., Scheu, S., Kuzyakov, Y., 2018. Carbon costs and benefits of Indonesian rainforest conversion to plantations. *Nat. Commun.* 9 (1), 2388. <https://doi.org/10.1038/s41467-018-04755-y>.
- Guo, D., Westra, S., 2016. Evapotranspiration: Modelling Actual, Potential and Reference Crop Evapotranspiration. R Package Version 1.9. Retrieved from <http://cran.r-project.org/package=Evapotranspiration>.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G., 2013. High-resolution global maps of 21st-century forest cover change. *Science* 342 (6160), 850–853. <https://doi.org/10.1126/science.1244693>.
- Hardanto, A., Röhl, A., Niu, F., Meijide, A., Hendrayanto, Hölscher, D., 2017a. Oil palm and rubber tree water use patterns: effects of topography and flooding. *Front. Plant Sci.* 8. <https://doi.org/10.3389/fpls.2017.00452>.
- Hardanto, A., Röhl, A., Hendrayanto, Hölscher, D., 2017b. Tree soil water uptake and transpiration in mono-cultural and jungle rubber stands of Sumatra. *For. Ecol. Manage.* 397, 67–77. <https://doi.org/10.1016/j.foreco.2017.04.032>.
- Hatton, T.J., Moore, S.J., Reece, P.H., 1995. Estimating stand transpiration in a eucalyptus-populnea woodland with the heat pulse method - measurement errors and sampling strategies. *Tree Physiol.* 15 (4), 219–227. <https://doi.org/10.1093/treephys/15.4.219>.
- Held, I.M., Soden, B.J.W., 2000. Water vapor feedback and global warming. *Annu. Rev. Environ. Environ.* 25, 441–475. <https://doi.org/10.1146/annurev.energy.25.1.441>.
- Inamdar, A.K., Ramanathan, V., 1998. Tropical and global scale interactions among water vapor, atmospheric greenhouse effect, and surface temperature. *J. Geophys. Res.* Atmos. 103 (D24), 32177–32194. <https://doi.org/10.1029/1998JD900007>.
- Jasechko, S., Sharp, Z.D., Gibson, J.J., Birks, S.J., Yi, Y., Fawcett, P.J., 2013. Terrestrial water fluxes dominated by transpiration. *Nature* 496 (7445), 347–350. <https://doi.org/10.1038/nature11983>.
- Jayasuriya, M.D.A., Dunn, G., Benyon, R., O'Shaughnessy, P.J., 1993. Some factors affecting water yield from mountain ash (*Eucalyptus regnans*) dominated forests in south-east Australia. *J. Hydrol. (Amst)* 150 (2–4), 345–367. [https://doi.org/10.1016/0022-1694\(93\)90116-Q](https://doi.org/10.1016/0022-1694(93)90116-Q).
- Joshi, L., Wibawa, G., Beukema, H., Williams, S., van Noordwijk, M., 2003. Technological change and biodiversity in the rubber agroecosystem of Sumatra. In: Vandermeer, J. (Ed.), *Tropical Agroecosystems*. CRC Press, FL, USA, pp. 133–157.
- Juhrbato, J., Leuschner, C., Hölscher, D., 2004. The relationship between maximal stomatal conductance and leaf traits in eight Southeast Asian early successional tree species. *For. Ecol. Manage.* 202 (1–3), 245–256. <https://doi.org/10.1016/j.foreco.2004.07.021>.
- Kobayashi, N., Kumagai, T., Miyazawa, Y., Matsumoto, K., Tateishi, M., Lim, T.K., Mudd, R.G., Ziegler, A.D., Giambelluca, T.W., Yin, S., 2014. Transpiration characteristics of a rubber plantation in central Cambodia. *Tree Physiol.* 34 (3), 285–301. <https://doi.org/10.1093/treephys/tpu009>.
- Köhler, M., Hanf, A., Barus, H., Hölscher, D., 2014. Cacao trees under different shade tree shelter: effects on water use. *Agrofor. Syst.* 88 (1), 63–73. <https://doi.org/10.1007/s10457-013-9656-3>.
- Kotowska, M.M., Leuschner, C., Triadiati, T., Meriem, S., Hertel, D., 2015. Quantifying above- and belowground biomass carbon loss with forest conversion in tropical lowlands of Sumatra (Indonesia). *Glob. Change Biol.* 21 (10), 3620–3634. <https://doi.org/10.1111/gcb.12979>.
- Kraft, G., 1984. Zur Lehre von der Durchforstungen, Schlagesellungen und Lichtungschieben, as seen in Assmann, E. (1961) *Waldertragskunde*. BLV, München-Bonn-Wien, Hannover, München-Bonn-Wien, pp. 83.
- Kreyling, J., Schweiger, A.H., Bahn, M., Ineson, P., Migliavacca, M., Morel-Journel, T., Christiansen, J.R., Schtickzelle, N., Larsen, K.S., 2018. To replicate, or not to replicate—that is the question: how to tackle nonlinear responses in ecological experiments. *Ecol. Lett.* 2018 (August). <https://doi.org/10.1111/ele.13134>.
- Kubitzka, C., Krishna, V.V., Alamsyah, Z., Qaim, M., 2018. The economics behind an ecological crisis: livelihood effects of oil palm expansion in Sumatra, Indonesia. *Hum. Ecol.* 46 (1), 107–116 2018.
- Kume, T., Takizawa, H., Yoshifuji, N., Tanaka, K., Tantasirin, C., Tanaka, N., Suzuki, M., 2007. Impact of soil drought on sap flow and water status of evergreen trees in a tropical monsoon forest in northern Thailand. *For. Ecol. Manage.* 238 (1–3), 220–230. <https://doi.org/10.1016/j.foreco.2006.10.019>.
- Kume, T., Tanaka, N., Kuraji, K., Komatsu, H., Yoshifuji, N., Saitoh, T.M., Suzuki, M., Kumagai, T., 2011. Ten-year evapotranspiration estimates in a Bornean tropical rainforest. *Agric. For. Meteorol.* 151 (9), 1183–1192. <https://doi.org/10.1016/j.agrformet.2011.04.005>.
- Kunert, N., Schwendenmann, L., Hölscher, D., 2010. Seasonal dynamics of tree sap flux and water use in nine species in Panamanian forest plantations. *Agric. For. Meteorol.* 150 (3), 411–419. <https://doi.org/10.1016/j.agrformet.2010.01.006>.
- Kunert, N., Aparecido, L.M.T., Wolff, S., Higuchi, N., Santos, J., dos, Araújo, de, A.C., Trumbore, S., 2017. A revised hydrological model for the Central Amazon: The importance of emergent canopy trees in the forest water budget. *Agric. For. Meteorol.* 239, 47–57. <https://doi.org/10.1016/j.agrformet.2017.03.002>.
- Lion, M., Kosugi, Y., Takanashi, S., Noguchi, S., Itoh, M., Katsuyama, M., Matsuo, N., Shamsuddin, S.A., 2017. Evapotranspiration and water source of a tropical rainforest in peninsular Malaysia. *Hydrol. Process.* 31 (24), 4338–4353. <https://doi.org/10.1002/hyp.11360>.
- Lu, P., Urban, L., Zhao, P., 2004. Granier's Thermal Dissipation Probe (TDP) method for measuring sap flow in trees: theory and practice. *Acta Bot. Sin.* 46 (6), 631–646. <https://doi.org/10.3389/fpls.2017.00452>.
- Manoli, G., Meijide, A., Huth, N., Knohl, A., Kosugi, Y., Burlando, P., Ghazoul, J., Fatichi, S., 2018. Ecohydrological changes after tropical forest conversion to oil palm. *Environ. Res. Lett.* 13 (6), 064035.
- Margono, B.A., Turubanova, S., Zhuravleva, I., Potapov, P., Tyukavina, A., Baccini, A., Goetz, S., Hansen, M.C., 2012. Mapping and monitoring deforestation and forest degradation in Sumatra (Indonesia) using Landsat time series data sets from 1990 to 2010. *Environ. Res. Lett.* 7 (3), 034010. <https://doi.org/10.1088/1748-9326/7/3/034010>.
- Margono, B.A., Potapov, P.V., Turubanova, S., Stolle, F., Hansen, M.C., 2014. Primary forest cover loss in Indonesia over 2000–2012. *Nat. Clim. Chang.* <https://doi.org/10.1038/nclimate2111>.

- 1038/ncimate2277.
- Martinelli, L.A., Piccolo, M.C., Townsend, A.R., Vitousek, P.M., Cuevas, E., McDowell, W., Robertson, G.P., Santos, O.C., Treseder, K., 1999. Nitrogen stable isotopic composition of leaves and soil: tropical versus temperate forests. *Biogeochemistry* 46 (1–3), 45–65. <https://doi.org/10.1007/BF01007573>.
- Matson, P.A., Vitousek, P.M., Ewel, J.J., Mazzarino, M.J., Robertson, G.P., 1987. Nitrogen transformations following tropical forest felling and burning on a volcanic soil. *Ecology* 68 (3), 491–502. <https://doi.org/10.2307/1938454>.
- Medhurst, J.L., Beadle, C.L., 2002. Sapwood hydraulic conductivity and leaf area - Sapwood area relationships following thinning of a Eucalyptus nitens plantation. *Plant Cell Environ.* 25 (8), 1011–1019. <https://doi.org/10.1046/j.1365-3040.2002.00880.x>.
- Meijide, A., Röhl, A., Fan, Y., Herbst, M., Niu, F., Tiedemann, F., June, T., Rauf, A., Hölscher, D., Knohl, A., 2017. Controls of water and energy fluxes in oil palm plantations: environmental variables and oil palm age. *Agric. For. Meteorol.* 239, 71–85. <https://doi.org/10.1016/j.agrformet.2017.02.034>.
- Meijide, A., Badu, C.S., Moyano, F., Tiralla, N., Gunawan, D., Knohl, A., 2018. Impact of forest conversion to oil palm and rubber plantations on microclimate and the role of the 2015 ENSO event. *Agric. For. Meteorol.* 252, 208–219.
- Meinzer, F.C., Bond, B.J., Warren, J.M., Woodruff, D.R., 2005. Does water transport scale universally with tree size? *Funct. Ecol.* 19 (4), 558–565. <https://doi.org/10.1111/j.1365-2435.2005.01017.x>.
- Merten, J., Röhl, A., Guillaume, T., Meijide, A., Tarigan, S., Agustá, H., Dieslich, C., Dittich, C., Faust, H., Gunawan, D., Hein, J., Hendrayanto, Knohl, A., Kuzakov, Y., Wiegand, K., Hölscher, D., 2016. Water scarcity and oil palm expansion: social views and environmental processes. *Ecol. Soc.* 21 (2). <https://doi.org/10.5751/ES-08214-210205>.
- Mueller, N.D., Butler, E.E., McKinnon, K.A., Rhines, A., Tingley, M., Holbrook, N.M., Huybers, P., 2016. Cooling of US Midwest summer temperature extremes from cropland intensification. *Nat. Clim. Change* 6 (3), 317. <https://doi.org/10.1038/ncclimate2825>.
- Mueller, N.D., Rhines, A., Butler, E.E., Ray, D.K., Siebert, S., Holbrook, N.M., Huybers, P., 2017. Global relationships between cropland intensification and summer temperature extremes over the last 50 years. *J. Clim.* 30 (18), 7505–7528. <https://doi.org/10.1175/JCLI-D-17-0096.1>.
- Nadezhda, N., Vandegehuchte, M.W., Steppe, K., 2012. Sap flux density measurements based on the heat field deformation method. *Trees Struct. Funct.* 26 (5), 1439–1448. <https://doi.org/10.1007/s00468-012-0718-3>.
- Niu, F., Röhl, A., Hardanto, A., Meijide, A., Köhler, M., Hendrayanto, Hölscher, D., 2015. Oil palm water use: calibration of a sap flux method and a field measurement scheme. *Tree Physiol.* 35 (5), 563–573. <https://doi.org/10.1093/treephys/tpv013>.
- Niu, F., Röhl, A., Meijide, A., Hendrayanto, Hölscher, D., 2017. Rubber tree transpiration in the lowlands of Sumatra. *Ecophysiology* 10 (7). <https://doi.org/10.1002/eco.1882>.
- O'Brien, J.J., Oberbauer, S.F., Clark, D.B., 2004. Whole tree xylem sap flow responses to multiple environmental variables in a wet tropical forest. *Plant Cell Environ.* 27 (5), 551–567. <https://doi.org/10.1111/j.1365-3040.2003.01160.x>.
- Oren, R., Phillips, N., Ewers, B.E., Pataki, D.E., Mezonigal, J.P., 1999. Sap-flux-scaled transpiration responses to light, vapor pressure deficit, and leaf area reduction in a flooded *Taxodium distichum* forest. *Tree Physiol.* 19 (6), 337–347. <https://doi.org/10.1093/treephys/19.6.337>.
- Priestley, C.H.B., Taylor, R.J., 1972. On the assessment of surface heat flux and evaporation using large-scale parameters. *Mon. Weather. Rev.* 100 (February), 81–92. [https://doi.org/10.1175/1520-0493\(1972\)100<0081:OTAOSH>2.3.CO;2](https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2).
- R Core Team, 2014. R: a Language and Environment for Statistical Computing. Retrieved from R Foundation for Statistical Computing, Vienna, Austria. <http://www.r-project.org/>.
- Radersma, S., de Ridder, N., 1996. Computed evapotranspiration of annual and perennial crops at different temporal and spatial scales using published parameter values. *Agric. Water Manage.* 31, 17–34. [https://doi.org/10.1016/0378-3774\(95\)00123-4](https://doi.org/10.1016/0378-3774(95)00123-4).
- Radersma, S., Ong, C.K., Coe, R., 2006. Water use of tree lines: importance of leaf area and micrometeorology in sub-humid Kenya. *Agrofor. Syst.* 66 (3), 179–189. <https://doi.org/10.1007/s10457-005-6643-3>.
- Roberts, S., Vertessy, R., Grayson, R., 2001. Transpiration from *Eucalyptus sieberi* (L. Johnson) forests of different age. *For. Ecol. Manage.* 143, 153–161. [https://doi.org/10.1016/S0378-1127\(00\)00514-4](https://doi.org/10.1016/S0378-1127(00)00514-4). special issue: THE SCIENCE OF MANAGING FORESTS TO SUSTAIN.
- Röhl, A., Niu, F., Meijide, A., Hardanto, A., Hendrayanto, Knohl, A., Hölscher, D., 2015. Transpiration in an oil palm landscape: effects of palm age. *Biogeosciences* 12 (19), 5619–5633. <https://doi.org/10.5194/bg-12-5619-2015>.
- Sabaja, C.R., Le Maire, G., June, T., Meijide, A., Rouspard, O., Knohl, A., 2017. Expansion of oil palm and other cash crops causes an increase of the land surface temperature in the Jambi province in Indonesia. *Biogeosciences* 14 (20), 4619–4635. <https://doi.org/10.5194/bg-14-4619-2017>.
- Sampaio, G., Nobre, C., Costa, M.H., Satyamurty, P., Soares-Filho, B.S., Cardoso, M., 2007. Regional climate change over eastern Amazonia caused by pasture and soybean cropland expansion. *Geophys. Res. Lett.* 34 (17). <https://doi.org/10.1029/2007GL030612>.
- Silvério, D.V., Brando, P.M., Macedo, M.N., Beck, P.S.A., Bustamante, M., Coe, M.T., 2015. Agricultural expansion dominates climate changes in southeastern Amazonia: the overlooked non-GHG forcing. *Environ. Res. Lett.* 10 (10). <https://doi.org/10.1088/1748-9326/10/10/104015>.
- Spracklen, D.V., Garcia-Carreras, L., 2015. The impact of Amazonian deforestation on Amazon basin rainfall. *Geophys. Res. Lett.* 42 (21), 9546–9552. <https://doi.org/10.1002/2015GL066063>.
- Tan, Z.H., Zhang, Y.P., Song, Q.H., Liu, W.J., Deng, X.B., Tang, J.W., Deng, Y., Zhou, W.J., Yang, L.Y., Yu, G.R., Sun, X.M., Liang, N.S., 2011. Rubber plantations act as water pumps in tropical China. *Geophys. Res. Lett.* 38 (24). <https://doi.org/10.1029/2011GL050006>.
- Tarigan, S., Wiegand, K., Sunarti, Slamet, B., 2018. Minimum forest cover required for sustainable water flow regulation of a watershed: a case study in Jambi Province, Indonesia. *Hydro. Earth Syst. Sci.* 22 (1), 581–594. <https://doi.org/10.5194/hess-22-581-2018>.
- Teuling, A.J., 2018. A forest evapotranspiration paradox investigated using lysimeter data. *Vadose Zone J.* 17 (1), 170031. <https://doi.org/10.2136/vzj2017.01.0031>.
- Teuscher, M., Gérard, A., Brose, U., Buchori, D., Clough, Y., Ehbrecht, M., Hölscher, D., Irawan, B., Sundawati, L., Wolini, M., Kreft, H., 2016. Experimental biodiversity enrichment in oil-palm-Dominated landscapes in Indonesia. *Front. Plant Sci.* 07. <https://doi.org/10.3389/fpls.2016.01538>.
- Lumley, T., Miller, A., 2009. Leaps: Regression Subset Selection. R Package Version 2.9. <http://CRAN.R-project.org/package=leaps>.
- Tromp-van Meerveld, H.J., McDonnell, J.J., 2006. On the interrelations between topography, soil depth, soil moisture, transpiration rates and species distribution at the hillslope scale. *Adv. Water Resour.* 29 (2), 293–310. <https://doi.org/10.1016/j.advwatres.2005.02.016>.
- Van Noordwijk, M., Khasanah, N., Dewi, S., 2017. Can intensification reduce emission intensity of biofuel through optimized fertilizer use? Theory and the case of oil palm in Indonesia. *Gcb Bioenergy* 9 (5), 940–952. <https://doi.org/10.1111/gcbb.12398>.
- Vertessy, R.A., Watson, F.G., Sharon, K.O., 2001. Factors determining relations between stand age and catchment water balance in mountain ash forests. *For. Ecol. Manage.* 143 (1–3), 13–26. [https://doi.org/10.1016/S0378-1127\(00\)00501-6](https://doi.org/10.1016/S0378-1127(00)00501-6).
- Villamor, G.B., Pontius, R.G., Van Noordwijk, M., 2014. Agroforest's growing role in reducing carbon losses from Jambi (Sumatra), Indonesia. *Region. Environ. Change* 14 (2), 825–834. <https://doi.org/10.1007/s10113-013-0525-4>.
- Woittiez, L.S., Turhina, S., Decsy, D., Slingerland, M., Van Noordwijk, M., Giller, K.E., 2018. Fertiliser application practices and nutrient deficiencies in smallholder oil palm plantations in Indonesia. *Exp. Agric.* <https://doi.org/10.1017/S0014479718000182>.
- Zemp, D.C., Schleussner, C.F., Barbosa, H.M.J., Rammig, A., 2017a. Deforestation effects on Amazon forest resilience. *Geophys. Res. Lett.* 44 (12), 6182–6190. <https://doi.org/10.1002/2017GL072955>.
- Zemp, D.C., Schleussner, C.F., Barbosa, H.M.J., Hirota, M., Montade, V., Sampaio, G., Staal, A., Wang-Erlandsson, L., Rammig, A., 2017b. Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks. *Nat. Commun.* 8. <https://doi.org/10.1038/ncomms14681>.

Transpiration on the rebound in lowland Sumatra

ORIGINALITY REPORT

0%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

Exclude quotes On

Exclude bibliography Off

Exclude assignment
template On

Exclude matches Off